

Metabolites produced by antagonistic microbes inhibit the principal avocado pathogens *in vitro*

Metabolitos producidos por microorganismos antagonistas son capaces de inhibir *in vitro* los principales patógenos del aguacate

Sara Ramírez R.¹, Julián David Arias M.², Juan Carlos Bedoya³, Ever Antoni Rueda L.⁴,
Claudia Yaneth Sánchez¹, and Sinar David Granada G.²

ABSTRACT

The demand for Hass avocado in the global market exceeds the supply by over 50%. Colombia has a remarkable advantage as a producer in the region due to its high yields. However, the productivity of this crop can be seriously affected by diseases such as root rot, caused by *Phytophthora cinnamomi*, postharvest body rot and stem end rot, caused by *Colletotrichum* sp. and *Phomopsis* sp., respectively. The potential of 76 bacterial isolates obtained from avocado rhizosphere to produce inhibitory metabolites against avocado's pathogens was evaluated. The antagonistic effect of the rhizobacteria against *P. cinnamomi*, *Colletotrichum* sp. and *Phomopsis* sp. was tested through dual cultures. Thirty-six percent of the tested isolates presented inhibition halos against *P. cinnamomi*, 36% against *Colletotrichum* sp. and 67% against *Phomopsis* sp. Additionally, three isolates were selected for fermentation tests using different broth cultures. The extracts obtained from fermentations in the minimal medium of isolates ARP5.1 and AED06 showed inhibitory activity against the evaluated pathogens, but this effect was not observed with the AED26 extract. The media supplemented with copper chloride did not enhance activity of the extracts. These results suggest that using microbial metabolic extracts is a viable alternative for controlling avocado pathogens *in vitro*.

Key words: *Persea americana*, postharvest diseases, *Phytophthora cinnamomi*, secondary metabolites, rhizobacteria.

RESUMEN

La demanda mundial de aguacate Hass supera la oferta en más de un 50%. A pesar de los altos rendimientos de este cultivo en Colombia, la productividad se ve seriamente afectada por enfermedades como la pudrición de la raíz, causada por *Phytophthora cinnamomi*, y las pudriciones poscosecha causadas por *Colletotrichum* sp. y *Phomopsis* sp. En este trabajo se evaluó el potencial de 76 aislamientos bacterianos obtenidos de la rizósfera de plantas de aguacate, para la producción de metabolitos con actividad inhibitoria frente a patógenos del cultivo. El efecto antagonista de las rizobacterias frente a *P. cinnamomi*, *Colletotrichum* sp. y *Phomopsis* sp. fue probado a través de cultivos duales. El 36% de los aislamientos evaluados mostraron halos de inhibición frente a *P. cinnamomi*, 36% frente a *Colletotrichum* sp. y 67% frente a *Phomopsis* sp. Tres de los aislamientos más promisorios se fermentaron en diferentes medios de cultivo líquidos. Los extractos de los aislamientos ARP5.1 y AED06 a partir de medio mínimo mostraron actividad inhibitoria frente a los patógenos. Los medios suplementados con cloruro de cobre no incrementaron la actividad de los extractos. Estos resultados demostraron la viabilidad de utilizar extractos metabólicos microbianos como una alternativa para controlar *in vitro* los patógenos de aguacate.

Palabras clave: *Persea americana*, enfermedades poscosecha, *Phytophthora cinnamomi*, metabolitos secundarios, rizobacterias.

Introduction

The global production of avocado exceeds 4 million t year⁻¹. In terms of the volume of the fruit and its acreage, avocado holds the fifth largest tropical fruit crop in the world. In 2010, Colombia was ranked fifth in avocado production worldwide (Yabrudy, 2012). Due to its high nutritional value, palatability, versatility, ease of preparation, wide

usability in culinary, pharmaceutical and cosmetic industries, this fruit is considered a product with great export potential for Colombia (Alfonso, 2008).

Despite its high production volume, Colombia still relies on imports to meet the domestic demand, which demonstrates the need for increasing avocado production (Alfonso, 2008; Yabrudy, 2012). Moreover, avocado crops are severely

Received for publication: 14 January, 2015. Accepted for publication: 30 March, 2015.

Doi: 10.15446/agron.colomb.v33n1.48241

¹ School of Chemistry, Politécnico Colombiano Jaime Isaza Cadavid. Medellín (Colombia)

² Plant Health and Biological Control Unit, Corporación para Investigaciones Biológicas. Medellín (Colombia). davidgranada888@gmail.com

³ Faculty of Health Sciences, Institución Universitaria Colegio Mayor de Antioquia. Medellín (Colombia)

⁴ Faculty of Sciences, Universidad del Tolima. Ibagué (Colombia)

affected by diseases such as root rot or dieback, stem-end rot and body rot, caused by *P. cinnamomi*, *Phomopsis* sp. and *Colletotrichum gloeosporioides*, respectively. The latter is responsible for the loss of up to 80% of the harvest in some crops (Demoz and Korsten, 2006; Jianga *et al.*, 2014).

Usually, disease control in avocado productive systems depends on agrochemical applications that have gradually caused important environmental damage and affected human health, turning plant health problems into a chronic threat (Morales, 2009; Romero-Correa *et al.*, 2014). Therefore, the use of compounds from nature, particularly the use of active compounds derived from plant extracts, has been positioning itself as an alternative to overcome these drawbacks (Grahovac *et al.*, 2014). However, the availability of plant material is generally low, making it difficult to reach large scale production and increasing manufacture costs (Subramani and Aalbersberg, 2012; Zabka and Pavela, 2013).

In recent years, the use of microorganisms as a source of bioactive substances has captured the attention of researchers (Bertrand *et al.*, 2014). Microbes have great potential for biological activities and the use of large-scale fermentation processes can reduce production costs of microbe-derived compounds. Antagonists are among plant associated microbes with a big potential to produce bioactive metabolites that can inhibit, partially or totally, the growth of pathogens (Van Eeden and Korsten, 2013). Some of them have been used in biological control systems with very good outcomes (Van Eeden and Korsten, 2013).

Therefore, finding an economically and environmentally viable alternative for obtaining compounds that are useful for the control of plant pathogens is urgent. Thus, the aim of this study was to assess the *in vitro* capability of antagonistic rhizobacteria associated with avocado crops of controlling and producing bioactive extracellular metabolites against *P. cinnamomi*, *Phomopsis* sp. and *Colletotrichum* sp.

Materials and methods

Culture media and reagents

Trypticase soy agar and broth (TSA and TSB), and Potato-Dextrose Agar (PDA) were purchased from Merck. Sabouraud broth was purchased from BD. Minimal medium (MM) components (200 mL M9 salts: 64.0 g L⁻¹ Na₂HPO₂·7H₂O, 15.0 g L⁻¹ KH₂PO₄, 2.5 g L⁻¹ NaCl, 5.0 g L⁻¹ NH₄Cl; 2.0 mL 1 M MgSO₄; 25.0 mL glucose 20%; 0.1 mL 1 M CaCl₂; 773 mL water). The solvents were purchased from Merck as well.

Microorganisms

The pathogen strains (*Phytophthora cinnamomi*, *Colletotrichum* sp. and *Phomopsis* sp.) were taken from the plant pathogen bank of Corporación para Investigaciones Biológicas, CIB. Samples for obtaining antagonistic microbes were taken from avocado producing farms located in Anserma (Caldas, Colombia) and in eastern Antioquia (Colombia). Nutritional roots and soil samples were isolated from apparently healthy avocado plants. Five to eight trees were sampled from each farm in a zigzag pattern. The samples were carried in properly labeled plastic bags and stored at 4°C until processed in the laboratory. The roots were washed with sterile distilled water and cut into 0.5 cm pieces. For the endophyte isolation, the roots were washed with distilled water, disinfected with 1% sodium hypochlorite for 1 min and cut into 0.5 cm pieces. The fragments were placed on TSA and incubated at 21°C for 24 to 48 h. After incubation, the single bacterial colonies grown on the surface of the culture medium were sub-cultured in a fresh medium. Each isolate was coded as “AED” in the case of endophytes and as “ARP” in the case of rhizoplane microbes. The purity of the isolates was checked with staining and microscopy techniques.

Antagonism assays

The selection of antagonistic isolates was performed using dual culture assays in petri dishes (Granada *et al.*, 2014). Briefly, petri dishes containing PDA were divided into four quadrants and each one was inoculated at 1 cm from the edge of the plate with a loopful of four different potential antagonists. A plug of the pathogen to be tested, grown during 8 d on PDA, was placed in the middle of the plate. A growth control consisting of the pathogen without any bacteria was performed as well. The inhibition zone around each potential antagonist was measured after 6 d of incubation at 24°C. The degree of inhibition was quantified on a scale from 0 to 3 (0 = no inhibition halo, 1 = halo from 0.1 to 1.0 cm, 2 = halo from 1.0 to 2.0 cm, 3 = halo from 2.0 to 4.0 cm). All procedures were conducted in triplicate with a repetition over time.

Fermentations and extracellular metabolite extraction

Pre-inoculum of the bacteria to be fermented were carried out in TSB and incubated for 24 h at 29°C and 200 rpm. Subsequently, 5 mL of the pre-inoculum were added to 125 mL erlenmeyer flasks containing 45 mL of three different media (minimal medium (MM), MM supplemented with copper chloride (1mM CuCl₂), TSB and TSB supplemented with copper chloride (1mM CuCl₂). The incubation conditions were 200 rpm at 29°C for 96 h. After fermentation,

the biomass was separated through centrifugation at 8,000 rpm for 5 min, the supernatants were freeze-dried and a solid-liquid extraction was performed using 20 mL of methanol. Methanolic solutions were dried in a vacuum assisted rotary evaporator.

Antimicrobial activity assays of metabolic extracts

Antimicrobial activity assays against *Colletotrichum* sp., *P. cinnamomi* and *Phomopsis* sp. were performed using 96-well microplates (Sanmartín-Negredo *et al.*, 2012; Villamil *et al.*, 2012). A BioRad microplate reader, model 680 XR (BioRad laboratories, Tokyo, Japan), at a 595 nm wavelength, was used. The *Colletotrichum* sp. inoculum was prepared using agar plugs in sterile distilled water to reach a concentration of $1 \cdot 10^4$ conidia/mL. The *P. cinnamomi* and *Phomopsis* sp. inoculums were prepared by mycelia fragmentation using sterile glass pearls to reach concentrations of $1 \cdot 10^4$ fragments/mL. Counting was achieved using a hemocytometer. Each plate well had the following added: 50 μ L of the pathogen, 100 μ L of the Sabouraud broth and 50 μ L of the microbial extract re-suspended in sterile distilled water to reach a concentration of 1% w/v. Controls with water instead of the extracts were performed and all of the treatments were carried out in quadruplicate. The microplates were read after 5 d of incubation at 24°C. The data were analyzed using ANOVA with Tukey post-hoc tests with VassarStats online software (available in <http://vassarstats.net/anova1u.html>).

Results and discussion

Antagonism assays

From the processed samples, 76 bacterial isolates were obtained. All of the pathogens tested showed inhibition halos at different degrees after being antagonized with the 76 bacterial isolates (Fig. 1). The 37% showed inhibition halos against *P. cinnamomi*, 36% against *Colletotrichum* sp. and 67% against *Phomopsis* sp. In the case of *P. cinnamomi*, 14.5% of isolates were ranked as degree three (3), 10.5% for *Colletotrichum* sp. and 42% *Phomopsis* sp. (Tab. 1), which indicates an increased susceptibility of *Phomopsis* sp. to be inhibited *in vitro*. It should be noted that several of the isolates showed inhibition zones against more than one pathogen.

TABLE 1. Summary of antagonism assay results of the 76 bacterial isolates.

Pathogen	<i>Phytophthora cinnamomi</i>				<i>Colletotrichum</i> sp.				<i>Phomopsis</i> sp.			
Inhibition degree	0	1	2	3	0	1	2	3	0	1	2	3
Number of isolates	48	10	6	12	51	13	4	8	24	6	16	30

Inhibition degree: 0, no inhibition halo; 1, halo from 0.1 to 1.0 cm; 2, halo from 1.0 to 2.0 cm; 3, halo from 2.0 to 4.0 cm.




FIGURE 1. Dual culture antagonism assays (three replicates per set of isolates). A, control of *P. cinnamomi*; B, dual culture assay for *P. cinnamomi* against four different bacterial isolates; C, control of *Colletotrichum* sp.; D, dual culture assay for *Colletotrichum* sp. against four different bacterial isolates; E, control of *Phomopsis* sp.; F, dual culture assay for *Phomopsis* sp. against four different bacterial isolates.

Antimicrobial assays of extracts from fermentations

Isolates AED06, AED26 and ARP5.1 showed the greatest inhibition degree and the broadest spectrum against the tested pathogens. Due to this, they were selected to be grown in broth medium in erlenmeyer flasks.

From the four tested media, the minimal medium (45 mL) was the most appropriate for producing inhibitory metabolites against pathogens used in this study. Although TSB showed a high biomass development, the extracts did not show significant inhibitory activity against pathogens. This may be due to the association of limitation of essential nutrients with the production of secondary metabolites. The depletion of nutrients generates stress responses in microbes, such as motility, secretion of extracellular enzymes, competence for genetic transformation, production of antibiotics or other bioactive compounds (Granada *et al.*, 2014; Thines *et al.*, 2014). Additionally, the media supplemented with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ did not show significant activity or diminished biomass production (Carvalho *et al.*, 2010). Nevertheless, several authors have reported a boost in antibiotic production and a positive regulation of secondary metabolism of various types of microorganisms when heavy metals such as copper are added to culture media (Paul and Banerjee, 1983; Tunac and Mcdaniel, 1985; Jafarzade *et al.*, 2013).

Conversely, copper is known to participate in the degradation of nucleic acids through destabilization and oxidative or hydrolytic breakdown of the double helix, possibly explaining the inhibition observed in the supplemented fermentations (Cervantes and Gutiérrez-Corona, 1994; Weinberg, 1997). However, copper's toxicity depends on the environmental conditions, concentration and type of microbes in the environment (Lippert, 1992).

Extract from the ARP5.1 isolate showed the highest inhibition activity against all of the tested pathogens, which was evidenced by the low optical density. Lower optical densities involve a small development of biomass; *i.e.* inhibition or less growth of the evaluated microbes. The characteristics of the limiting conditions of salts and glucose in the minimal medium probably stimulated the production of molecules involved in the antagonistic mechanisms (Ruyters *et al.*, 2013). The growth of the *P. cinnamomi* treated with the AED26 extract was also significantly lower ($P \leq 0.05$), as compared with the control (Fig. 2), but it did not have an effect on the other tested pathogens. This is a surprising result because this isolate was selected due to its inhibition halos in dual culture tests. This is probably owing to the culture media used

that impairs the production of bioactive metabolites on this isolate. Moreover, it has been reported that some bacteria are capable of producing such compounds in response to the presence of other microorganisms or in response to aspects of the environment (Ryan and Dow, 2008). Therefore, in the absence of competitors, compounds with a biocidal capacity are not produced by the antagonist and further studies must be performed on this type of isolates.




FIGURE 2. Activity assays against *P. cinnamomi* using the minimal medium for production of extracts. Control: *P. cinnamomi* without an extract. Means with different letters indicate significant differences according to the Tukey test ($P \leq 0.05$) ($n = 4$). Bars indicate standard error.

The AED06 extract showed high inhibition against *Colletotrichum* sp. and *Phomopsis* sp., but a low effect on *P. cinnamomi* (Figs. 3 and 4). *P. cinnamomi* is not considered a fungus, but is a pathogen of the Oomycete class; this might imply that metabolites produced by AED06 have a narrow spectrum and are only active against true fungi. Conversely, metabolites produced by ARP5.1 have a broader spectrum and their mode of action might be able to kill the different tested pathogens generally.

This study showed that healthy avocado trees are associated with microbes capable of *in vitro* inhibition against the growth of pathogens such as *P. cinnamomi*, *Colletotrichum* sp. and *Phomopsis* sp., and that one of the mechanisms for inhibition is the production of secondary metabolites. Furthermore, the minimal medium represented the most suitable substrate for the production of inhibitory extracts by the tested bacteria. Additionally, it was observed that the potential of a microbe to produce inhibitory metabolites not only depends on what is evidenced from dual culture assays using petri dishes, but may also possibly be a response to the presence of other competing microbes. Thus, the production of said metabolites in an




FIGURE 3. Activity assays against *Colletotrichum* sp. using the minimal medium for the production of extracts. Control: *Colletotrichum* sp. without an extract. Means with different letters indicate significant differences according to the Tukey test ($P \leq 0.05$) ($n = 4$). Bars indicate standard error.

axenic liquid medium may be limited when there are not any competing microbes.

Acknowledgement

The authors would like to thank the Institución Universitaria Colegio Mayor de Antioquia, especially the Faculty of Health Sciences, for funding this research.

Literature cited

- Alfonso B, J.A. 2008. Manual técnico del cultivo de aguacate Hass (*Persea americana* L.). Fundación Hondureña de Investigación Agrícola (FHIA), La Lima, Honduras.
- Bertrand, S., N. Bohni, S. Schnee, O. Schumpp, K. Gindro, and J.L. Wolfender. 2014. Metabolite induction via microorganism co-culture: a potential way to enhance chemical diversity for drug discovery. *Biotechnol. Adv.* 32, 1180-1204. Doi: 10.1016/j.biotechadv.2014.03.001
- Carvalho, A.L.U., F.H.P.C. Oliveira, R.L.R. Mariano, E.R. Gouveia, and A.M. Souto-Maior. 2010. Growth, sporulation and production of bioactive compounds by *Bacillus subtilis* R14. *Braz. Arch. Biol. Technol.* 53, 643-652. Doi: 10.1590/S1516-89132010000300020
- Cervantes, C. and F. Gutiérrez-Corona. 1994. Copper resistance mechanisms in bacteria and fungi. *FEMS Microbiol. Rev.* 14, 121-137. Doi: 10.1111/j.1574-6976.1994.tb00083.x
- Demoz, B.T. and L. Korsten. 2006. *Bacillus subtilis* attachment, colonization, and survival on avocado flowers and its mode of action on stem-end rot pathogens. *Biol. Control* 37, 68-74. Doi: 10.1016/j.biocontrol.2005.11.010
- Grahovac, J., M. Grahovac, J. Dodić, B. Bajić, and J. Balaž. 2014. Optimization of cultivation medium for enhanced production of antifungal metabolites by *Streptomyces hygroscopicus*. *Crop Prot.* 65, 143-152. Doi: 10.1016/j.cropro.2014.07.020
- Granada G, S.D., A. Rueda L., and C.A. Peláez. 2014. Antimicrobial activity of extracellular metabolites from antagonistic bacteria isolated from potato (*Solanum phureja*) crops. *Summa Phytopathol.* 40, 212-220. Doi: 10.1590/0100-5405/1953




FIGURE 4. Activity assays against *Phomopsis* sp. using the minimal medium for production of extracts. Control: *Phomopsis* sp. without an extract. Means with different letters indicate significant differences according to the Tukey test ($P \leq 0.05$) ($n = 4$). Bars indicate standard error.

- Jafarzade, M., N.A. Yahya, F. Shayesteh, G. Usup, and A. Ahmad. 2013. Influence of culture conditions and medium composition on the production of antibacterial compounds by marine *Serratia* sp. WPR3. *J. Microbiol.* 51, 373-379. Doi: 10.1007/s12275-013-2440-2
- Jianga, J., H. Zhaia, H. Li, Z. Wang, Y. Chen, N. Hong, G. Wang, G.N. Chofong, and W. Xu. 2014. Identification and characterization of *Colletotrichum fructicola* causing black spots on young fruits related to bitter rot of pear (*Pyrus bretschneideri* Rehd.) in China. *Crop Prot.* 58, 41-48. Doi: 10.1016/j.cropro.2014.01.003
- Lippert, B. 1992. From cisplatin to artificial nucleases - the role of metal ion-nucleic acid interactions in biology. *Biometals* 5, 195-208. Doi: 10.1007/BF01061218
- Morales, J. 2009. Enfermedades de importancia económica en el cultivo de aguacate. pp. 15-31. In: Libro de Resúmenes, III Congreso Latinoamericano del Aguacate. Medellín, Colombia.
- Paul, A.K. and A.K. Banerjee. 1983. Determination of optimum conditions for antibiotic production by *Streptomyces galbus*. *Folia Microbiol.* 28, 397-405. Doi: 10.1007/BF02879489
- Romero-Correa, M.T., R. Villa-Gómez, E. Castro-Mercado, and E. García-Pineda. 2014. The avocado defense compound phenol-2,4-bis (1,1-dimethylethyl) is induced by arachidonic acid and acts via the inhibition of hydrogen peroxide production by pathogens. *Physiol. Mol. Plant Pathol.* 87, 32-41. Doi: 10.1016/j.pmp.2014.05.003
- Ruyters, S., P. Salaets, K. Oorts, and E. Smolders. 2013. Copper toxicity in soils under established vineyards in Europe: a survey. *Sci. Total Environ.* 443, 470-477. Doi: 10.1016/j.scitotenv.2012.11.001
- Ryan, R.P. and J.M. Dow. 2008. Diffusible signals and interspecies communication in bacteria. *Microbiology* 154, 1845-1858. Doi: 10.1099/mic.0.2008/017871-0
- Sanmartín-Negredo, P., X. López, M.P. Pemberthy, D. Granada-Sair, and E.A. Rueda-Lorza. 2012. Análisis del modo de acción de la capacidad antagonista de *Trichoderma asperellum* sobre *Colletotrichum gloeosporioides* y *Fusarium* sp. *Rev. Tumbaga* 2, 29-49.
- Subramani, R. and W. Aalbersberg. 2012. Marine actinomycetes: an ongoing source of novel bioactive metabolites. *Microbiol. Res.* 167, 571-580. Doi: 10.1016/j.micres.2012.06.005

- Thines, E., H. Anke, and R.W.S. Weber. 2004. Fungal secondary metabolites as inhibitors of infection-related morphogenesis in phytopathogenic fungi. *Mycol. Res.* 108, 14-25. Doi: 10.1017/S0953756203008943
- Tunac, J.B. and L.E. McDaniel. 1985. Effect of phosphate and copper on the fermentation of hydroheptin. *Appl. Environ. Microb.* 50, 1192-1195.
- Van Eeden, M. and L. Korsten. 2013. Factors determining use of biological disease control measures by the avocado industry in South Africa. *Crop Prot.* 51, 7-13. Doi: 10.1016/j.cropro.2013.03.011
- Villamil C., J.E., J.O. Blanco V., and S.E. Viteri R. 2012. Evaluación *in vitro* de microorganismos nativos por su antagonismo contra *Moniliophthora roreri* Cif. & Par. en cacao (*Theobroma cacao* L.). *Rev. Fac. Nal. Agr. Medellin* 65, 6305-6315.
- Weinberg, E.D. 1997. Mineral element control of microbial secondary metabolism. pp. 289-316. In: Weinberg, E.D. (ed.). *Microorganisms and minerals*. Marcel Dekker, New York, NY.
- Yabrudy V., J. 2012. El aguacate en Colombia: Estudio de caso de los Montes de María, en el Caribe colombiano. *Documentos de Trabajo de Economía Regional* 171, 1-45.
- Zabka, M. and R. Pavela. 2013. Antifungal efficacy of some natural phenolic compounds against significant pathogenic and toxinogenic filamentous fungi. *Chemosphere* 93, 1051-1056. Doi: 10.1016/j.chemosphere.2013.05.076