

***Melanagromyza obtusa* (Diptera: Agromyzidae), a new record for Colombia***Melanagromyza obtusa* (Diptera: Agromyzidae), nuevo registro para ColombiaJavier O. Martínez-Alava<sup>1</sup>, Francisco Serna<sup>1</sup>, and A. Lucía Pérez B.<sup>1</sup>

## ABSTRACT

*Melanagromyza obtusa* (Diptera: Agromyzidae) is recorded for the first time in Colombia. Distribution of this species is mainly Asiatic although it has been recently reported in Florida (The United States of America), the Dominican Republic, Puerto Rico, Haiti, Panama, and Peru. In countries such as India, Indonesia (Java) and Malaysia, it is recognized as an important pest in economic crops of *Cajanus cajan* (Fabaceae) and, to a lesser degree, in other Fabaceae, such as *Flemingia macrophylla*.

**Key words:** taxonomy, geographic distribution, invasive species.

## RESUMEN

*Melanagromyza obtusa* (Diptera: Agromyzidae) se registra por primera vez para Colombia. Esta especie de distribución principalmente asiática presenta registros recientes para Florida (Estados Unidos), República Dominicana, Puerto Rico, Haití, Panamá y Perú. En países como India, Indonesia (Java) y Malasia se conoce como plaga de importancia económica en cultivos de *Cajanus cajan* (Fabaceae) y en menor grado en otras Fabáceas como *Flemingia macrophylla*.

**Palabras clave:** taxonomía, distribución geográfica, plaga introducida.

*Melanagromyza* Hendel, 1920 is distributed worldwide, being one of the more species-rich genus among the family, with more than 380 described species (Boucher, 2010; Koçak and Sasakawa, 2010; Shi and Gaimari, 2015). Approximately 160 species of this genus are recorded from the New World (Shi and Gaimari, 2015). They are borers in roots, stems, trunks, branches, flower petioles, fruits, ovaries or seeds (Korytkowski, 2014; Shi and Gaimari 2015).

Characteristics used to recognize this genus can be found in Shi and Gaimari (2015). *Melanagromyza obtusa* (Fig. 1) is characterized by a green-copper color; ocellar triangle green, well developed, anteriorly limiting with the margin of the lunule, and with its anterior edge blunt and flat; fronto-orbital bristles numerous, with the anterior ones proclinate; arista long, with conspicuous pubescence; wing length 2.4 to 2.8 mm; hypandrium and aedeagal apodeme long; apodeme of the hypandrium very long. Projection at tip of epandrium rounded, with long hairs; aedeagus hyaline; basiphallus with unusual long side arms, which are extending up to the distiphallus; dorsal bridge of basiphallus seems to be missing (Malloch, 1914; Spencer, 1961, 1973; Dempewolf, 2004).

*Melanagromyza obtusa* (Malloch, 1914) is recognized as an important economic pest in crops of *Cajanus cajan* (L.) Huth (Fabaceae) in India, Indonesia (Java) and Malaysia,

and in a lesser degree in other Fabaceae, including *Flemingia macrophylla* (Willd.) Kuntze ex Merr., and *Phaseolus radiatus* L. (Kulkarni, 1966; Spencer, 1973; Ipe, 1974; Singh and Gupta, 1981; Shanower *et al.*, 1998; Sharma *et al.*, 2003; Moudgal *et al.*, 2008; Palacios-Torres *et al.*, 2010; Araúz *et al.*, 2013; Korytkowski, 2014; EPPO, 2016).

Larvae of *M. obtusa* are borer of seeds and pods of *F. macrophylla*. Adults were obtained from the seeds inside pods of *F. macrophylla*, maintained in the Lab of the UNAB museum. Biological material was collected at the Macagual Research Center of the Universidad de la Amazonia (Uniamazonia), in the municipality of Florencia, department of Caqueta, Colombia.

Morphological characteristics of the specimens were studied with the aid of a NIKON SMZ-1 stereomicroscope (Nikon GmbH, Düsseldorf, Germany). For taxonomic identification, keys, descriptions and diagnosis in Spencer (1961, 1963, 1983, 1984), Spencer and Steyskal (1986), Sanabria (1994), Boucher (2010), Korytkowski (2014), Lonsdale (2014), and Shi and Gaimari (2015) were employed; and for the species level identification, Malloch (1914) and Spencer (1961, 1973) were followed.


In order to examine the male genitalia, the methodology in Korytkowski (2014) and Shi and Gaimari (2015) was

Received for publication: 11 Abril, 2016. Accepted for publication: 30 June, 2016.

Doi: 10.15446/agron.colomb.v34n2.56958

<sup>1</sup> Grupo Sistemática de Insectos Agronomía (SIA). Museo Entomológico UNAB, Faculty of Agricultural Sciences, Universidad Nacional de Colombia. Bogota (Colombia). jomartineza@unal.edu.co





**FIGURE 1.** *Melanagromyza obtusa* (Malloch, 1914). A, female, lateral view; B, male phallic complex, lateral view.

employed. In the UNAB museum, voucher specimens were deposited, following the curatorial procedures suggested in Martínez and Serna (2015).

This is the first record of *M. obtusa* for Colombia. The species distribution is known from Asia (China-Taiwan), Flores Island (Indonesia), India (Assam, Bihar, Delhi, Maharashtra, Uttar Pradesh), Java (Indonesia), Malaysia, Sri Lanka, Bangladesh, Burma, Nepal, Pakistan, Philippines, Thailand, Vietnam, New Guinea, and Japan); also it is known in Australia (Queensland), Papua New Guinea, and in recent years records of this species are known from Florida (USA), the Dominican Republic, Puerto Rico, Haiti, Panama, and Peru (Kulkarni, 1966; Spencer, 1973; Ipe, 1974; Singh and Gupta, 1981; Shanower *et al.*, 1998; Spencer, 1999; Sharma *et al.*, 2003; Palacios-Torres *et al.*, 2010; Araúz *et al.*, 2013; Korytkowski, 2014; EPP0, 2016).

**Material examined:** *Melanagromyza obtusa*, 6♀♀ 5♂♂, **COLOMBIA, Caqueta**, Florencia, Macagual Research Center, 1°29'58.70" N and 75°39'46.49" W, 12-Sep-2014, In *Flemingia macrophylla* (Willd.) Kuntze ex Merr. (Fabaceae), J. Martínez [UNAB No. Catal. 1826]; 1♀ 1♂, **Putumayo**, Villagarzon, Vda. San Rafael, Fca. El Escondite, 0°47'41" N 76°35'8" W, 317 m a.s.l., In: *Flemingia macrophylla* (Fabaceae), 29-Feb-2016, J. Martínez [UNAB No. Catal. 1833].

### Acknowledgements

The authors thank the UNAB (Universidad Nacional de Colombia, Bogota) Entomological Museum for providing the necessary tools for the rearing and the identification of specimens, and for the curatorial procedures and Colciencias and the Universidad Nacional de Colombia for sponsoring the first author for his doctorate under the doctoral training program in Colombia.

### Literature cited

- Araúz, N., A. Santos M., R.A. Cambra T., and J.A. Bernal V. 2013. Insectos plagas y parasitoides asociados al cultivo de Guandú (*Cajanus cajan* (L.) millsp. Fabaceae) en Chiriquí, República de Panamá. *Tecnociencia* 15, 5-18.
- Boucher, S. 2010. Family Agromyzidae (leaf-mining flies). pp. 1057–1071. In: Brown, B.V., A. Borkent, J.M. Cumming, D.M. Wood, N.E. Woodley, and M. Zumbado (eds.). *Manual of Central American Diptera*. Vol. 2. NRC Research Press, Ottawa.
- Dempewolf, M. 2004. Arthropods of economic importance - Agromyzidae of the world. In: <http://wbd.etibioinformatics.nl/bis/agromyzidae.php?selected=beschrijving&menuentry=soorten&id=99>; consulted: february, 2016.
- EPP0. 2016. EPP0 global database. In: <https://gd.eppo.int>; consulted: february, 2016.
- Ipe, I.M. 1974. Morphological, behavioural and biological studies of *Melanagromyza obtusa* (Malloch) (Diptera: Agromyzidae) on *Cajanus indicus* Spreng. *Zeitschr. Angew. Entomol.* 75, 89-98.
- Koçak, E. and M. Sasakawa. 2010. Two species of *Melanagromyza* (Diptera: Agromyzidae), with descriptions of immature stages from *Hera cleum* (Apiaceae) and new records from Turkey. *Entomol. News* 121, 262-266.
- Korytkowski, C.A. 2014. Contribución al conocimiento de los Agromyzidae (Diptera: Muscomorpha) en el Perú. *Rev. Peru. Entomol.* 49, 1-106.
- Kulkarni, S.M. 1966. Infestation, sex-ratio and damage by *Melanagromyza obtusa* (Diptera: Agromyzidae) to *Moghania macrophylla* seeds in the field. *Entomol. Exp. Appl.* 9, 323-326.
- Lonsdale, O. 2014. Redefinition and synonymy of genera in the *Ophiomyia* genus group, with the description of *Euhexomyza* new genus (Diptera: Agromyzidae). *Can. Entomol.* 146, 481-513. Doi: 10.4039/tce.2014.2
- Malloch, J.R. 1914. Formosan Agromyzidae. *Annales Musei Nationalis Hungarici* 12, 306-36.
- Martínez A., J.O. and F. Serna. 2015. Managing insect collections. Micropezidae (Diptera: Nerioidea) of the Entomological Museum UNAB. *Agron. Colomb.* 33, 339-347, Doi: 10.15446/agron.colomb.v33n3.52432

- Moudgal, R.K., R.K. Lakra, B. Dahiya, and M.K. Dhillon. 2008. Physico-chemical traits of *Cajanus cajan* (L.) Millsp. Pod wall affecting *Melanagromyza obtusa* (Malloch) damage. *Euphytica* 161, 429-436. Doi: 10.1007/s10681-007-9601-5
- Palacios-Torres, R.E., J.M. Valdez-Carrasco, J. Étienne, R. Vega-Muñoz, J. Marín-Sánchez, A. Castañeda-Vildozola, and C. Nava-Díaz. 2010. Nuevos registros de plantas hospederas y distribución geográfica de *Melanagromyza floris* Spencer, *M. tomaterae* Steyskal, *M. viridis* (Frost) y *Ophiomyia lantanae* (Froggatt) (Diptera: Agromyzidae) en México. *Acta Zool. Mex. New Ser.* 26(1), 59-71.
- Sanabria, A. 1994. Insectos minadores (Diptera: Agromyzidae) de la Sabana de Bogotá (Cundinamarca Colombia). *Rev. Colomb. Entomol.* 20(1), 61-100.
- Shanower, T.G., S.S. Lalt, and V.R. Bhagwat. 1998. Biology and management of *Melanagromyza obtusa* (Malloch) (Diptera: Agromyzidae). *Crop Prot.* 17(3), 249-263.
- Sharma, H. C., G. Pampapathy, and L.J. Reddy. 2003. Wild relatives of pigeonpea as a source of resistance to the pod fly (*Melanagromyza obtusa* Malloch) and pod wasp (*Tanaostigmodes cajaninae* La Salle). *Gen. Res. Crop Evol.* 50, 817-824.
- Shi, L. and S.D. Gaimari. 2015. Revision of the genus *Melanagromyza* in California, with descriptions of three new species (Diptera: Agromyzidae). *Zootaxa* 4005(1), 1-102.
- Singh, O. P. and J.P. Gupta 1981. Contribution to the Cytogenetics of an Indian Insect Pest. I. Karyotype and Salivary Chromosomes of *Melanagromyza obtusa* Malloch (Diptera: Agromyzidae). *Caryologia: Int. J. Cytol. Cytosyst. Cytogen.* 34(3), 275-282, Doi: 10.1080/00087114.1981.10796892
- Spencer, K.A. 1961. A synopsis of the Oriental Agromyzidae (Diptera). *Trans. Royal Entomol. Soc. London* 113, 55-100.
- Spencer, K.A. 1963. A synopsis of the Neotropical Agromyzidae. *Trans. Royal Entomol. Soc. London* 115, 291-389.
- Spencer, K.A. 1973. The Agromyzidae (Diptera) of economic importance. *Ser. Entomol.* 9, i-xi + 1-418.
- Spencer, K.A. 1983. Leaf mining Agromyzidae (Diptera) in Costa Rica. *Rev. Biol. Trop.* 31(1), 41-67.
- Spencer, K.A. 1984. The Agromyzidae (Diptera) of Colombia, including a new species attacking potato in Bolivia. *Rev. Colomb. Entomol.* 10(1-2), 3-33.
- Spencer, K.A. 1999. Family Agromyzidae (Australasian/Oceanian Diptera Catalog). En: <http://hbs.bishopmuseum.org/aocat/agromyzidae.html>; consulted: february, 2016.
- Spencer, K.A. and G.C. Steyskal. 1986. Manual of the Agromyzidae (Diptera) of the United States. U.S.D.A. Agriculture Handbook 638, 1-478.