

Radish (*Raphanus sativus* L.) morphophysiology under salinity stress and ascorbic acid treatments

Morfofisiología del rábano (*Raphanus sativus* L.) bajo estrés salino y tratamientos con ácido ascórbico

Ana Gabriela de Sousa Basílio¹, Leonardo Vieira de Sousa¹, Toshik Iarley da Silva^{2*}, Joana Gomes de Moura¹, Anderson Carlos de Melo Gonçalves³, José Sebastião de Melo Filho¹, Ygor Henrique Leal¹, and Thiago Jardelino Dias⁴

ABSTRACT

The use of saline or low-quality water in agriculture is an alternative to increasing water demand, especially in arid or semi-arid regions. However, the use of water with high levels of salts causes disturbances in plants, which can lead to their death; thus, alternatives to mitigate these effects are relevant in current agriculture. Currently, antioxidants are used to mitigate the effects of salts in plants, and among them ascorbic acid has been frequently mentioned. Therefore, the aim of this study was to evaluate the effect of irrigation with saline water combined with applications of ascorbic acid on the development and photosynthetic activity of radish (*Raphanus sativus* L.) plants. This experiment was carried out in a greenhouse with a randomized block design, with the treatments distributed in a 5×5 incomplete factorial scheme, composed of five electrical conductivities of the irrigation water (EC_w): 0.50, 1.30, 3.25, 5.20 and 6.00 dS m⁻¹, and five ascorbic acid (AA) doses: 0.00, 0.29, 1.00, 1.71, and 2.00 mM. The evaluated variables were: shoot height, leaf number, tuberous root diameter, chlorophyll a, b and total content, chlorophyll a/b ratio, initial fluorescence, maximum fluorescence, variable fluorescence and quantum yield of photosystem II. The saline water influenced the analyzed variables in the radish crop regardless of the ascorbic acid application. The ascorbic acid was not efficient in attenuating the deleterious effect of salinity in the irrigation water on the development and fluorescence of the radish. However, it was observed that the concentration of 1.00 mM of ascorbic acid promoted an increase in chlorophyll a, b and total in the salt-stressed radish plants.

Key words: abiotic stress, growth, photosynthesis, vegetable, crop.

RESUMEN

El uso de aguas salinas en la agricultura surge como alternativa al aumento en la demanda de agua, principalmente en regiones áridas o semiáridas. Sin embargo, el uso de agua con alto contenido de sales causa disturbios en las plantas pudiendo llevar hasta la muerte de las mismas. Por esta razón, se buscan alternativas para atenuar tales efectos. Actualmente, los antioxidantes se utilizan para mitigar los efectos de las sales en las plantas; entre estos antioxidantes el ácido ascórbico ha sido frecuentemente mencionado. Basado en lo expuesto, este trabajo tuvo por objetivo evaluar el efecto del riego con aguas salinas combinado con la aplicación de ácido ascórbico, sobre el desarrollo y actividades fotosintéticas de plantas de rábano (*Raphanus sativus* L.). El experimento fue desarrollado en ambiente protegido en bloques al azar con esquema factorial incompleto 5×5, con cinco conductividades eléctricas del agua de riego (EC_w: 0.50, 1.30, 3.25, 5.20 y 6.00 dS m⁻¹) y cinco dosis de ácido ascórbico (AA: 0.00, 0.29, 1.00, 1.71 y 2.00 mM). Las características evaluadas fueron altura de plantas, número de hojas, diámetro de la raíz tuberosa, índice de clorofila a, b, y total, relación de clorofila a/b, fluorescencia inicial, fluorescencia máxima, fluorescencia variable y rendimiento cuántico del fotosistema II. El uso de agua salina influenció todas las variables analizadas en el cultivo del rábano independiente de la utilización del ácido ascórbico. El uso del ácido ascórbico no fue eficiente como atenuante de los efectos de la salinidad del agua de riego sobre el desarrollo y la fluorescencia del cultivo del rábano. Sin embargo, se observó que la aplicación del ácido ascórbico en la dosis de hasta 1.00 mM promueve el aumento en los índices de clorofilas a, b y total del rábano sometido al estrés salino.

Palabras clave: estrés abiótico, crecimiento, fotosíntesis, hortaliza.

Introduction

The world's growing population in recent years has been accompanied by increased food demand, andoleraceous

crops are important to meeting this growing demand. In addition, oleraceous crops contribute directly to family farming activities (Noda and Noda, 2016).

Received for publication: 26 March, 2018. Accepted for publication: 6 November, 2018

Doi: 10.15446/agron.colomb.v36n3.74149

¹ Federal University of Paraíba, Center for Agrarian Sciences, Department of Plant Science, Areia (PB), (Brazil).

² Federal University of Viçosa, Department of Plant Science, Viçosa (MG), (Brazil).

³ Federal University of Roraima, Department of Plant Science, Boa Vista (RR), (Brazil).

⁴ Federal University of Paraíba, Center for Human, Social and Agrarian Sciences, Department of Agriculture, Bananeiras (PB), (Brazil).

* Corresponding author: iarley.toshik@gmail.com



The radish (*Raphanus sativus* L.) is an edible root vegetable of the Brassicaceae family, originated in the Mediterranean region, which presents globular swollen tap root with white or reddish skin. The radish is a fast-growing vegetable that can be harvested 25 to 35 d after sowing. The fast harvest cycle makes it particularly suitable for small and medium oleraceous producers, who can cultivate them along with longer cycle crops (Filgueira, 2008).

The radish contains significant amounts of vitamins A, C and B, and minerals such as calcium, phosphorus, magnesium, iron, sodium and potassium, which are essential nutrients to humans (Vidigal *et al.*, 2007). The radish crop, like other oleraceous crops, requires quality water to fully grow.

Irrigated agricultural systems require water availability and quality. The occurrence of low precipitation rates and saline soil conditions in arid and semi-arid regions around the world prevents salts from leaching, causing their accumulation within soil and water. High salt concentration conditions cause nutritional imbalance, reduced transpiration and photosynthesis rates, lower leaf CO₂ availability, and restricted stomatal conductance, which result in morphologically reduced plant biomass (Silva *et al.*, 2013; Silva *et al.*, 2018).

Salinity stress comprises two main components. The osmotic component, in which a high salt concentration in the soil solution reduces water availability to plants, and the ionic component, which is related to the toxicity of the ions released by the salts (Silveira *et al.*, 2010; Prisco *et al.*, 2017). Taking into account the necessity of using low quality water in irrigation systems, several studies have attempted to find ways to make these waters suitable for plant development and productivity (Oliveira *et al.*, 2016; Rodrigues *et al.*, 2017; Veras *et al.*, 2018).

Ascorbic acid is a major antioxidant found in plants and plays an important role in several cellular processes. The foliar application of ascorbic acid stimulates photosynthetic pigment production and plant growth under stress conditions. Several studies have indicated that ascorbic acid plays a fundamental role in protecting plants from multiple environmental stresses, including salinity stress (Maia *et al.*, 2012; Melo *et al.*, 2014).

In the literature, there are several reports on the use of ascorbic acid for attenuating the deleterious effects of saline stress, such as in *Triticum aestivum* (Abbasi and Fakhani, 2015; Siddiqui *et al.*, 2018), *Zea mays* (Billah *et al.*, 2017), *Linum usitatissimum* (El-Bassiouny and Sadak, 2015), *Citrus aurantium* (Kostopoulou *et al.*, 2015), and *Hordeum vulgare* (Agami, 2014). Based on this, the aim of this study was to evaluate the mitigating effect of ascorbic acid on the morphophysiological characteristics of salt stressed radish (*Raphanus sativus* L.).

Materials and methods

This experiment was carried out from October to November, 2017 in a greenhouse at the Agricultural Sciences Center of the Federal University of Paraiba (UFPB), located in the city of Areia, Paraiba, Brazil. The soil used in the experiment was collected in the 0-20 cm depth from the Plant Science Department land in the Federal University of Paraiba. This soil was classified as Latosol (EMBRAPA, 2014), and its chemical analysis is shown in Table 1.

The radish variety used was the hybrid Margaret Queen Kobayashi (ISLA®). Five seeds were sown per plastic bag at a depth of 2 cm, and emergence occurred 5 d later. Then, the treatments with the saline water irrigation, according to the predetermined EC_w values, were initiated. Ten d after emergence, the plants were thinned to one plant per bag.

The experimental design was a randomized block with the treatments distributed in a 5x5 factorial scheme, generated with the Central Box experiment matrix (Bortoluzzi and Alvarez, 1997). The factorial design was composed of five electrical conductivities of the irrigation water (EC_w): 0.50, 1.30, 3.25, 5.20 and 6.00 dS m⁻¹, and five ascorbic acid (AA) doses: 0.00, 0.29, 1.00, 1.71 and 2.00 mM, using four replicates. Plastic bags with a 3.0 dm³ capacity, containing one plant each, represented the experiment units.

The water with the EC_w of 0.50 dS m⁻¹ was obtained from the UFPB water supply system. A mixture of NaCl, CaCl₂.2H₂O and MgCl₂.6H₂O at a 7:2:1 ratio was added to the 0.50 dS m⁻¹ water to prepare the saline treatment water with the higher EC_w values (1.30, 3.25, 5.20 and 6.00 dS m⁻¹) (Medeiros, 1992). A micro processed portable conductivity meter CD-860 (Instrutherm, Brazil) was used to prepare

TABLE 1. Soil analysis report.

pH (H ₂ O)	OM (%)	P -----mg dm ⁻³ -----	K ⁺	Na ⁺	Ca ²⁺	Mg ²⁺	Al ³⁺	H ⁺ + Al ³⁺	BS	CEC	BSA (%)
-----cmol _c dm ⁻³ -----											
6.20	2.48	24.85	78.42	0.07	3.90	1.90	0.00	2.43	6.07	8.50	71.46

OM - organic matter, BS - base sum, CEC - cation exchange capacity, BSA - base saturation.

the saline waters. Irrigation management was measured using a drainage lysimeter (Alves *et al.*, 2017).

The radish leaves were sprayed with ascorbic acid until completely wet 14 d after seedling emergence. The non-ionic surfactant Tween 80[®] was added to the ascorbic acid solution for a better absorption effect at a concentration of 0.0002%.

At 15 and 30 d after seedling emergence, the following development variables were evaluated: shoot height, measured from base to leaf apex using a graduated ruler in centimeters; leaf number, counting all the leaves from each plant; tuberous root diameter determined at 2 cm of the soil using a digital pachymeter; chlorophyll a, chlorophyll b, total chlorophyll content and chlorophyll a/b ratio. The chlorophyll contents were determined with a portable electronic chlorophyll meter model CFL 1030, ClorofiLOG[®] (Falker, Brazil), and one reading was performed on each plant in totally expanded leaves.

The initial fluorescence (F_0), maximum fluorescence (F_m), variable fluorescence (F_v) and quantum yield of photosystem II (F_v/F_m) were measured at 28 d after seedling emergence. Chlorophyll fluorescence measurements (F_0 , F_m , F_v and F_v/F_m) were obtained using the modulated fluorometer Plant Efficiency Analyser - PEA II[®] (Hansatech Instruments Co., UK), and the readings were performed between 09:00 and 10:00 in the morning, on young, fully expanded, non-harmed and well lighted leaves. The fluorescence parameters were measured after a period of 30 min of dark adaptation with foliar tweezers (Konrad *et al.*, 2005). The obtained data were submitted to analysis of variance and regression using the software R (R Core Team, 2017).

Results and discussion

It was observed that the saline irrigation water had an effect on the development variables assessed at 15 and 30 d after seedling emergence. The shoot height, leaf number and tuberous root diameter were reduced with the increasing EC_w values (Fig. 1). Differences between the two assessment periods on the above-mentioned variables were observed once they were development variables, and over a period of 15 d there was an increase in size.

Aerial development is an important characteristic in salt stressed plant evaluation because it provides important information on plant response to stressful conditions (Jamil *et al.*, 2007). For the aerial part development variables, shoot height (Fig. 1A) had a higher value than the one predicted within the confidence interval in the electrical conductivity of 5.20 dS m⁻¹, with plants reaching a mean value of 10.5

cm. At the electrical conductivity of 6.00 dS m⁻¹, the radish plants reached an average value of 8 cm. Different results were observed by Putti *et al.* (2016), who registered no significant difference in radish behavior when submitted to 5.00 dS m⁻¹ level of salinity.

At 15 DAE, the leaf number (Fig.1B) remained constant at all EC_w levels. At 30 DAE, the leaf number decreased as the salinity increased. A similar result was found by Nascimento *et al.* (2015), who registered reduced leaf numbers in pepper (*Capsicum annum* L.) subjected to increasing salinity levels. Reduced leaf growth is a plant defense mechanism under stress conditions, such as water deficit and salinity, and it diminishes transpirational water loss (Taiz *et al.*, 2017). These results showed leaf sensitiveness to salinity stress, which made them reduced both in size and number as the salt concentration increased in the irrigation water.

For tuberous root diameter (Fig. 1C), a slight reduction was observed as the concentration of salts in the irrigation water increased at 15 and 30 DAE. The results obtained in this study corroborate those obtained by Bacarin *et al.* (2007), who studied the salinity effect on the growth and development of radish grown in a nutrient solution, and verified a reduction in leaf area and tuberous root diameter in response to the salinity stress.

Figure 2A shows shoot height as a function of the ascorbic acid treatment. The highest values were 7.8 cm at 15 DAE and 10.6 cm at 30 DAE, obtained in the absence of AA and at the dose of 1.71 mM, respectively. The lowest shoot height values were observed at the highest AA dose (2.00 mM), with 7.3 and 8.8 cm at 15 and 30 DAE, respectively.

For leaf number, a constant number of leaves was observed for all AA doses applied at 15 DAE. At 30 DAE, the radish plants treated with 0.29 mM of AA had a slight increase in the leaf number when compared to the control treatment plants (0.00 mM), showing that, as the concentrations of AA increased, a 13% decrease was observed in the leaf number, as compared to the control treatment (Fig. 2B).

Ascorbic acid is closely related to photosynthesis and respiration processes. The AA concentration in plant tissue varies as a function of plant response and different stress conditions, such as intensity and species sensitiveness to stress (Venkatesh *et al.*, 2012). Plants that present a low ascorbic acid synthesis capacity are very sensitive to environmental stress conditions, thus their growth and development are affected and, in some situations, an exogenous AA application is necessary in order to supply this antioxidant deficiency (Gao and Zhang, 2008).




FIGURE 1. Shoot height (A), leaf number (B) and tuberous root diameter (C) of radish (*Raphanus sativus* L.) subjected to different electrical conductivities of the irrigation water (ECw) at 15 and 30 DAE.




FIGURE 2. Shoot height (A), leaf number (B) and tuberous root diameter (C) of radish (*Raphanus sativus* L.) subjected to different ascorbic acid doses at 15 and 30 DAE.

At 15 DAE, increases in the AA concentrations resulted in a slightly reduced tuberous root diameter. The control plants presented a 0.9 mm tuberous root diameter, whereas the radish plants treated with 2.0 mM of AA presented a 0.7 mm tuberous root diameter. Different results were observed at 30 DAE, when the control plants presented a 34 mm tuberous root diameter, and the plants treated with 2.0 mM of AA reached 38 mm (Fig. 2C).

The foliar a, b, total and a/b chlorophyll indices (FCI) were influenced by the addition of salts in the irrigation water (Fig. 3). In general, the plants subjected to the highest electrical conductivity of the irrigation water (6.0 dS m⁻¹) and evaluated at 30 DAE showed the highest chlorophyll a, b and total values. For the assessment period, no differences were observed for the chlorophyll a or total chlorophyll. However, differences for chlorophyll b in electrical conductivities around 5.2 dS m⁻¹ and higher were observed. For

the chlorophyll a/b ratio, differences were only registered in the electrical conductivity of 3.25 dS m⁻¹.

No difference was observed in the chlorophyll a content as a function of time; in the first evaluation (15 DAE), the lowest value (24.62) was found in the ECw of 0.50 dS m⁻¹, and the highest value (25.91) was seen in the ECw of 1.30 dS m⁻¹, with a decreasing response from this electrical conductivity. In the second evaluation period (30 DAE), the minimum and maximum values were 23.75 and 27.54 in the ECw of 1.30 dS m⁻¹ and 6.00 dS m⁻¹, respectively (Fig. 3A).

The chlorophyll b content (Fig. 3B) presented the highest values of 5.92 at 15 DAE and 7.68 at 30 DAE, both occurring in the ECw of 6.00 dS m⁻¹. The assessment period influenced the chlorophyll b values in relation to the ECw treatment. When comparing the results obtained at 15 and 30 DAE, an increase for all tested ECw was observed,




FIGURE 3. Chlorophyll a (A), chlorophyll b (B), total chlorophyll (C) and chlorophyll a/b ratio (D) of radish (*Raphanus sativus* L.) subjected to different electrical conductivities of the irrigation water (ECw) at 15 and 30 DAE.

except for 1.30 dS m⁻¹. These increments were 10.8, 15.1, 20.2 and 32.8% for the EC_w of 0.5, 3.25, 5.20 and 6.00 dS m⁻¹, respectively.

Figure 3C shows that the lowest value for the total chlorophyll content at 15 DAE was 30.2, obtained in the EC_w of 0.50 dS m⁻¹, while the highest value (32.1) was found in the EC_w of 1.30 dS m⁻¹. At 30 DAE, the lowest value was 28.8, obtained in the EC_w of 1.30 dS m⁻¹, whereas the highest value of 35.1 was observed in the EC_w of 6.00 dS m⁻¹. The chlorophyll a/b ratio presented higher values at 15 DAE, with a maximum and minimum value of 4.7 and 4.25, obtained in the EC_w of 1.30 and 6.00 dS m⁻¹, respectively. In the second evaluation (30 DAE), the highest value was 4.28, and the lowest was 3.67, obtained in the EC_w of 1.30 and 0.50 dS m⁻¹, respectively (Fig. 3D).

These increases in the plant chlorophyll content when submitted to higher EC_w levels are contrary to those found by

some authors (Chaparzadeh and Behboud, 2015; Jasim *et al.*, 2016; Kasim *et al.*, 2016). However, their studies were conducted with higher salinity levels, and they did not clarify the radish variety used. According to Jamil *et al.* (2007), the chlorophyll content is reduced in plants that are sensitive to salinity and increases in tolerant plants.

The content of chlorophyll a, chlorophyll b, total chlorophyll and the chlorophyll a/b ratio were affected by the ascorbic acid treatment. The content of chlorophyll a and total presented no difference in relation to the assessment period. For the chlorophyll b content, there was a difference between 1.0 and 1.71 mM of AA, whereas for the chlorophyll a/b ratio, there was a difference in the 0.29, 1.0 and 1.71 mM of AA (Fig. 4).

At 15 DAE, the maximum value of chlorophyll a was 26.83, obtained when submitted to 0.29 mM of AA. From this AA dose on, the chlorophyll a content decreased and reached




FIGURE 4. Chlorophyll a (A), chlorophyll b (B), total chlorophyll (C) and chlorophyll a/b ratio (D) of radish (*Raphanus sativus* L.) subjected to different ascorbic acid doses at 15 and 30 DAE.

a minimum value of 25.09 at the dose of 1.71 mM. At 30 DAE, the lowest chlorophyll a content value (24.47) was observed in the absence of AA, and the highest (27.62) was obtained with the AA dose of 1.00 mM, with decreasing values from this dose on (Fig. 4A).

Figure 4B shows that the chlorophyll b content was affected during the evaluation periods. At 30 DAE, the chlorophyll b values were higher, with increases in all AA doses in relation to the evaluation performed at 15 DAE, representing increments of 5.6; 11.3; 18.6; 16.3 and 19.2% for the 0.00; 0.29; 1.00; 1.71 and 2.00 mM of AA, respectively.

The total chlorophyll presented little variation in relation to the different doses of AA and the two assessment periods (Fig. 4C). At 15 DAE, the highest value was 32.52, obtained in the AA dose of 0.29 mM, and, from this dose on, the total chlorophyll content started to decrease, with 30.08 being the lowest value at the dose of 1.71 mM. At 30 DAE,

the minimum value was 30.07, obtained in the absence of AA with a quadratic effect, and the maximum value was 34.65 with 1.00 mM of AA.

The lowest chlorophyll a/b ratios were observed in the AA dose of 1.00 mM, 4.28 and 3.79 at 15 and 30 DAE, respectively. The highest values for each assessment period were 4.78 at 15 DAE and 4.16 at 30 DAE, obtained with the doses of 0.29 mM and absence of AA, respectively (Fig. 4D). The chlorophyll content increase up to the 1.00 mM dose can be explained by the role of AA in regulating plant biochemical/physiological reactions (Khan *et al.*, 2006). However, higher doses of AA do not provide the same effects.

For initial fluorescence, it was observed that the maximum efficiency (93.6) was obtained under the electrical conductivity of 4.3 dS m⁻¹. The maximum and variable fluorescence presented mean values of 428.25 and 342.68, respectively (Fig. 5A-C). Distinct results were found in




FIGURE 5. A) Initial fluorescence (F_0); B) Maximum fluorescence (F_m); C) Variable fluorescence (F_v); and D) Quantum yield of photosystem II (F_v/F_m) of radish (*Raphanus sativus* L.) subjected to different electrical conductivities of the irrigation water (ECw).

the literature. When studying the NaCl effect, up to the concentration of 220 mM on salt-stressed radish plants, Bacarin *et al.* (2007) found no significant differences in the initial fluorescence as the salinity level increased.

For the quantum yield of photosystem II, the highest value (0.806) was obtained in the EC_w of 1.30 dS m⁻¹, while the lowest value was 0.792 in the EC_w of 6.00 dS m⁻¹ (Fig. 5D). Jamil *et al.* (2007) reported decreases in the quantum yield of photosystem II of radish plants as the salinity level increased.

Conclusions

The saline water influenced the analyzed variables in the radish (*Raphanus sativus* L.) plants regardless of the ascorbic acid application. The ascorbic acid was not efficient in attenuating the deleterious effect of salinity in the irrigation water on the development and fluorescence of the radish plants. However, the concentration of 1.00 mM of ascorbic acid promoted increased chlorophyll a, b and total in the salt-stressed radish plants.

Literature cited

- Abbasi, M. and E. Fakhani. 2015. Role of salicylic acid and ascorbic acid in the alleviation of salinity stress in wheat (*Triticum aestivum* L.). JBES 6, 107-113.
- Agami, R.A. 2014. Applications of ascorbic acid or proline increase resistance to salt stress in barley seedlings. Biol. Plantarum 58(2), 341-347. Doi: 10.1007/s10535-014-0392-y
- Alves, E.S., D.F. Lima, J.A.S. Barreto, D.P. Santos, and M.A.L. Santos. 2017. Determinação do coeficiente de cultivo para a cultura do rabanete através de lisimetria de drenagem. Irriga 22(1), 194-203. Doi: 10.15809/irriga.2017v22n1p194-203
- Bacarin, M.A., A.R. Falqueto, C.L. Moraes, P. Marini, and T.R. Löwe. 2007. Crescimento e fotossíntese em plantas de rabanete sob estresse salino. Rev. Bras. Agroc. 13(4), 473-479.
- Billah, M., M.M. Rohman, N. Hossain, and M.S. Uddin. 2017. Exogenous ascorbic acid improved tolerance in maize (*Zea mays* L.) by increasing antioxidant activity under salinity stress. Afr. J. Agr. Res. 12(17), 1437-1446.
- Bortoluzzi, A.L. and V.V.H. Alvarez. 1997. Pesquisa em casa de vegetação e em campo: matrizes experimentais. Departamento de Solo/CCA - UFV, Viçosa, Brazil.
- Chaparzadeh, N. and E.H. Behboud. 2015. Evidence for enhancement of salinity induced oxidative damages by salicylic acid in radish (*Raphanus sativus* L.). J. Plant Physiol. Breed. 5, 23-33.
- El-Bassiouny, H. and M.S. Sadak. 2015. Impact of foliar application of ascorbic acid and α-tocopherol on antioxidant activity and some biochemical aspects of flax cultivars under salinity stress. Acta Biol. Colomb. 20(2), 209-222. Doi: 10.15446/abc.v20n2.43868
- EMBRAPA - Empresa Brasileira de Pesquisa Agropecuária. 2014. Sistema brasileiro de classificação de solo. 4 ed. Brazil.
- Filgueira, F.A.R. 2008. Novo manual de olericultura: Agrotecnologia moderna na produção e comercialização de hortaliças. 3 ed. UFV, Viçosa, Brazil.
- Gao, Q. and L. Zhang. 2008. Ultraviolet-B-induced oxidative stress and antioxidant defense system responses in ascorbate-deficient *vtc1* mutants of *Arabidopsis thaliana*. J. Plant Physiol. 165, 138-148. Doi: 10.1016/j.jplph.2007.04.002
- Jamil, M., K.J. Lee, J.M. Kim, H.S. Kim, and E.S. Rha. 2007. Salinity reduced growth PS2 photochemistry and chlorophyll content in radish. Sci. Agricola 64(2), 111-118. Doi: 10.1590/S0103-90162007000200002
- Jasim, A.H., W.M.A. Al Timmen, and A.S. Abid. 2016. Effect of salt stress on plant growth and free endogenous hormones of primed radish (*Raphanus Sativus* L.) seeds with salicylic acid. Int. J. Chem. Tech. Res. 9(6), 339-346.
- Kasim, W.A., K.M. Saad-Allah, and M. Hamouda. 2016. Seed priming with extracts of two seaweeds alleviates the physiological and molecular impacts of salinity stress on radish (*Raphanus Sativus*). Int. J. Agric. Biol. 18(3), p. 653-660. Doi: 10.17957/IJAB/15.0152
- Khan, A., M.S.A. Ahmad, H.R. Athar, and M. Ashraf. 2006. Interactive effect of foliar applied ascorbic acid and salt stress on wheat (*Triticum aestivum* L.) at seedling stage. Pak. J. Bot. 38(5), 1407-1414.
- Konrad, M.L.F., J.A.B. Silva, P.R. Furlani, and E.C. Machado. 2005. Trocas gasosas e fluorescência da clorofila em seis cultivares de cafeeiro sob estresse de alumínio. Bragantia 64(3), 339-347.
- Kostopoulou, Z., I. Therios, E. Roumeliotis, A.K. Kanellis, and A. Molassiotis. 2015. Melatonin combined with ascorbic acid provides salt adaptation in *Citrus aurantium* L. seedlings. Plant Physiol. Biochem. 86, 155-165. Doi: 10.1016/j.plaphy.2014.11.021
- Maia, J.M., S.L. Ferreira-Silva, E.L. Voigt, C.D. Macedo, L.F.A. Ponte, and J.A.G. Silveira. 2012. Atividade de enzimas antioxidantes e inibição do crescimento radicular de feijão caupi sob diferentes níveis de salinidade. Act. Bot. Brasilica 26(2), 342-349. Doi: 10.1590/S0102-33062012000200010
- Medeiros, J.F. 1992. Qualidade da água de irrigação utilizada nas propriedades assistidas pelo "GAT" nos Estados do RN, PB, CE e avaliação da salinidade dos solos. 1992. MSc. thesis, Universidade Federal da Paraíba, Campina Grande, Brazil
- Melo, G.M., M.R. Barbosa, A.L.F. Dias, L. Willadino, and T.R. Camara. 2014. Pré-condicionamento *in vitro* de plantas de cana-de-açúcar (*Saccharum* spp.) para tolerância ao estresse salino. Rev. Bras. Eng. Agric. Amb. 18, 27-33.
- Nascimento, L.B., J.F. Medeiros, S.S.V. Alves, B.L.C. Lima, and J.L.A. Silva. 2015. Desenvolvimento inicial da cultura do pimentão influenciado pela salinidade da água de irrigação em dois tipos de solos. Agropec. Cient. Semi-Árido 11, 37-43.
- Noda, H. and S.N. Noda. 2016. Agricultura familiar tradicional e conservação da sócio-biodiversidade amazônica. Interações 4(6), 55-66.
- Oliveira, D.S., T.J. Dias, E.P. Oliveira, H.C. Santos, W.B. Magalhaes, B.F. Matos, L.M.C. Sousa, and J.S. Melo Filho. 2016. Growth and physiology of peanut (*Arachis hypogaea* L.) irrigated with

- saline water and biofertilizer application times. *Afric. J. Agric. Res.* 11(44), 4517-4524.
- Prisco, J.T., E. Gomes Filho, and R.S. Miranda. 2017. Physiology and biochemistry of plants growing under salt stress. pp. 425-448. In: Gheyi, H.R., N.S. Dias, C.F. Lacerda, and E. Gomes Filho (eds.). *Manejo da salinidade na agricultura: Estudos básicos e aplicados*. INCTSal, Fortaleza, Brazil.
- Putti, F.F., J.F. Silva Júnior, R. Ludwig, L.R.A. Gabriel Filho, C.P. Cremasco, and A.E. Klar. 2014. Avaliação da cultura do rabanete ao longo do ciclo submetido em diferentes níveis de salinidade. *J. Agron. Sci.* 3(2), p. 80-90.
- R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.
- Rodrigues, R.M., L.F. Cavalcante, A.G.L. Souto, H.R. Gheyi, and F.O. Mesquita. 2017. Growth and regrowth of neem after cutting in saline-sodic soil treated with organics inputs. *Rev. Caatinga* 30, 116-124. Doi: 10.1590/1983-21252017v30n113rc
- Siddiqui, M.H., S.A. Alamri, M.Y. Al-Khaishany, M.A. Al-Qutami, and H.M. Ali. 2018. Ascorbic acid application improves salinity stress tolerance in wheat. *Chiang Mai J. Sci.* 45(3), 1296-1306.
- Silva, E.M., C.J.G.S. Lima, S.N. Duarte, F.S. Barbosa, and R. Maschio. 2013. Níveis de salinidade e manejo da fertirrigação sobre características da berinjela cultivada em ambiente protegido. *Rev. Ciênc. Agron.* 44, 150-158.
- Silva, T.I., J.S. Melo Filho, A.C.M. Gonçalves, L.V. Sousa, J.G. Moura, T.J. Dias, J.C. Alvarez-Pizarro, W.E. Pereira, and R.M.N. Mendonça. 2018. Salicylic acid effect on *Ocimum basilicum* L. during growth in salt stress and its relationship between phytomass and gas exchange. *J. Exp. Agric. Inter.* 22(4), 1-10. Doi: 10.9734/JEAI/2018/40901
- Silveira, J.A.G., S.L.F. Silva, E.N. Silva, and R.A. Viegas. 2010. Mecanismos biomoleculares envolvidos com a resistência ao estresse salino em plantas. pp. 166-185. In: Gheyi, H.R., N.S. Dias, and C.F. Lacerda (eds.). *Manejo da salinidade na agricultura: estudos básicos e aplicados*. INCTSal, Fortaleza, Brazil.
- Taiz, L., E. Zeiger, I.A. Møller, and A. Murphy. 2017. *Fisiologia e desenvolvimento vegetal*. Artmed, Porto Alegre, Brazil.
- Venkatesh, J., C.P. Upadhyaya, J.W. Yu, A. Hemavathi, D.H. Kim, R.J. Strasser, and S.W. Park. 2012. Chlorophyll a fluorescence transient analysis of transgenic potato overexpressing D-galacturonic acid reductase gene for salinity stress tolerance. *Hort. Envir. Biotec.* 53(4), 320-328. Doi: 10.1007/s13580-012-0035-1
- Veras, M.L.M., J.S. Melo Filho, L.S. Alves, T.H.S. Irineu, and T.J. Dias. 2018. Growth and dry matter pitombeira seedlings under salinity levels and application of biofertilizer. *Com. Scientia* 8(3), 486-492. Doi: 10.14295/cs.v8i3.1881
- Vidigal, S.M. and M.W. Pedrosa. 2007. Rabanete. pp. 661-664. In: Paula Júnior T.J., and M. Venzon. (eds.) *101 culturas - Manual de tecnologias agrícolas*. EPAMIG, Belo Horizonte, Brazil.