

A model of sulphur solubility for hydrous mafic melts: application to the determination of magmatic fluid compositions of Italian volcanoes

Bruno Scaillet and Michel Pichavant
ISTO-CNRS, UMR 6613, Orléans, France

Abstract

We present an empirical model of sulphur solubility that allows us to calculate fS_2 if P , T , fO_2 and the melt composition, including H_2O and S , are known. The model is calibrated against three main experimental data bases consisting in both dry and hydrous silicate melts. Its prime goal is to calculate the fS_2 of hydrous basalts that currently lack experimental constraints of their sulphur solubility behaviour. Application of the model to Stromboli, Vesuvius, Vulcano and Etna eruptive products shows that the primitive magmas found at these volcanoes record fS_2 in the range 0.1-1 bar. In contrast, at all volcanoes the magmatic evolution is marked by dramatic variations in fS_2 that spreads over up to 9 orders of magnitude. The fS_2 can either increase during differentiation or decrease during decompression to shallow reservoirs, and seems to be related to closed *versus* open conduit conditions, respectively. The calculated fS_2 shows that the Italian magmas are undersaturated in a FeS melt, except during closed conduit conditions, in which case differentiation may eventually reach conditions of sulphide melt saturation. The knowledge of fS_2 , fO_2 and fH_2O allows us to calculate the fluid phase composition coexisting with magmas at depth in the C-O-H-S system. Calculated fluids show a wide range in composition, with CO_2 mole fractions of up to 0.97. Except at shallow levels, the fluid phase is generally dominated by CO_2 and H_2O species, the mole fractions of SO_2 and H_2S rarely exceeding 0.05 each. The comparison between calculated fluid compositions and volcanic gases shows that such an approach should provide constraints on both the depth and mode of degassing, as well as on the amount of free fluid in magma reservoirs. Under the assumption of a single step separation of the gas phase in a closed-system condition, the application to Stromboli and Etna suggests that the main reservoirs feeding the eruptions and persistent volcanic plumes at these volcanoes might contain as much as 5 wt% of a free fluid phase. Consideration of the magma budget needed to balance the amounts of volatiles emitted in the light of these results shows that the amount of non-erupted magma could be overestimated by as much as one order of magnitude.

Key words sulphur – hydrous basalts – volcanic gas – Italy

1. Introduction

The determination of the composition of the fluids that coexist with magmas at depth requires

1) a good definition of the magma pre-eruptive conditions (P , T , fO_2 , fS_2 , fH_2O) and 2) solubility models for the main volatile species. The combination of these two sets of information allows us to calculate the fluid composition through the consideration of fluid-melt equilibria (e.g., Scaillet and Evans, 1999). For any volatile species dissolved in a silicate melt that is saturated in a fluid, equilibrium conditions demand that the fugacity f_i of species i in the melt equals its fugacity in the fluid. For water we have therefore

$$fH_2O_{\text{melt}} = fH_2O_{\text{fluid}}. \quad (1.1)$$

Mailing address: Dr. Bruno Scaillet, ISTO-CNRS, UMR 6613, 1A Rue de la Férollerie, 45071 Orléans cedex 02, France; e-mail: bscaille@cnrs-orleans.fr

The relationships between fugacity (or activity) and concentration in silicate melts have been established mainly for H₂O and CO₂ for a wide range of melt compositions (*e.g.*, Dixon *et al.*, 1995; Papale, 1997; Zhang, 1999). Once the fugacity of a given volatile species has been determined, its partial pressure, P_i , can be determined via the simple eq. (1.2)

$$P_i = \frac{f_i}{\gamma_i} \quad (1.2)$$

where γ_i is the fugacity coefficient of species i in the fluid.

In the system C-O-H, the other main species to be considered is CO₂, except if fO_2 is significantly below FMQ, in which case reduced species such as H₂, CO, and CH₄ can be present as well (see Holloway, 1977). Therefore, considering that H₂O and CO₂ are the main volatile species in the fluid, we have the following equations:

$$P_{H_2O} = \frac{f_{H_2O}}{\gamma_{H_2O}} \quad (1.3)$$

$$P_{CO_2} = \frac{f_{CO_2}}{\gamma_{CO_2}} \quad (1.4)$$

Since the fluid can be treated as a binary H₂O-CO₂ system, we have also the constraint, X_i being the mole fraction of species i in the fluid

$$X_{H_2O} + X_{CO_2} = 1. \quad (1.5)$$

Recalling that $P_i = X_i P_{\text{fluid}}$, eqs. (1.3) to (1.5) can be solved simultaneously to calculate both P_{H_2O} and P_{CO_2} and thus P_{fluid} since by definition

$$P_{\text{fluid}} = \sum P_i. \quad (1.6)$$

What is calculated through this approach is thus the pressure at which the melt becomes saturated in a fluid phase and the composition of this fluid. This method offers thus the possibility to constrain both the minimum pressure of magma reservoir and the fluid composition that is likely to escape this reservoir during an eruptive process. Obviously those two information are of considerable interest for the assessment of volcanic hazards. This approach was first applied using modern solubility models for H₂O

and CO₂ by Anderson *et al.* (1989) to the Bishop tuff eruption. Since then, several petrological studies have estimated the pressure depth at which magmas become fluid-saturated, assuming that the fluid phase is a binary H₂O-CO₂ mixture (*e.g.*, Roggensack *et al.*, 1997).

Most arc magmas are, however, notoriously rich in sulphur. Over the last 15 years, several melt inclusions studies have shown that the pre-eruptive melt sulphur content of arc basalts is equal to, or even higher than, the amount of dissolved CO₂ (*e.g.*, Roggensack *et al.*, 1997; Marianelli *et al.*, 1995, 1999). This shows that sulphur bearing volatile species must be incorporated into the above approach for a more complete description of fluid-melt equilibria. The nature and proportion of S-bearing species, mostly H₂S and SO₂, strongly depend on fO_2 , as for C-bearing species (see Holloway, 1977; Symonds *et al.*, 1994). Unlike C-species, however, the redox range in which the H₂S/SO₂ ratio varies is exactly within the redox range recorded by most arc-magmas (*i.e.* NNO to NNO+2). This fO_2 sensitivity introduces an additional complication in the approach outlined above, since the role of fO_2 on volatile speciation must be explicitly taken into account. The calculation of the pressure for fluid saturation of a magma saturated in a C-O-H-S fluid at an fO_2 equal to or higher than NNO will require to solve the following equation:

$$P_{\text{fluid}} = P_{H_2O} + P_{CO_2} + P_{SO_2} + P_{H_2S} \quad (1.7)$$

if the contribution of CO, CH₄, H₂, S₂, and O₂ partial pressures are ignored, which is a valid approximation under the redox conditions considered here ($fO_2 > \text{NNO}$). The SO₂ and H₂S species abundances are controlled by the following equilibria:



which show that what is needed to calculate S-bearing species fugacities is fS_2 or one over the three S-bearing species fugacities (assuming that both fO_2 and fH_2 are known, which are two

quantities generally known or obtainable if the pre-eruptive conditions are determined in magmas).

The preceding section shows that there is a need for a solubility model for S in silicate melts. Up to recently, there was only one such a model (Wallace and Carmichael, 1992) which was calibrated for dry basalts and fO_2 below NNO, using mostly the experimental data of Haughton *et al.* (1974). Clemente *et al.* (2004) have developed both a thermodynamic and an empirical sulphur solubility model for hydrous rhyolitic compositions. In the intermediate composition range, that is between basalt and rhyolite, only a few studies have been carried out at one bar (*e.g.*, Katsura and Nagashima, 1974) and even fewer at high pressure (Carroll and Rutherford, 1987; Luhr, 1990; Carroll and Webster, 1994). Of the work done at high pressure, only that of Luhr (1990) provides sulphur solubility data obtained under known fS_2 estimated from either Fe-FeS equilibria or equilibrium assemblages involving anhydrite. Clearly, it is this mid-compositional range of hydrous mafic to intermediate magmas that still lacks experimental coverage, and this is unfortunate because most of arc magmas belong to this category. Moretti *et al.* (2003) devised a general solubility model that should be applicable to a wide range of silicate melt compositions, including those hydrous. However, although based on a rigorous, as well as promising, thermodynamic description of melt-fluid equilibria, the current version of the model is quite complex to implement and is presently restricted to conditions of melt undersaturation with respect to a Fe-S-O sulphide melt.

In the present paper, we follow a more pragmatic approach and derive an empirical solubility model by considering the existing experimental database on geologically relevant silicate liquids. We have deliberately ignored the wealth of data existing in the metallurgical literature, since 1) it is exclusively based on dry compositions, 2) it concerns silicate melt compositions far outside the compositional spectrum displayed by terrestrial magmas, and 3) it covers temperatures greater than those typical of Earth's magmatism. Our first target is to build a model that allows us to compute the fS_2 of basaltic to intermediate hydrous melts, using

as input parameters P , T , fO_2 and melt composition. Although we believe that the model described below retrieves the correct order of magnitude in terms of fS_2 , we envision it as an intermediate step that should help define future experimental strategies to obtain high pressure solubility data onto which more rigorous thermodynamic approaches, such as that of Moretti *et al.* (2003), will be calibrated. The second target of this report is the application of this model to Italian volcanoes for which there exists a combined data set of pre-eruption H_2O , CO_2 and S concentrations, in addition to other intensive parameters (T and fO_2). We considered Stromboli, Vesuvius, Vulcano and Etna volcanoes, for which there are melt inclusion constraints on pre- to syn-eruptive volatile abundances. We calculated the equilibrium fluid composition corresponding to the depth at which the melt inclusion were saturated in fluid. The rationale here is clear: to constrain fluid compositions from depth to near surface conditions and compare it with the observed gas composition measured at the exit (see Scaillet and Pichavant, 2003). The combination of the two sets of data (calculation and observation) should allow a better use of volcanic gases as monitoring tools during on-going volcanic crisis.

2. An empirical model for hydrous basaltic melts

We attempted to devise a general empirical model in which the different effects of fO_2 , fS_2 and T on the melt sulphur content, as well as the melt composition, are explicitly taken into account. Several equations were tested and the following was found to give satisfactory results

$$\log S = aP + bT + c\Delta NNO^3 + d\Delta NNO^2 + e\Delta NNO\Delta FFS + f\Delta FFS + \sum g_i W_i \quad (2.1)$$

where S is the total sulphur concentration in ppm, P the pressure in bar, T the temperature in °C, ΔNNO and ΔFFS are the referenced fO_2 and fS_2 as explained below, W_i represents the weight % of oxide i , and a , b , c , d , e , f and g_i are fitted parameters listed in table I. The summa-

Table I. Regression coefficients for the empirical model of sulphur solubility.

P	7.28E-06
NNO^3	0.00488128
NNO^2	0.0818873
NNOFFS	-0.0224068
T	0.00084107
FFS	0.22801636
SiO_2	-0.012467
Al_2O_3	-0.0015766
Fe_2O_3	0.37362348
FeO	0.0674383
MgO	0.01121929
CaO	0.02000831
Na_2O	0.05644745
K_2O	-0.0248037
TiO_2	0.00672403
H_2O	0.06868295
OH	0.05778453

tion is carried over all major oxides, including FeO , Fe_2O_3 , OH^- and H_2O . The $\text{Fe}^{2+}/\text{Fe}^{3+}$ ratios of experimental glasses were calculated using the empirical model of Kilinc *et al.* (1984), whereas the water contribution was split into hydroxyl and molecular water using the method of Zhang (1999). The third order polynomial function is necessary to reproduce the dissymmetrical inverted bell-shaped pattern (*e.g.*, Katsura and Nagashima, 1974), while the crossed $f\text{O}_2$ - $f\text{S}_2$ term is needed to take into account the effect of varying $f\text{S}_2$ on the relationship between $f\text{O}_2$ and S in melt (see Clemente *et al.* 2004). Using mole fractions instead of weight% oxides, or any other intensive parameters (*i.e.* $f\text{H}_2$ in lieu of $f\text{O}_2$), does not improve the quality of the fit. Similarly, to account for the compositional dependence various norm projections were tested including the CIPW norm, but none was found to decrease the residuals. As for $f\text{O}_2$, which is referenced to the NNO solid buffer, we have referenced $f\text{S}_2$ to the iron-troilite solid buffer (Fe-FeS or FFS) such that

$$\Delta\text{FFS} = \log f\text{S}_2(\text{melt}) + \log f\text{S}_2(\text{FFS}) \quad (2.2)$$

in which the $f\text{S}_2$ of the Fe-FeS buffer is given by (Froese and Gunter, 1976)

$$\log f\text{S}_2(\text{FFS}) = \left\{ (2/1.987T) [-0.2655(P-1) - 35910 + 12.56T] \right\} / 2.302581 \quad (2.3)$$

where P is in bar and $T(K)$ is the absolute temperature.

We derived a set of fitted $a \dots g_i$ parameters for eq. (2.1) by linear regression considering the data bases of Luhr (1990), O'Neill and Mavrogenes (2002), Katsura and Nagashima (1974) on rhyodacite melts, in addition to those of Clemente *et al.* (2004). As stated above, although the existing experimental database is considerably larger, we restricted the regression procedure to the above studies for the following reasons: first, all works were specifically aimed at exploring the behaviour of sulphur in geological melts and are thus of direct relevance to magmatic contexts; second, they were carried out within P - T - $f\text{O}_2$ - $f\text{S}_2$ - $f\text{H}_2\text{O}$ conditions that closely approach those under which natural magmas evolve; third, use of the complete database is not warranted since much of it concerns slags or synthetic silicate melts of metallurgical, rather than geological, interest. Because our model is empirical, including this database will have the undesirable but unavoidable consequence that one atmosphere data overwhelmingly dominate over those obtained at high pressure and thus that the fitted parameters be biased toward anhydrous compositions. Preliminary fitting procedures used the database of Haughton *et al.* (1974). However, the extensive work of O'Neill and Mavrogenes (2002) on dry mafic melts at one bar has shown that a substantial part of the experimental database gathered by Haughton *et al.* (1974) does not obey expected thermodynamic relationships, and for this reason we have not included this work in our database.

The selected database consists therefore of 64 hydrous rhyolitic compositions (Clemente *et al.* (2004) plus three charges of Scaillet *et al.* (1998) for which $f\text{S}_2$ has been measured using the same technique than in Clemente *et al.* (2004)), 72 hydrous rhyolite-andesite compositions (Luhr, 1990), 333 dry mafic to ultramafic

compositions (O'Neill and Mavrogenes, 2002), and 5 dry rhyodacite compositions of Katsura and Nagashima (1974). There is therefore a 30/70 repartition between hydrous and dry compositions. The experiments of Luhr (1990) and Clemente *et al.* (2004) are the only ones in which both fO_2 and fS_2 are known for hydrous compositions, and cover an fO_2 range between NNO-1 up to NNO+3, a total pressure of 100-400 MPa, a temperature 800-1000°C, for melt

water contents up to 8 wt%, the melt compositions spanning a SiO_2 range of 57-80 wt% on an anhydrous basis. The experiments of Katsura and Nagashima (1974) were included in the regression in an effort to constrain the sulphur solubility in silicic compositions under dry, *i.e.* low pressure, conditions. The database of O'Neill and Mavrogenes was chosen among others (*e.g.*, Buchanan and Nolan, 1979; apart from that of Haughton *et al.*, 1974, for the reason given above) because of its systematic variation of melt composition at constant fO_2 and fS_2 , which allows a better evaluation of the compositional dependence of sulphur solubility (*i.e.* the g_i terms in eq. (2.1)). The fitted parameters are listed in table I.

Figure 1a-c shows that the model reproduces observed sulphur concentrations of the whole data set over more than 3 orders of magnitude (10-11900 ppm), with no apparent systematic divergence from the 1/1 correlation line within any of the three main data subsets, except in the low concentration range. Considering that the database covers a wide compositional range (SiO_2 ranging from 35 to 80 wt%), as well as widely differing P - T -fluid fugacities conditions, the agreement between observed and calculated sulphur concentrations is considered satisfactory. In terms of fS_2 , the model reproduces measured fS_2 within an average of 0.65 log unit, over more than 15 log units when normalised to the FFS solid buffer. Inspection of fig. 1a-c shows nevertheless that hydrous melts are not as well reproduced by the model, which is a reflection of the difficulty in accurately controlling fS_2 in high pressure experiments, since there is no buffering technique which allows to measure fS_2 with a precision similar to that of fO_2 or fH_2 . Given that there exist solubility models for rhyolite compositions (Clemente *et al.*, 2003) and for dry mafic melts (*e.g.*, Wallace and Carmichael, 1992; O'Neill and Mavrogenes, 2002), this empirical model is intended primarily to be used for melt compositions currently lacking experimental constraints, that is, hydrous mafic melts.

Equation (2.1) is merely a convenient mathematical way to describe the interdependence between the various parameters that control the sulphur solubility in silicate melts. As such the




Fig. 1a-c. Comparison between measured and calculated melt sulphur concentrations for the three main experimental data sub set used to calibrate the empirical model (eq. (2.1), see text). a) rhyolitic compositions (Clemente *et al.*, 2004; Scaillet *et al.*, 1998; Katsura and Nagashima, 1978); b) intermediate compositions (Luhr, 1990); c) basaltic composition (O'Neill and Mavrogenes, 2002).

fitted parameters have clearly no thermodynamical meaning. Yet, it is useful to consider at this stage the individual effect of some of the parameters on the calculated fS_2 (or sulphur solubility) to appreciate their relative importance on sulphur behaviour in hydrous magmas. All parameters with a positive value increase the sulphur solubility, when everything else is kept constant. For instance at constant fO_2 , fS_2 , P and melt composition, a temperature increase produces an increase in melt sulphur content. The same is true for P which suggests that, in hydrous systems, an increase in pressure increases the sulphur solubility, a trend opposite to that found for dry mafic melts saturated in an immiscible sulphide liquid (see Mavrogenes and O'Neill, 1999). Of the melt components, only SiO_2 , Al_2O_3 and K_2O have negative signs. Although this is as expected for SiO_2 and Al_2O_3 because both components increase the degree of melt polymerisation, and thus they decrease its capacity to exchange sulphur ions with free oxygens, the negative sign found for K_2O is more surprising since this element is often positively correlated with melt sulphur content (see Métrich and Clocchiatti, 1996). Whether this effect is real or is merely a reflection of a compositional bias in the database cannot be solved in the present study. The large value found for Fe_2O_3 may indicate preferential association of oxidized sulfur groups with Fe^{3+} melt components but we stress that in the database, oxidized melts are mostly intermediate to silicic compositions with a low bulk FeO and thus a relatively low Fe_2O_3 content. Experiments performed on iron-rich basalts have been mostly performed at low fO_2 , and the behaviour of sulphur in dry oxidized basalt still demands experimental investigations. A last aspect of interest concerns the role of water. Both molecular H_2O and hydroxyl group have strong positive values which suggests that addition of water to a silicate melt held at constant P - T - fO_2 - fS_2 conditions significantly increases its melt sulphur content. We note, however, that because high pressure experiments are also those that are hydrous in the database, correctly discriminating the effects of pressure and water content on sulphur solubility is not yet possible. Overall the empirical model can be considered as calibrat-

ed in the pressure range 1-4000 bar, temperature range 800-1400°C, melt water content range 0-10 wt%, fO_2 range NNO-2 to NNO+3, and melt sulphur contents 50-10000 ppm. Melt compositions not yet considered and to which the model should be applied with caution are evolved alkali-rich magmas, such as phonolites and peralkaline rhyolites.

3. Calculation of fluid phase composition

To calculate the fluid phase composition we follow the same approach as Scaillet and Pichavant (2003) who determined the fluid compositions of a number of intermediate to silicic arc magmas. They also attempted to constrain the fluid phase of some hydrous arc basalts, using a simplified version of the Wallace and Carmichael (1992) model. The calculations were performed considering that the fluid composition can be described in the C-O-H-S system. We thus ignore the contribution of halogens, in particular F and Cl, although we recognise that those species may play an important role in the evolution of magmatic fluids (see Carroll and Webster, 1994), especially in alkali-rich magmas (*e.g.*, Métrich, 1990; Webster and De Vivo, 2002).

The species considered are H_2O , H_2 , CO_2 , CO , CH_4 , H_2S , SO_2 , S_2 and O_2 . To calculate the fluid compositions we use as input parameters, fO_2 , fS_2 and fH_2O . The thermodynamic model of water solubility for basalt of Dixon *et al.* (1995) is used to calculate fH_2O from the measured melt H_2O content. The fO_2 is constrained from petrological studies (*i.e.*, spinel-olivine-melt equilibria; Ballhaus *et al.*, 1991) or sulphur speciation (*e.g.*, Métrich and Clocchiatti, 1996). The fS_2 is calculated from eq. (2.1).

At any fixed P and T , fixing fO_2 , fS_2 and fH_2O allows us to calculate the fugacities of all remaining species. For pressures below 5 kbar, we use the Modified Redlich Kwong equation of state (see Holloway, 1977; Flowers, 1979) with mixing rules as given by Ferry and Baumgartner (1987). At higher pressure, we use the corresponding state equation of Shi and Saxena (1992) and Saxena and Fei (1987) to calculate the fugacities of pure species. The fugacities of the species in the fluid mixture were derived us-

ing the Lewis and Randall rule: the fugacity coefficient of species i in the mixture equals that of pure species i at the same P and T .

Because the fluid pressure is unknown, the calculation must be done iteratively. The following procedure was adopted. We first fix P , T , and calculate the corresponding $f\text{H}_2\text{O}$ and $f\text{S}_2$ from melt inclusion data. This results in a $f\text{CO}_2$ which is used to calculate the equilibrium CO_2 content of the melt, using the solubility model of Dixon *et al.* (1995). This CO_2 content is then compared to that measured in the melt inclusion and if the difference between observed and calculated values exceeds 10 ppm, the calculation is performed again at a different pressure until the test is fulfilled. Because P variations affect both $f\text{S}_2$ and $f\text{H}_2\text{O}$ for a given set of T - $f\text{O}_2$ -melt composition values, tests are also performed on the difference between observed and calculated values in melt S and H_2O contents. The pressures of fluid saturation listed below reproduce observed H_2O , CO_2 and S melt contents to ± 0.01 wt%, 10 ppm, 1 ppm, respectively.

For alkali-rich basalts, we corrected for the alkali effect using the following method which is based on the approach of Dixon (1997) to evaluate the increase in CO_2 solubility resulting from an increase in melt alkalinity. For a given calculated $f\text{CO}_2$, we first determine the melt CO_2 content using the solubility model of Dixon *et al.* (1995) which is calibrated on MORB composition. This amount of CO_2 is then adjusted using the following empirical fit:

$$\text{CO}_2(\text{ppm}) = \text{CO}_{2\text{-MORB}}(\Pi - 0.472)0.8092 * \Pi + \text{CO}_{2\text{-MORB}} \quad (3.1)$$

where $\text{CO}_{2\text{-MORB}}$ is the CO_2 content of a MORB melt at P and T (in ppm) calculated using the model of Dixon *et al.* (1995), and Π is a compositional parameter devised by Dixon (1997) to evaluate the effect of metal cations on CO_2 solubility, and is equal to

$$\Pi = -6.5(\text{Si}^{4+} + \text{Al}^{3+}) + 20.17(\text{Ca}^{2+} + 0.8\text{K}^+ + 0.4\text{Mg}^{2+} + 0.4\text{Fe}^{2+}) \quad (3.2)$$

where the Si, Al ... are cationic fractions, and

Fe^{2+} is computed from total iron expressed as FeO_{tot} (only Fe^{2+} in Dixon, 1997). Equation (3.1) was derived by linear regression of solubility data (see Holloway and Blank, 1994; Dixon, 1997) obtained at 1 kbar and 1200°C on MORB tholeiite, Kilauea tholeiite, basanite and leucitite melts which encompass a Π range of 0.47 (MORB) to 2.35 (leucitite). Equation (3.1) back calculates experimental solubility data with an average standard deviation of 80 ppm at 1 kbar. Inspection of experimental data shows that in fact the CO_2 solubility of mafic melts is constant within analytical uncertainties for Π in the range 0.47-0.79, so that we have applied eq. (3.1) only to melt compositions having a Π value higher than 0.8. Although the correction factor is likely to vary with pressure and temperature, we have applied eq. (3.1) at all P and T since there are not enough data to properly evaluate this effect over a large range of P - T conditions.

We note, in addition, that the compositions used to derive eq. (3.1) are significantly less alkali-rich than some of the basalts erupted at Italian volcanoes: in particular, the tephrites of Vesuvius have K_2O contents almost twice higher than that of the most alkali-rich composition used to derive eq. (3.1). The strong correlation observed in experimental mafic melts between




Fig. 2. Correlation between dissolved CO_2 and the K_2O content of experimental basaltic melts used to determine the solubility of CO_2 at 1000 bar and 1200°C . See Dixon (1997) for the source of data. Vesuvius tephrites, between 4 and 5 wt% K_2O , could have CO_2 solubilities close, or in excess, to 4000 ppm, or nearly 10 times more than MOR basalts (*ca.* 500 ppm under these conditions).

K₂O and CO₂ content (fig. 2) suggests that K₂O may have a decisive role on CO₂ solubility in mafic melts perhaps far outweighing that of other metal cations. Simple extrapolation of the trend shown on fig. 2 would imply that CO₂ solubility in a tephrite with 5 wt% K₂O at 1 kb and 1200°C could be nearly ten times that of MORB or 3000–4000 ppm, while application of the method of Dixon (1997) with $\Pi=1$ predicts a much smaller increase in CO₂ solubility (from ca. 500 ppm for a MORB to 724 ppm for a tephrite). In support of this enhanced CO₂ solubility are the phase equilibria of Trigila and De Benedetti (1993), which have shown that addition of CO₂ to dry tephrite at 2 kbar depresses the liquidus temperatures of pyroxene, leucite and plagioclase by 50–75°C with estimated CO₂ concentrations in melt in excess of 1 wt% (see also Freda *et al.*, 1997, for similar findings on phonotephritic magmas of the Alban Hills). A freezing point depression effect in a given silicate-CO₂ system implies a significant solubility of CO₂ in the melt. Although such a potassium effect on CO₂ solubility cannot be rigorously modeled at present, the above observations suggest that the pressures of fluid saturation calculated here could be largely overestimated for mafic melt compositions that contain significantly more K₂O than the rock melts so far investigated in experimental studies on CO₂ solubility.

In summary we believe that mafic melts moderately enriched in K₂O relative to MORB can have their CO₂ solubility, and thus their pressure of fluid saturation, correctly estimated (to within 0.5 kbar) by the method explained above. This is the case for most of the mafic magmas erupted at Etna, Stromboli and Vulcano. In contrast, there exists the possibility that the pressures of fluid saturation calculated in this work for K₂O-rich mafic magmas such as Vesuvius tephrites, are largely overestimated, perhaps by as much as several kilobars. This indicates that there is an urgent need of experimental constraints on CO₂ solubility in alkali-rich basalts.

4. Source of data

In this section we briefly review the present status of knowledge on pre-eruptive H₂O-CO₂-

S melt concentrations as well as of P - T - fO_2 of the different volcanoes considered. We are primarily interested in works where those three volatiles have been determined (*e.g.*, Marianelli *et al.*, 1999; Métrich *et al.*, 2001). However, we also considered melt inclusion data sets lacking CO₂ determination but for which both H₂O and S are known (*e.g.*, Marianelli *et al.*, 1995; Métrich *et al.*, 1993; Cioni *et al.*, 1995). Although the lack of CO₂ constraints prevents the calculation of the pressure of fluid saturation and fluid composition, H₂O and S abundances can still be used to infer fS_2 , provided there are T - fO_2 constraints. In such a case, the pressure for calculation in eq. (2.1) is that corresponding to the H₂O solubility. While this pressure is clearly a minimum, in terms of fS_2 calculation the error introduced by this uncertainty is minor, compared to that due to fO_2 : a pressure increase of 4 kbar decreases the calculated $\log fS_2$ by 0.5 log unit. In the following, we focus primarily on mafic primitive melts, since it is the injection of mafic fresh magmas that is more likely to re-awaken or trigger a new explosive event, as illustrated by the April 5th, 2003 event at Stromboli.

4.1. Stromboli

We restricted our calculations to melt inclusions analysed by Métrich *et al.* (2001), who studied the products of paroxysmic eruptions such as that of the August 23, 1998 explosion, during which crystal-rich scoria and highly vesicular yellow pumices were erupted. The first type of rock is interpreted to represent quenched fragments of a crystal-rich degassed magma that was resident in a very shallow magma chamber. This reservoir was intercepted by a rising batch of crystal-poor and volatile-rich magma, now represented by the yellow pumice. Melt inclusion constraints on volatile abundances were obtained on both type of magmas, giving pressure for fluid saturation up to 4 kbar for the volatile-rich yellow pumice (Métrich *et al.*, 2001; Bertagnini *et al.*, 2003). Temperature constraints come from melt inclusion homogenisation during heating stage experiments which indicate that the yellow

pumice was at 1125-1140°C, whereas the crystal-rich scoria was at a slightly lower temperature of 1100-1125°C. The pre-eruptive temperature for the yellow pumice has been confirmed by recent phase equilibrium work (Di Carlo *et al.*, 2004). Here we use temperatures of 1140 and 1100°C for the pumice and scoria rocks, respectively. The redox state of Stromboli magmas was inferred through the determination of sulphur speciation (Métrich and Clocchiatti, 1996; Métrich *et al.*, 2002) which yields an average fO_2 of NNO+0.95. For the calculation of fluid phase corresponding to the scoria or degassed magmas, we assumed that the pre-eruptive melt contained 40-50 ppm CO_2 , since this species could not be detected by FTIR in scoria melt inclusions.

4.2. Vesuvius

For Vesuvius there is a large body of data on melt inclusion chemistry (*e.g.*, Vagelli *et al.*, 1992, 1993; Cioni *et al.*, 1995, 1998; Marianelli *et al.*, 1995, 1999; Belkin *et al.*, 1998; Lima *et al.*, 1999; Signorelli and Capaccioni, 1999; Signorelli *et al.*, 1999; Cioni, 2000; Raia *et al.*, 2000; Webster *et al.*, 2001), but for Stromboli only recently has the CO_2 in melt inclusions been determined using micro beam techniques (Marianelli *et al.*, 1999; Signorelli *et al.*, 1999; Cioni, 2000). We did not consider works done on felsic phonolites, however, since eq. (2.1) is not yet calibrated for these compositions. To minimise errors arising from differences in sulphur and water determinations between different research groups, we used only the data sets of Cioni *et al.* (1995), and Marianelli *et al.* (1995, 1999) in which at least both S and H_2O were determined using the same instrument and analytical procedure. Most of the analysed inclusions correspond to primitive melts for which heating stages experiments give homogenisation temperatures of *ca.* 1100-1150°C (*e.g.*, Cioni *et al.*, 1995; Marianelli *et al.*, 1995). For melt inclusions lacking such information, temperatures were determined using the CaO in melt geothermometer of Cioni *et al.* (1998). The redox state is known indirectly from sulphur speciation determination either on bulk

rocks, lava or pumices (Marini *et al.*, 1998), or in melt inclusions (Métrich *et al.*, 2002), both yielding an fO_2 of *ca.* NNO+1.2.

4.3. Vulcano

Melt inclusions of Vulcano magmas have been analysed by Clocchiatti *et al.* (1994a,b) and Gioncada *et al.* (1998). The latter work showed that the CO_2 content of primitive melt inclusions is below the FTIR detection limit, which yields a maximum CO_2 content of about 50 ppm. The storage conditions of the reservoir that fed the 1888-1890 eruption, during which rhyolite to trachytic magmas were ejected, were inferred by Clocchiatti *et al.* (1994a) to be: a pressure depth of 80 MPa, melt water contents in the range 1-1.5 wt%, and temperatures between 1000-1100°C. This shallow reservoir is inferred to be connected to a deeper one where intermediate and mafic magmas reside. These are assumed to be equivalent to primitive melt inclusions in La Sommata basalts, whose homogenisation temperature is $1180 \pm 20^\circ C$ (Gioncada *et al.*, 1998). The depth of the felsic upper reservoir provides an upper bound for the mafic part, which is estimated to lie at 70-110 MPa (Gioncada *et al.*, 1998). Measured melt water contents of the least differentiated magmas (La Sommata melt inclusions) are in the range 2.1-3.8 wt%, that is they are higher than their potential felsic derivatives (Clocchiatti *et al.*, 1994a; Gioncada *et al.*, 1998), suggesting an open system behaviour with respect to volatiles (Clocchiatti *et al.*, 1994b). A similar observation can be made for sulphur which reaches maximum values of 2872 ppm in the most mafic melts analysed (Clocchiatti *et al.*, 1994b). The redox state of mafic magmas has been inferred from sulphur speciation to be at NNO+0.72 (Métrich and Clocchiatti, 1996; Métrich *et al.*, 2002).

To calculate the fluid phase compositions we used the melt inclusions analyses reported by Clocchiatti *et al.* (1994a) assuming that they all contained *ca.* 40-50 ppm dissolved CO_2 . Thus, the calculated fluid compositions correspond to the maximum CO_2 content that can be expected at Vulcano given the available melt

inclusion constraints. In addition, we calculated the fS_2 corresponding to melt inclusions of Clocchiatti *et al.* (1994b), by assuming that mafic melts have a pre-eruptive water content of 1 wt%, whereas that of more felsic compositions (shoshonite to latite-rhyolites) has been set at 1 wt%.

4.4. Etna

Unlike the three previous volcanoes, there are currently no published combined sets of pre-eruptive H₂O-CO₂-S contents of Etna mafic melts. Existing melt inclusion studies have analysed either S and Cl (*e.g.*, Clocchiatti and Métrich, 1984), H₂O and S (Métrich *et al.*, 1993), CO₂ (Métrich and Mosbah, 1988), or S and Cl (Kamenetsky and Clocchiatti, 1996) but none has determined the concentration of those volatiles collectively in single melt inclusions. A literature survey shows that only the alkali basalt erupted in 1892 (Mt. Maletto), which is among the most primitive basalts yet erupted on Etna (Armienti *et al.*, 1988), has had its pre-eruptive volatile content determined in full, albeit not necessarily on the same melt inclusions (Métrich and Mosbah, 1988 for CO₂; and Métrich *et al.*, 1993, for H₂O and S). There are thus surprisingly very few constraints on the volatile concentrations that could be used to determine fluid saturation pressures of Etnaean magmas, either past or present. Therefore, we adopted a different strategy relative to the three other volcanoes. We performed thermodynamic calculations at various pressures, assuming that a fluid phase is present, using available H₂O and S melt concentrations coupled to petrological T and fO_2 constraints. The pressure range over which calculations were performed is derived from geophysical and petrological constraints, as summarised in the next sections.

The melt water contents of alkali basalts of Etna have been determined to be in the range 1-2.3 wt% (Trigila *et al.*, 1990; Métrich *et al.*, 1993), the higher value being regarded as the pristine pre-eruptive melt water content, based on phase equilibrium constraints (Métrich and Rutherford, 1998). Geochemical modelling has shown that these alkali basalts can be derived

by 12-13 wt% of olivine fractionation from a picritic basalt (Armienti *et al.*, 1988), which therefore suggests that the water content of primary magmas at Etna is close to 0.8-2 wt%, assuming closed system fractionation. As stated above, only melt inclusions of the 1892 eruption have had their CO₂ content determined, with an average value of 588 ppm (Métrich and Mosbah, 1988). This value is considered as a minimum because of the occurrence of carbonate crystals in the quenched melt inclusions (Métrich and Mosbah, 1988). Additional evidence for magmatic CO₂ comes from its occurrence in the volcanic plume or in diffuse soil emanations (*e.g.*, Allard *et al.*, 1991; Bruno *et al.*, 2001), and from the study of fluid inclusions associated to melt inclusions in lava phenocrysts or ultramafic nodules (Clocchiatti and Métrich, 1984; Sobolev *et al.*, 1991; Frezzotti *et al.*, 1991; Clocchiatti *et al.*, 1992). Such fluid inclusions generally belong either to a low (H₂O-CO₂ or CO₂) or a high (CO₂) density population. The high density inclusions yield maximum entrapment pressures of 6-7 kbar of a nearly pure CO₂ fluid (Clocchiatti *et al.*, 1992), whereas the low one yields pressures in the range 1-3 kbar (Clocchiatti and Métrich, 1984; Sobolev *et al.*, 1991; Frezzotti *et al.*, 1991). These observations suggest that any fluid phase coexisting with magma at depth must be CO₂-rich.

For sulphur there is a considerable amount of analytical data (*e.g.*, Métrich and Clocchiatti, 1989; Métrich *et al.*, 1993), the highest concentration recorded in melt inclusions being 3800 ppm (as quoted in Clocchiatti *et al.*, 1992), though most have sulphur contents in the range 2500-3500 ppm (Clocchiatti and Métrich, 1984; Métrich and Clocchiatti, 1989, 1996). Generally the sulphur-rich melt inclusions are enclosed in the most magnesian olivines and they do not coexist with immiscible sulphide liquids (Métrich and Clocchiatti, 1989), while sulphur-poor inclusions hosted by Fe-rich olivine are occasionally saturated in a Cu-rich sulphide melt (Métrich and Clocchiatti, 1989).

Petrological, geochemical, and geophysical evidence (*e.g.*, Armienti *et al.*, 1988; Bonaccorso, 1996, 2001; Tanguy *et al.*, 1997; Murru *et al.*, 1999) all points to the probable existence of a

large deep reservoir beneath the volcano at a pressure depth range of 7-10 kbar, in which most of the primitive Etnean magmas stagnate and fractionate to hawaiites (or trachybasalts). Such hawaiites constitute the bulk of the magmas outpoured over the last 300000 years. Similarly, geophysical, geodetic, geochemical, melt/fluid inclusions, and phase equilibrium constraints, suggest that there are at least two additional shallow reservoirs feeding historical as well as on-going eruptions that lie at a pressure depth of about 1 and 3-4 kbar (*e.g.*, Sobolev *et al.*, 1991; Bonnacorso, 1996; Métrich and Rutherford, 1998; Murru *et al.*, 1999; Caracausi *et al.*, 2003). Temperatures of magma extrusion for the recent period have been estimated from a variety of approaches including direct measurement of lava flow (Tanguy *et al.*, 1997), melt inclusion homogenisation (*e.g.*, Sobolev *et al.*, 1991; Clocchiatti *et al.*, 1992) and thermodynamical and phase equilibrium constraints (Trigila *et al.*, 1990; Métrich and Rutherford, 1998) and all fall within a T range of 1070-1100°C. Temperatures of primitive alkali magmas stored at 7-10 kbar are estimated to be around 1200°C (Kamenetsky and Clocchiatti, 1996).

Redox conditions of Etnean basalts have been determined either through spinel-olivine-orthopyroxene equilibria (*e.g.*, Kamenetsky and Clocchiatti, 1996) or via the determination of sulphur speciation in glass inclusions (Métrich and Clocchiatti, 1996). The latter approach gives an fO_2 corresponding to NNO+0.35. The fO_2 retrieved from spinel bearing assemblages in primitive melt inclusions from Mt. Maletto alkali basalt encompasses a range NNO up to NNO+1, but we note that the fO_2 calculated for another similar alkali basalt (Mt. Spagnolo) falls between NNO+1 to NNO+2, suggesting possibly more oxidizing conditions than those adopted here. Sato and Moore (1973) made intrinsic fO_2 (and fS_2) measurements in gas streams from two hornito vents during the 1970 eruption and they found an fO_2 of NNO-0.45, which Gerlach (1980) showed to be compatible with the restored gas compositions measured by Huntingdon (1973) during the same event.

In summary, because of the lack of confident pre-eruptive CO_2 determinations, thermodynamic calculations of fluid composition co-

existing with Etnean mafic magmas were performed in the pressure range 400-10000 bar, in keeping with the petrological and geophysical constraints summarised above. Temperatures were either 1150°C (1 wt% H_2O) or 1130°C (2.23 wt% H_2O), based on melt inclusion heating stage experiments (Métrich and Clocchiatti, 1996). We considered melts water contents of 1 and 2.23 wt% (Métrich *et al.*, 1993) and melt sulphur contents of 1000-3400 ppm, with melt compositions as reported by Métrich *et al.* (1993). We explored fO_2 conditions from NNO-0.45 to NNO+0.35, to ascertain the effect of fO_2 on both fS_2 and fluid composition.

5. Trends in fS_2 and sulphide saturation

In all subsequent diagrams we use the ratio CaO/Al_2O_3 as a differentiation index to illustrate variations in fS_2 . The sulphide saturation is computed from the following equilibrium:



using thermodynamic data for liquid FeO and FeS as given in O'Neill and Mavrogenes (2002). For all melt compositions we assume that 90% of iron is FeO, and we use an activity coefficient for FeO of 1.4 (O'Neill and Mavrogenes, 2002). Given the relatively high redox state of the magmas, there may be less Fe^{2+} than the assumed value and thus the activity of FeS calculated (a_{FeS}) here should be considered as maximum values. The calculated a_{FeS} corresponds to pure FeS, however. Saturation in a sulphide liquid could arise if other components are present in solution, such as Cu or Ni.

5.1. Stromboli

In fig. 3, two trends in fS_2 can be distinguished; MI of pumices define a flat trend with fS_2 values in the range 0.01 to 1 bar, the most primitive melt having the highest fS_2 . The pumice trend joins that defined by the scoria MI, which is much more steep and spreads over nine orders of magnitude fS_2 at a nearly constant CaO/Al_2O_3 ratio. This spread in fS_2 re-




Fig. 3. Evolution of fS_2 with CaO/Al_2O_3 ratio of Stromboli magmas. Because of massive clinopyroxene crystallisation in those magmas, fractionation is marked by a continuous decrease in the CaO/Al_2O_3 ratio. The fS_2 is calculated from melt inclusion data (Métrich *et al.*, 2001) and eq. (2.1) (see text). Pumice corresponds to magma emitted during paroxysmal events whereas scoria represents magmas erupted during Strombolian type eruptions.




Fig. 4. Calculated activities of liquid FeS plotted versus the CaO/Al_2O_3 ratio for the four volcanoes considered (see text for explanations). Note that almost all magmas are below saturation conditions with an immiscible sulphide liquid.

flects mostly the variable concentrations of sulphur analysed in various melt inclusions, since melt compositions in scoria, including water content, varies little (Métrich *et al.*, 2001). The huge spread records the progressive degassing in sulphur of the magma that is stored in the shallow reservoir. The very steep

slope indicates that degassing operates with little chemical modification of the environment. Inclusions in the scoria displaying the highest fS_2 may represent remnants of deep magma batches that intrude the upper reservoir or record incipient degassing of the deep magma batches during uprise. Melt inclusions in pumices were trapped at depths exceeding 3 kbar (see below). The fS_2 variations in this deeper reservoir appear to be much more restricted. The slight decrease observed corresponds to either selective loss of sulphur or, and more probably, to an increase in melt water content during fractionation, since water decrease fS_2 at a given melt sulphur content (see eq. (2.1)). The calculated a_{FeS} ranges from 0.1 down to 0.0001 (fig. 4), suggesting that mafic melts at Stromboli are far from reaching sulphide saturation. Removal of sulphur via sulphide disposal during fractionation is therefore probably not an efficient process at Stromboli.

5.2. Vesuvius

The data on Vesuvius are shown in fig. 5, being distinguished by the year of eruption. Two main trends emerge. The first corresponds to the 1906-1944 eruptions, whereas the second corresponds to the Pollena-Pompeii-Avellino events. The 1906-1944 trend is somewhat analogous to the Stromboli in that less differentiated MI have higher and broadly constant fS_2 which, below a CaO/Al_2O_3 ratio of 0.7, sharply decrease down to a $\log fS_2$ of -7 . In contrast, MI analysed in Pollena-Pompeii-Avellino define a broad trend of increasing fS_2 as fractionation proceeds, although for the Pompeii MI a significant scatter is apparent, reaching values of fS_2 in excess of 100 bar. It is interesting to note that the first trend corresponds to events which characterise open conduit conditions, with semi-persistent Strombolian activity, as opposed to the other events which typify the establishment of a closed system following obstruction of the conduit during a long repose time preceding Plinian to sub-Plinian eruptions (*e.g.*, Civetta *et al.*, 1991; Civetta and Santacroce, 1992; Cioni *et al.*, 1998). On this basis, the trend of decreasing fS_2 can be interpreted as reflecting




Fig. 5. Evolution of fS_2 with CaO/Al_2O_3 ratio of Vesuvius magmas. The fS_2 is calculated from melt inclusion data (Cioni *et al.*, 1995; Marianelli *et al.*, 1995, 1999) and eq. (2.1) (see text). Different symbols correspond to the main eruptive events. The Pollena, Pompei and Avellino eruptions correspond to closed conduit conditions, while the 1944 and 1906 events are characteristic of open conduit conditions.

continuous loss of sulphur (and other volatiles) during magma uprise and emission to the surface (open conduit), in a way similar to that seen at Stromboli, whereas that of increasing fS_2 reflects a closed system in which sulphur has perhaps a dominant incompatible behaviour (its concentrations in the melt and fluid increase). As for Stromboli, all but one calculated a_{FeS} are significantly below unity (fig. 4) which again suggests that conditions for sulphide saturation are not reached in Vesuvius magmas, although they tend to approach it under closed conduit conditions. This may be one reason for the observed increase in fS_2 during differentiation.

5.3. Vulcano

The Vulcano data define two distinct groups (fig. 6); one corresponding to the most mafic MI (basaltic compositions) which cluster at fS_2 around 0.1 bar, and the second that corresponds to more evolved MI (shoshonitic to rhyolite compositions) that span values of fS_2 between several bars down to 10^{-8} bar, defining a trend negatively correlated with the CaO/Al_2O_3 ratio (that is fS_2 increases with differentiation). This

second trend is similar to that observed for the Pollena-Pompei-Avellino magmas discussed above. However, there is a significant gap between mafic and felsic magmas and it is unclear from the data whether this gap is real or if the fS_2 of the two groups overlap at a CaO/Al_2O_3 ratio of around 1. If this were true, then mafic melts must evolve towards lower fS_2 during fractionation. This hypothetical decrease could represent decompression of volatile-rich magmas during open conduit conditions, such as at Stromboli. Thereafter, once the partially degassed mafic batches have been emplaced at shallow level they further fractionate under closed system conditions during which fS_2 increases. Alternatively, the two groups may represent different levels of magma fractionation, one deep (both high and constant CaO/Al_2O_3 ratio and fS_2) and one shallow (increasing fS_2 with fractionation). The a_{FeS} calculated are in general much lower than unity ($a_{FeS} < 0.01$), but they do increase with fractionation such that the most felsic magmas may approach FeS saturation (fig. 4).




Fig. 6. Evolution of fS_2 with CaO/Al_2O_3 ratio of Vulcano magmas. The fS_2 is calculated from melt inclusion data (Clocchiatti *et al.*, 1994a,b; Giocada *et al.*, 1998) and eq. (2.1) (see text). Note that the most primitive melts (high CaO/Al_2O_3 ratio) cluster at around fS_2 of 0.1 bar, whereas the rest of melts defines a negative trend of increasing fS_2 with differentiation.




Fig. 7. Evolution of fS_2 with CaO/Al_2O_3 ratio of Etna magmas. The fS_2 is calculated from melt inclusion data (Métrich *et al.*, 1993) and eq. (2.1) (see text).

5.4. Etna

The Etna data (fig. 7) do not define any obvious trend, but this is largely due to the low number of MI analysed so far. The most primitive MI again have fS_2 in the range 0.1 to 1 bar, as observed for the other volcanoes. The calculated a_{FeS} values are the highest of the four volcanoes, ranging from 0.1 up to 0.95, or close to FeS saturation (fig. 4).

6. Composition of fluids

The compositions of fluids calculated for MI with known H_2O , S and CO_2 contents are listed in table II together with input fH_2 , fS_2 and fH_2O and the corresponding pressure of fluid saturation. In all cases, the computed mole fraction of CH_4 is several orders of magnitude lower than that of either CO_2 or CO , and this species is not considered further in this work.

6.1. Stromboli

The pressures of fluid saturation of the MI in pumices range from 2100 bar to near 3500 bar (table II, fig. 8), *i.e.* they are comparable to those calculated by Métrich *et al.* (2001) and Bertagnini *et al.* (2003), despite the fact that sulphur bearing species were not taken into account in these

works. This is due to the fact that, although melt sulphur contents are relatively high, the corresponding fS_2 , and thus the molar abundances of S-bearing species, remain low. In the calculated examples, the mole fractions of SO_2+H_2S never exceeds 0.1 (fig. 8). The mole fraction of H_2O remains relatively constant, at about 0.3, while that of CO_2 slightly increases with pressure, exceeding 0.6 at near 3500 bar. The fluid coexisting with MI in scoria, calculated with the assumption of 50 ppm CO_2 dissolved, corresponds to entrapment pressures of 100 bar, which is equivalent to a depth beneath the crater of *ca.* 300 m, *i.e.* similar to previous estimates (*e.g.*, Harris and Stevenson, 1997). Given that the central part of the volcano is permanently flushed with CO_2 -rich gases, as remote sensing of the volcanic plume indicates (Allard *et al.*, 1994), it is unlikely that magmas stored in the shallow reservoir are CO_2 -free. Therefore the 50 ppm threshold appears to be a limiting condition which corresponds to the maximum possible pressure depth at which the upper reservoir that feeds the typical Strombolian activity is lying. Deeper conditions will inevitably produce MI with detectable CO_2 contents with the FTIR method. Under these conditions it appears that the fluid phase is extremely rich in CO_2 , with X_{CO_2} in excess of 0.9 (fig. 8). In contrast, the water content of the shallow magmatic fluid is very low being even lower than SO_2 in some instances.

Figure 9 shows the evolution of both $H_2O/ /SO_{2tot}$ and CO_2/SO_{2tot} ($SO_{2tot}=SO_2+H_2S$) mole ratios with pressure. MI of pumices define a broad positive trend in both ratios, whose extrapolation to near surface conditions corresponds to gases with $H_2O/ /SO_{2tot}$ and CO_2/SO_{2tot} ratios between 0.1 and 1. Such ratios are at the lower end of the range displayed by fluids coexisting with scoria MI. In these, the $H_2O/ /SO_{2tot}$ can exceed 10 while the CO_2/SO_{2tot} can go over 1000. Clearly such a large spread is chiefly due to the variable contents of sulphur dissolved in MI in scoria. The fact that the pumice MI trend intersects the lower end of the scoria trend suggests that this ratio range of 0.1-1 represents the fluid compositions of magmas just injected in the upper reservoir, which have not yet undergone extensive volatile degassing associated to normal Strombolian activity.

Table II. Composition of fluids of Stromboli, Vesuvius, Vulcano and Etna volcanoes, based on melt inclusion constraints.

	<i>P</i> bar	<i>T</i> °C	Δ NO	Δ FFS	<i>f</i> H ₂ bar	<i>f</i> S ₂ bar	<i>f</i> H ₂ O bar	XH ₂	XH ₂ O	XCO ₂	XCO	XSO ₂	XH ₂ S	XS ₂	H ₂ O _n wt%	S _m ppm	CO _{2-m} ppm	S _{fluid} wt%
Stromboli																		
ST82p-ohn52a	2930	1150	0.95	4.597	2.100	0.065	900	0.00044	0.34311	0.64264	0.00307	0.01000	0.00069	0.00001	2.7	1730	1087	0.978
ST82p-ohn52a	2900	1150	0.95	4.569	2.300	0.062	960	0.00048	0.36716	0.61933	0.00299	0.00929	0.00074	0.00001	2.8	1730	1031	0.931
ST82p-ohn9b	3430	1150	0.95	4.123	2.400	0.020	1020	0.00047	0.32820	0.66379	0.00324	0.00402	0.00035	0.00000	2.8	2050	1689	0.3949
ST82p-ohn50	2570	1150	0.95	3.989	2.000	0.017	865	0.00044	0.37623	0.61287	0.00271	0.00734	0.00038	0.00001	2.7	1230	894	0.721
ST79p-ohn30	2190	1100	0.95	5.793	1.600	0.180	660	0.00050	0.33507	0.59010	0.00294	0.06767	0.00036	0.00000	2.1	2110	1107	6.283
ST82s-Ohn2	100	1100	0.95	3.465	0.005	0.003	2	0.00005	0.02053	0.89200	0.00461	0.08275	0.00002	0.00003	0.15	740	50	5.885
ST82s-Ohn3	100	1100	0.95	-2.392	0.005	0.000	2	0.00005	0.02060	0.97385	0.00504	0.00046	0.00000	0.00000	0.16	50	50	0.0338
ST82s-Ohn4	100	1100	0.95	2.028	0.005	0.003	2	0.00005	0.02058	0.94935	0.00491	0.02510	0.00001	0.00000	0.16	890	50	1.83
Vesuvius																		
V597-109-O1	4000	1150	1.2	5.140	1.700	0.198	1100	0.00023	0.29534	0.67997	0.00223	0.02157	0.00063	0.00003	2.9	1800	2500	1.931
V597-109-O1	3300	1150	1.2	4.698	1.350	0.078	810	0.00024	0.27594	0.70327	0.00251	0.01760	0.00042	0.00001	2.5	1600	2100	1.555
V597-109-O1	3220	1150	1.2	4.776	1.200	0.094	700	0.00023	0.24751	0.72945	0.00273	0.01963	0.00043	0.00000	2.3	1700	1900	1.6969
V597-109-O1	5250	1150	1.2	4.327	2.600	0.023	1650	0.00022	0.31140	0.68238	0.00228	0.00351	0.00020	0.00000	3.4	2300	3100	0.3711
V597-109-O1	6400	1150	1.2	3.566	2.600	0.003	1680	0.00016	0.24063	0.75557	0.00268	0.00079	0.00005	0.00000	3.2	1000	3500	0.0015
V597-109-D1	3520	1150	1.2	4.069	1.600	0.017	930	0.00026	0.29380	0.69639	0.00254	0.00678	0.00021	0.00000	2.66	1200	1900	0.615
V597-109-D1	5360	1150	1.2	4.830	1.470	0.071	920	0.00012	0.16591	0.82509	0.00317	0.00552	0.00019	0.00001	2.4	1600	4300	0.4608
V51906-O1	3100	1150	1.2	4.974	1.900	0.151	1100	0.00036	0.38545	0.58675	0.00201	0.02454	0.00087	0.00003	3	1700	1800	2.3677
V51906-O1	3300	1150	1.2	4.448	2.150	0.044	1220	0.00037	0.39883	0.58727	0.00203	0.01102	0.00048	0.00001	3.2	1800	1700	1.0898
GL3-3-Lc	900	1150	1.2	1.164	0.270	0.000	150	0.00026	0.18988	0.80404	0.00325	0.00257	0.00001	0.00000	1.15	300	383	0.212
Vulcano																		
IV/O131	885	1180	0.72	3.961	2.650	0.038	815	0.00243	0.03510	0.00021	0.00021	0.02951	0.00024	0.00004	2.88	2300	48	5.0374
IVol4	725	1180	0.72	4.605	2.150	0.170	630	0.00251	0.87752	0.04119	0.00028	0.07325	0.00505	0.00020	2.51	1980	39	11.19
6FM-M Rosso	185	1160	0.72	0.509	0.290	0.000	85	0.00150	0.46785	0.52443	0.00405	0.00215	0.00002	0.00002	0.93	200	51	0.2188
Tuffi Bruni inf	400	1100	0.72	0.961	0.420	0.000	130	0.00180	0.60220	0.39227	0.00250	0.00119	0.00004	0.00000	1.160	440	46	0.1396
Tuffi Bruni sup	400	1100	0.72	5.673	0.730	0.500	228	0.00165	0.58339	0.25893	0.00158	0.14457	0.00074	0.00014	1.500	730	53	15.774
Vulcanello	210	1100	0.72	2.715	0.420	0.001	130	0.00172	0.57700	0.40978	0.00262	0.00859	0.00030	0.00000	1.150	650	50	0.980
Peccite rhyolite	220	1000	0.72	5.009	0.330	0.014	109	0.00140	0.51130	0.46300	0.00222	0.01971	0.00231	0.00006	1.000	100	55	2.286
Etna																		
I wt% H ₂ O in melt																		
E79043	400	1150	-0.45	6.654	1.450	12.030	110	0.00334	0.28634	0.53668	0.01570	0.06513	0.06476	0.02805	1.06	3400	114	15.91
E79043	1000	1150	-0.45	6.677	1.600	11.330	125	0.00137	0.14253	0.77273	0.02249	0.02405	0.02702	0.00981	1.06	3400	423	5.610
E79043	500	1100	0.35	6.733	0.550	5.590	110	0.00099	0.23352	0.57256	0.00573	0.15907	0.01788	0.01026	1.06	3400	153	15.41
E79043	400	1150	0.35	6.552	0.560	9.515	110	0.00128	0.27834	0.27919	0.00310	0.39352	0.02244	0.02216	1.06	3400	58	32.91
E79043	500	1150	0.35	6.549	0.550	9.267	110	0.00099	0.22679	0.41710	0.00457	0.16200	0.01731	0.01700	1.06	3400	110	26.44
E79043	1000	1150	0.35	6.407	0.630	6.091	125	0.00054	0.14055	0.72342	0.00821	0.11421	0.00782	0.00053	1.06	3400	394	9.979
E79043	3000	1150	0.35	6.341	0.680	3.593	145	0.00014	0.05645	0.91146	0.01048	0.01903	0.00169	0.00074	1.06	3400	1774	1.6621
E79043	7000	1150	0.35	6.209	0.880	1.250	220	0.00002	0.02131	0.96598	0.00981	0.00271	0.00011	0.00005	1.06	3400	5404	0.216
E79043	10000	1150	0.35	6.183	1.080	0.454	300	0.00002	0.01557	0.97320	0.01050	0.00066	0.00003	0.00001	1.06	3400	8484	0.051
E79043	1000	1150	0.35	4.067	0.630	0.023	125	0.00054	0.14485	0.83820	0.00961	0.00635	0.00044	0.00002	1.06	1000	457	0.598
2.23 wt% H₂O in melt																		
1892-OH11-8	500	1130	0.35	5.431	2.300	0.490	450	0.00410	0.91339	0.00725	0.00007	0.05635	0.01799	0.00088	2.120	3310	2	11.570
1892-OH11-8	1000	1130	0.35	5.414	2.550	0.430	520	0.00210	0.55216	0.40819	0.00379	0.02476	0.00875	0.00034	2.230	3310	211	3.657
1892-OH11-8	3000	1130	0.35	5.348	2.900	0.250	635	0.00060	0.24644	0.73937	0.00710	0.00434	0.00210	0.00005	2.230	3310	1318	0.558
1892-OH11-8	7000	1130	0.35	5.216	3.700	0.087	950	0.00010	0.09270	0.89730	0.00854	0.00108	0.00024	0.00001	2.230	3310	5056	0.104
1892-OH11-8	10000	1130	0.35	5.117	4.600	0.039	1300	0.00007	0.06770	0.92257	0.00944	0.00017	0.00004	0.00000	2.230	3310	8111	0.016
1892-OH11-8	600	1130	-0.45	5.824	6.550	1.190	500	0.00094	0.84808	0.06289	0.00153	0.01080	0.00656	0.00173	2.230	3310	18	12.079
1892-OH11-8	1000	1130	-0.45	5.811	6.500	1.070	520	0.00514	0.55087	0.39255	0.00932	0.00602	0.00322	0.00086	2.230	3310	204	4.720
1892-OH11-8	1000	1130	0.35	3.053	2.550	0.002	520	0.00201	0.55485	0.43689	0.00406	0.00162	0.00057	0.00000	2.230	1000	227	0.239

Sources of melt inclusion volatiles are: Stromboli (Métrich *et al.*, 2001); Vesuvius (Marianelli *et al.*, 1999); Vulcano (Clocchiatti *et al.*, 1994a,b; Giocciada *et al.*, 1998); Etna (Métrich *et al.*, 1993).




Fig. 8. Composition of fluids in equilibrium with melt inclusions from scoria and pumice (Métrich *et al.*, 2001). X_i is the mole fraction of species i . Symbols plotting at 100 bar correspond to melt inclusions in scoria, all other to melt inclusions in pumice. The compositions of fluids in scoria were calculated assuming a maximum CO_2 content of 50 ppm.




Fig. 9. Pressure evolution of $\text{H}_2\text{O}/\text{SO}_{2\text{tot}}$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios of fluids in equilibrium with melt inclusion of Stromboli melt inclusions. The thick grey continuous and dashed lines represent average C/S and H/S ratio measured on a hot fumerole or within the dilute plume at Stromboli (Allard *et al.*, 1994).

6.2. Vesuvius

The mole fractions of CO_2 , H_2O and $\text{SO}_{2\text{tot}}$ of the 1944 eruption are plotted against pressures of fluid saturation on fig. 10. Calculated pressures range from slightly below 1000 bar to over 6500 bar, being again similar to those cal-

culated by Marianelli *et al.* (1999). There is no clear trend with pressure, but it is interesting to note that even relatively low pressure MI yield fluid compositions very rich in CO_2 . In all computed examples, the SO_2 and H_2S mole fractions remain below 0.05, whereas the mole fraction of H_2O ranges between 0.2 up to 0.4.

The $\text{H}_2\text{O}/\text{SO}_{2\text{tot}}$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios decrease with pressure down to values of about 20 and 50, respectively, at 3000 bar (fig. 11), and then increase again with further pressure decrease. The second part of the evolution remains hypothetical, however, since it is based on a single MI datum at 900 bar.




Fig. 10. Composition of fluids in equilibrium with melt inclusions of the 1944 Vesuvius eruption (Marianelli *et al.*, 1999). X_i is the mole fraction of species i . Note the rather constant fluid composition over the pressure interval.




Fig. 11. Pressure evolution of $\text{H}_2\text{O}/\text{SO}_{2\text{tot}}$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios of fluids in equilibrium with melt inclusion of the 1944 Vesuvius eruption.




Fig. 12. Composition of fluids in equilibrium with melt inclusions of Vulcano magmas (Clocchiatti *et al.*, 1994a,b; Gioncada *et al.*, 1998). X_i is the mole fraction of species i . Fluid compositions have been calculated assuming that the melt inclusions contain 50 ppm dissolved CO_2 . This gives maximum CO_2 contents of the fluid. Note the continuous evolution of the fluid composition with pressure.




Fig. 13. Pressure evolution of $\text{H}_2\text{O}/\text{SO}_{2\text{tot}}$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios of fluids in equilibrium with melt inclusions of Vulcano magmas.

6.3. Vulcano

The fluid composition of Vulcano magmas is shown in fig. 12. Given the lack of detectable CO_2 , and the generally low melt H_2O contents (<3 wt%), the calculated pressures for fluid saturation are low compared to Stromboli and Vesuvius, ranging from 900 down to 200 bar, in agreement with previous findings (Clocchiatti *et al.*, 1994a; Gioncada *et al.*, 1998). The data set shows a regular variation of fluid composition as

pressure decreases. Between 900 and 200 bar, the mole fraction of H_2O decreases continuously, from over 0.9 down to 0.5, being paralleled by a concomitant increase in CO_2 . The mole fraction of $\text{SO}_{2\text{tot}}$ first increases up to 400 bar, where it reaches a value of 0.15, and then decreases down to 0.05 at 200 bar.

The evolution of the $\text{H}_2\text{O}/\text{CO}_{2\text{tot}}$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios with pressure is shown on fig. 13. The $\text{H}_2\text{O}/\text{CO}_{2\text{tot}}$ decreases continuously with pressure, reaching values of 1 at 200 bar. In contrast, the $\text{CO}_2/\text{SO}_{2\text{tot}}$ ratio increases with decreasing pressures, from 1 at 900 bar up to 400 at 200 bar: note that this ratio displays considerable variation at 200 bar, as observed for Stromboli. This reflects variable depletion of sulphur during shallow level degassing of the magmas

6.4. Etna

Figure 14 shows the evolution of the mole fractions of H_2O , CO_2 and S_{tot} with pressure calculated for melt water contents of 1 and 2.23 wt%, f_{O_2} at NNO+0.35, and melt sulphur contents of 3300-3400 ppm. We stress again that calculations assume the presence of a fluid phase in the entire pressure range. In both cases, the fluid composition is dominated by CO_2 at pressures higher than 2000 bar. At pressures higher than 7000 bar, the fluid consists of more than 90% of CO_2 . For instance at 10000 bar, the calculated fluid composition corresponding to 1 wt% dissolved H_2O has a mole fraction of CO_2 of 0.97, that is the fluid is nearly pure CO_2 , in agreement with fluid inclusion constraints on primitive Etna basalts (Clocchiatti *et al.*, 1992). Under such conditions the calculated CO_2 content of the basalt melt is 0.85 wt% (table II). It is only below 1000 bar (2.23 wt% H_2O) or 400 bar (1 wt% H_2O) that CO_2 is lower than H_2O . For water-rich conditions, the mole fraction of H_2O always exceeds that of $\text{SO}_{2\text{tot}}$, while at a melt H_2O content of 1 wt%, $\text{SO}_{2\text{tot}}$ equals or even exceeds H_2O at pressures lower than 500 bar.

The difference in fluid composition arising from different melt H_2O contents is illustrated on fig. 15 where the mole fraction of H_2O is plotted

against that of CO_2 . Note that the mole fractions plotted correspond to actual value that take into account the sulphur species, whose total mole fraction reach values in excess of 0.44 (fig. 14a,b, table II). This is why all the data plot below the mixing line corresponding to a binary $\text{H}_2\text{O}-\text{CO}_2$. Clearly, the amount of dissolved water significantly affects the fluid composition: at 500 bar the calculated mole fraction of H_2O varies from 0.92 for a melt H_2O content of 2.2 wt%, down to 0.22 for a melt H_2O content of 1 wt%. However, the effect of melt H_2O content vanishes at high pressure, so that at or above 7000 bar, the calculated fluid compositions for the two melt water contents diverge by less than 0.05 in their mole fraction of H_2O . Variation in $f\text{O}_2$ in the range considered, does not strongly




Fig. 14a,b. Evolution of the mole fractions of H_2O , CO_2 and $\text{H}_2\text{S}+\text{SO}_2$ with pressure calculated for melt water contents of 1 (a) and 2.23 (b) wt%, $f\text{O}_2$ at NNO+0.35, and melt sulphur contents of 3300-3400 ppm, corresponding to primitive Etnean magmas. X_i is the mole fraction of species i .




Fig. 15. Evolution of the mole fraction of H_2O against that of CO_2 calculated for melt water contents of 1 and 2.23 wt%, $f\text{O}_2$ at NNO+0.35, and melt sulphur contents of 3300-3400 ppm, corresponding to primitive Etnean magmas. The mole fractions correspond to actual values, that take into account the sulphur species, whose total mole fraction reach values in excess of 0.44 (fig. 14a,b, table II). This is why all the data plot below a mixing line corresponding to a binary $\text{H}_2\text{O}-\text{CO}_2$. Dashed lines represent isobaric conditions calculated for different melt water contents.




Fig. 16. Pressure evolution of the $\text{H}_2\text{O}/\text{CO}_2$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios for melts having either 1 or 2.23 wt% dissolved H_2O and 3300-3400 ppm of dissolved sulphur at NNO+0.35.

affect the fluid compositions, apart from the $\text{SO}_2/\text{H}_2\text{S}$ ratio. In contrast, a lower sulphur content of the melt (that is lower $f\text{S}_2$) obviously decreases significantly the abundances of sulphur bearing species and, by implication, increases the CO_2/SO_2 ratio (table II).

Figure 16 shows the dramatic effect that pressure exerts on the $\text{H}_2\text{O}/\text{CO}_2$ and $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratios. The $\text{H}_2\text{O}/\text{CO}_2$ ratio increases from 0.01-0.1 at 10000 bar to 1-100 at 400-500 bar, whereas the $\text{CO}_2/\text{SO}_{2\text{tot}}$ decreases within this pressure interval from 1000 at 10000 bar down to 0.8-0.1 at 400-500 bar. The variation in melt water contents affects the $\text{H}_2\text{O}/\text{CO}_2$ ratio more than the $\text{CO}_2/\text{SO}_{2\text{tot}}$ one, and the latter should be more accurate if used for barometric purposes (*i.e.* depth of main stage of volatile exsolution, see below).

7. Comparison with volcanic gases

The comparison with volcanic gases can be only made for Stromboli, Vulcano and Etna, since hot fumaroles or volcanic plumes have not been emitted at Vesuvius since its last eruption in 1944.

7.1. Stromboli

For Stromboli, direct sampling of fumaroles in the vent is prevented by hazardous access (Allard *et al.*, 1994). The hottest fumarole sampled had a temperature of 410°C which is 700°C lower than the estimated temperature of magma last equilibration in the shallow reservoir. This fumarole has a $\text{H}_2\text{O}/\text{SO}_2$ mole ratio of 11 and a CO_2/SO_2 ratio of 19. Both ratios fall within the range calculated for magmatic fluids in the shallow reservoir (fig. 9) and are thus compatible with an origin from such a depth. The average ratio of CO_2/SO_2 in the dilute plume is 11, which fits also with a shallow reservoir origin, being at the lower end of the calculated fluid composition at 100 bar. The above reasoning, however, implicitly assumes that the volcanic plume composition is dominated by the composition of fluids at depth, and ignores the possible contribution of the melt degassing during ascent from the deep/shallow reservoirs to the surface. In other words, it assumes that the volcanic plume (or hot fumaroles) is fed by bubbles that escape the magma at a certain depth, reaching the surface unmodified (equivalent to a single step degassing at depth under closed system condi-

tions). In the present case, this depth could lie anywhere between 100 and 3000 bar, since in this pressure range magmatic fluids having a CO_2/SO_2 ratio of 10 can be produced. Although such a scenario can apply to a magma intrusion at rest, like during periods of quiescent degassing with no magma emission, this is unlikely to happen in a decompressing magma body, in view of the exceedingly low melt viscosity of hydrous basalts that should allow volatiles degassing, and continuous melt-fluid re-equilibration, even on the short time scales characteristic of Strombolian eruptions.

We have thus calculated the bulk fluid composition arising from melt+fluid contribution, for various melt/fluid ratios, assuming that the melt can fully degass upon eruption or ascent (see also Scaillet and Pichavant, 2003). Figure 17 shows the results in terms of CO_2/SO_2 mole ratios, which is a ratio typically measured in volcanic plumes (Allard *et al.*, 1994). Also shown are the CO_2/SO_2 mole ratios of the melt and fluid phases. Excluding kinetic effects, any fluid vented at Stromboli must lie between




Fig. 17. Effect of the amount of fluid in the reservoir on the final fluid compositions (resulting from the contribution of volatiles from the melt and fluid, melt+fluid), expressed in terms of the $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratio (see text), for Stromboli magmas. The compositions of fluid and melt phases are shown as closed symbols. A fluid amount of 0.1 wt% in the deep reservoir results in a final fluid having a $\text{CO}_2/\text{SO}_{2\text{tot}}$ ratio of 1, whereas an amount of 4 wt% results in a final $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratio of 10, similar to that measured in the volcanic plume of Stromboli (thick grey line, from Allard *et al.*, 1994).

these two poles. As shown on fig. 17, if the deep magma contains 0.1 wt%, the resulting bulk fluid has a CO₂/SO₂ ratio of 1, ie significantly lower than the average one measured in the plume (11). In contrast, if the magma at depth contains 4 wt% fluid, then the resulting CO₂/SO₂ ratio is similar to that measured, with an average at 11. Higher fluid contents at depth would further increase the ratio: a fluid content of 10 wt% would give a CO₂/SO₂ ratio of 22. We thus suggest that in order to produce a volcanic plume with a CO₂/SO₂ ratio of 11, the main magma reservoir must be fluid-saturated with a fluid amount of ca. 4 wt%.

Petrological and geophysical data (see Francalanci *et al.*, 1989) suggest the existence of a large reservoir at 10-14 km depth, in agreement with MI constraints (Métrich *et al.*, 2001; Bertagnini *et al.*, 2003), in which the magmas injected in the shallow reservoir are produced. On this basis, it is probable that such a reservoir is the major source of volatile emissions at Stromboli (Bertagnini *et al.*, 2003). Thus, the above analysis suggests that such a reservoir must contain a significant amount of free fluid in order to reconcile observations (volcanic plume composition, MI volatile contents) and the thermodynamic constraints (this work) on fluid composition. The total volatile content of Strombolian magmas can be estimated using the fluid composition calculated in this work, the amount of volatiles dissolved in the melt (Métrich *et al.*, 2001) and assuming a fluid content of 4 wt%. This gives the following range (corresponding to different volatile contents of MI in pumice): H₂O_{tot}: 2.86-3.45 wt%; CO₂_{tot}: 2.94-3.46 wt%; SO₂_{tot}: 0.29-0.90 wt% (or 1450-4500 ppm sulphur). The H₂O and S values are close to those analysed in MI, which is due to the fact that the concentrations of both volatiles are comparable in melt and fluid. In contrast, the bulk CO₂ content is more than one order of magnitude higher than that dissolved in MI (0.089-0.169 wt%), which is due to the CO₂-rich character (XCO₂>0.6) of the fluid phase at Stromboli. The very high bulk CO₂ content suggests that even the most primitive magmas at Stromboli, *i.e.* those generated at mantle pressures, are fluid-saturated as well.

7.2. Vulcano

The geochemistry of volcanic fumeroles at Vulcano has been extensively discussed and modelled using various approaches and chemical tracers (*e.g.*, Capasso *et al.*, 1994, 2001; Chiodini *et al.*, 1995; Todesco, 1997; Nuccio *et al.*, 1999; Giggenbach *et al.*, 2001; Nuccio and Paonita, 2001; Di Liberto *et al.*, 2002; Diliberto *et al.*, 2002; Paonita *et al.*, 2002). There is a general agreement that the fumerole compositions at Vulcano result from mixing between a magmatic source and a boiling hydrothermal system. Here we focus on the highest temperature fumeroles, in which the magmatic signature is less likely to be affected by such a mixing. The hottest fumeroles reached a temperature of near 700°C, and consisted in 85-90% H₂O, 5-15% CO₂ with minor amounts of sulphur, in the order of 3 % (Capasso *et al.*, 1994), having an *f*O₂ of NNO+1 or very close to that inferred from sulphur speciation (NNO+0.72). Although quite variable, the measured CO₂/SO₂ mole ratio is ca. 10 (*e.g.*, Giggenbach *et al.*, 2001), or similar to that of Stromboli. If taken at face value, the 9/1 ratio of H₂O/CO₂ suggests an origin of such fluids from a reservoir located at 800-900 bar (fig. 12), since lower pressure would correspond to higher mole fraction of CO₂. It must be remembered, however, that this is based on the assumption that MI at Vulcano have a maximum of 50 ppm dissolved CO₂. Clearly, lower dissolved CO₂ would allow water-rich fluids to be produced at lower pressures. However, geochemical modelling (Nuccio *et al.*, 1999) has shown that the hydrothermal contribution is quite high even in the hottest fumeroles (up to 30%), which implies that the H₂O mole fraction of fumeroles are probably maximum values. Similarly, if the CO₂/SO₂ ratio is used to infer the depth of fluid exsolution, then a pressure of 200-300 bar is retrieved (fig. 13), but again, lower CO₂ in MI would decrease this estimate. On the other hand, magmatic gas scrubbing by aquifers is known to strongly deplete magmatic gases of their SO₂ (Symonds *et al.*, 2001) which indicates that the sulphur content of fumeroles are minimum values. This will counteract the effect due to an overestimation of CO₂ in MI, leading to an increase in the CO₂/SO₂ ratio.

As for Stromboli, the above comparison ignores the melt contribution to the volatile budget and that due to decompression or to crystallisation. However, compared to Stromboli, a proper evaluation of this contribution is hampered by the compositional variability of erupted magmas at Vulcano (*e.g.*, Clocchiatti *et al.*, 1994a). That is, the evaluation of melt contribution depends on whether the upper reservoir stores rhyolite or some more mafic composition. If for instance, we assume that the main degassing magmatic source is rhyolite melt, then a fluid amount of 0.5 wt% in the shallow reservoir would be needed in order to release a fluid with appropriate mole ratios of H₂O/CO₂ and CO₂/SO₂. Clearly, however, the uncertainty concerning the chemical nature of the plumbing system, coupled with the lack of precise CO₂ determination in MI, prevent us obtaining robust constraints on bulk volatile compositions and their possible depth of exsolution. To these sources of error must be added the hydrothermal contribution, that modifies to various extents the composition of magmatic fluids (Nuccio *et al.*, 1999). We must therefore await for more precise data on CO₂ content in MI before doing any conclusive statement.

7.3. Etna

Although we lack CO₂ constraints in MI, the case for Etna is made simpler than that of Vulcano, because of the chemical homogeneity of magmas erupted and because the numerous eruptions since 1970 have allowed continuous monitoring of the volcanic plume for CO₂ and SO₂ species, among other volatiles, using modern analytical tools. In addition, contrary to the two previous volcanoes, there is at least one complete data set of hot gas composition available for the 1970 eruption (Huntington, 1973; Gerlach, 1980). The temperature of gas collection (1075°C) is virtually identical to that of magma extrusion which greatly facilitates the comparison with the thermodynamic calculation performed here, since it can be anticipated that the magmatic component in these gases is largely dominant. We first consider this data set and then compare our calculations with remote sensing results. As for Stromboli and Vulcano,

we start by assuming that fluids correspond to deep exsolution of a stagnant magma body and then we evaluate the effect of melt contribution arising during decompression.

The average composition of volcanic gases sampled by Huntington (1973) and restored by Gerlach (1980) (excluding the extremes), reads as (in mole fraction): H₂O=0.48±0.03; CO₂=0.24±0.05, and SO₂=0.26±0.05 (*n*=14). A remarkable feature of this composition is its very high SO₂ content, that equals or even exceeds CO₂, underscoring the potential of Etnean magmas to release sulphur-rich fluids. Inspection of fig. 14a,b shows that production of such a fluid requires both relatively low melt H₂O contents and low pressure conditions, essentially below 1000 bar. Fluids equilibrated with a melt H₂O content of 2.23 wt%, reach a maximum mole fraction of SO₂ below 10%, and are thus unlikely to be the source of gases sampled in 1970. In contrast, at a melt H₂O content of 1 wt%, sulphur-rich gas compositions can be produced at pressures below 1000 bar: the fluid composition calculated at 400 bar (table II) closely approaches the gases measured by Huntington (1973) which in turn suggests that those gases could have been released from a magma body lying at such a depth.

The 1970 data set of gas compositions is unique, and for more recent times information on the chemical composition of volcanic gases has been obtained mostly via remote sensing of volcanic plumes or through analyses of diffuse and peripheral gas emissions (*e.g.*, Allard *et al.*, 1991; Bruno *et al.*, 1999, 2001). However, Allard *et al.* (1991) report a CO₂/SO₂ mole ratio of *ca.* 10 measured in hot (1090°C) crater gases during the 1986 eruption, which is significantly higher than that measured in the 1973 eruption (Huntington, 1973; Gerlach, 1980). The CO₂/SO₂ ratio measured in the summit crater plume ranges from 10 to as high as 38, but Allard *et al.* (1991) consider the lower value as more representative of the average gas output. Such a ratio is attained at around 1000 bar for a melt H₂O content of 2.23 wt%, and at slightly higher pressures for 1 wt% H₂O dissolved in melt, around 1500 bar. The full compositional data corresponding to these gases are not known, in particular the mole fraction of H₂O and SO₂,

which prevents from further constraining the depth of degassing.

We now consider the melt contribution. Figure 18a,b displays the CO_2/SO_2 mole ratio of the bulk fluid released during decompression of a magma containing either 1 or 5 wt% fluid, for a variety of starting pressures, in the interval 400-10000 bar. Also shown are the CO_2/SO_2




Fig. 18a,b. Effect of the amount of fluid in the reservoir on the final fluid compositions (resulting from the contribution of volatiles from the melt and fluid, melt+fluid), expressed in terms of $\text{CO}_2/\text{SO}_{2\text{tot}}$ mole ratio (see text). Calculations were done for melt H_2O contents of 1 (a) and 2.23 (b) wt%, 3300-3400 ppm dissolved sulphur, NNO+0.35, which should correspond to primitive Etnaean magmas. An amount of 1 wt% fluid in the deep reservoir results in a final fluid having a $\text{CO}_2/\text{SO}_{2\text{tot}}$ ratio of 2-3. In contrast, for magmas stored at pressures higher than 2000 bar, an amount of 5 wt% results in a final fluid $\text{CO}_2/\text{SO}_{2\text{tot}}$ ratio of 10 upon near surface degassing, similar to that measured in the volcanic plume at Etna (thick grey line Allard *et al.*, 1991).

mole ratios corresponding to the melt and fluid phases, calculated at the corresponding pressures. The calculations were done for melt H_2O contents of 1 and 2.23 wt% having 3300-3400 ppm sulphur dissolved. Both melt H_2O contents yield similar results, however (fig. 18a,b). Under such conditions, the simulations show that any magma having 1 wt% fluid at any depth falls short in producing a bulk fluid having the observed CO_2/SO_2 ratio of 10, the computed values being in the order of 2-3 at best. In contrast, if the magma has an amount of fluid of 5 wt% and is stored at a depth higher than 3000 bar, then its decompression to near atmospheric conditions should release a fluid having the appropriate CO_2/SO_2 mole ratio. It is interesting to note that beyond 3000 bar, the curves are very steep, illustrating that the calculation is insensitive to the choice of the starting pressure. Also noteworthy is the fact that at pressures below 2000 bar, the CO_2/SO_2 ratio sharply decreases as pressure decreases, such that magmas stored at 400-500 bar with fluids amount of 1-5 wt% would release fluids with CO_2/SO_2 ratios lower than 1, or unlike those measured in the volcanic plume.

The previous calculation assumes obviously that the rising magma exsolves continuously its volatiles but that those volatiles do not escape the parent magma until near atmospheric conditions are attained (*i.e.* a closed system assumption). In other words this corresponds to the case where the differential ascent velocities of a rising magma batch and its exsolved bubbles are negligible. Given the low viscosity of hydrous basalt melt and the significant proportion of fluid inferred (which should correspond to a large volume fraction of bubbles in the magma, see Vergnolle and Jaupart, 1986), this assumption is clearly questionable. Yet, progressive escape of bubbles at high pressure would deplete the magma in CO_2 relative to sulphur, since the CO_2/SO_2 ratio of the fluid is always much higher than 1 at pressures higher than 500 bar (fig. 18a,b). Early depletion of a CO_2 -rich gas phase (*i.e.* open system) would therefore lead to a significant decrease in the final bulk CO_2/SO_2 ratio produced by the degassing lava. We therefore conclude that the fluid amounts derived above can be considered minimum estimates.

If the measured CO_2/SO_2 ratio measured in the volcanic plume or hot fumeroles is indeed representative of magmatic conditions at depth and correspond to full degassing of either out-poured lava or crystallising magma, we can use the above results to infer the bulk volatile content of the magma. To do so we first consider that the main reservoir feeding the Etna eruption lies at 3000 bar, in accordance with the geophysical, geodetic, petrological and geochemical evidence summarised previously. For a reservoir at 3000 bar storing a magma with 1 wt% dissolved H_2O and coexisting with 5 wt% fluid the bulk volatile content is: $\text{H}_2\text{O}_{\text{tot}} = 1.05$ wt%, $\text{CO}_{2\text{tot}} = 4.89$ wt%, $\text{SO}_{2\text{tot}} = 0.81$ wt%. For a magma with 2.23 wt% dissolved H_2O , the figures are: $\text{H}_2\text{O}_{\text{tot}} = 2.70$ wt%, $\text{CO}_{2\text{tot}} = 4.48$ wt%, $\text{SO}_{2\text{tot}} = 0.68$ wt%. A reservoir at 1000 bar with a magma at 1 wt% H_2O in melt would lead to: $\text{H}_2\text{O}_{\text{tot}} = 1.31$ wt%, $\text{CO}_{2\text{tot}} = 3.81$ wt%, $\text{SO}_{2\text{tot}} = 1.57$ wt%, that is a lower pressure increases both H_2O and SO_2 yields relative to CO_2 . The bulk content calculated for H_2O does not depart significantly from that in MI, whereas the bulk contents of CO_2 and SO_2 are considerably higher than the amounts dissolved in the melt (see table II).

In summary, it appears that the gas compositions measured at Etna can be produced in a variety of ways which somehow reflect the complexity and large vertical extent of the plumbing system of the volcano. In particular, the existence of several levels of magma storage in the crust is likely to induce a variety of fluid composition since the pressure of gas release may exert a strong control on its composition (*e.g.*, Caracausi *et al.*, 2003; this work). There are no reasons magmas erupted at Etna over the last decades should have all followed the same ascent path and ponded at constant storage levels. The difference in the CO_2/SO_2 ratios between the 1970 gases and those emitted during the 1975-1995 period may be a reflection of the complex feeding system. The 1970 gas composition can be reproduced by storage and degassing at 400 bar of a relatively water-poor but sulphur-rich magma. In contrast, gases emitted in more recent eruptions require storage at slightly greater depths, around 1000-1500 bar, if they were expelled from this level with no further melt contribution. If melt

accompanied gas escape, then the source was more likely at 3000 bar and filled with relatively undegassed magma (sulphur-rich), coexisting with a significant amount of fluid in order to yield a volcanic plume with an average CO_2/SO_2 ratio of about 10. These are not the only possibilities, however. It could be considered that the magma stored at 3000 bar contained only 1000 ppm dissolved sulphur, which would increase the CO_2/SO_2 ratio (table II). The only way to distinguish between these different scenarios is to have a combined H_2O - S - CO_2 data set on MI, as well as a more detailed knowledge about the possible variation of pre-eruptive melt H_2O contents between eruptive events.

8. Discussion and conclusions

The empirical model presented in this study has allowed us to calculate the sulphur fugacities of hydrous basaltic magmas. There is little control, however, on the values retrieved since even at near surface conditions, $f\text{S}_2$ is a rarely measured parameter in volcanic gases. Pioneering attempts of Sato and Moore (1973) on the 1970 Etna gases failed to measure the $f\text{S}_2$ of the hottest gases emitted and they could only obtain data up to 860°C. Yet, as Gerlach (1980) has shown, the $f\text{S}_2$ measured fits in with that restored from gas composition and thus with our own calculation. Apart from this case, however, we are left to accept the calculated values at face value, until experimental data on the relationships between $f\text{S}_2$ and sulphur solubility in hydrous mafic melts become available.

We have seen that the knowledge of $f\text{S}_2$, $f\text{O}_2$ and $f\text{H}_2\text{O}$ (or $f\text{CO}_2$) enables us to characterise the composition of gas phase at depth which is useful information for the interpretation of volcanic gas data. Here also, however, it needs to be stressed that the calculations are heavily dependent upon the accuracy of the solubility models employed to calculate $f\text{H}_2\text{O}$ and $f\text{CO}_2$. The latter parameter is clearly critical for assessing the pressures of fluid saturation: in this respect, it appears that the CO_2 solubility in K-rich basalts is still not appropriately known. This is a research trend which should be given

the highest priority in the near future for a correct quantitative modeling of volatiles behaviour at Italian volcanoes. Similarly, a full description of degassing processes in shallow magma reservoirs demands that CO₂ contents in MI be measured beyond the concentration levels allowed by the FTIR tool. We suggest that future volatile studies implement analytical tools to such an end.

Accepting that our model retrieves fS_2 within the correct order of magnitude then the calculations show that, except in special circumstances, the addition of sulphur does not strongly change pressure estimates with respect to model calculations which consider only H₂O and CO₂ species. As we have seen, this is due to the generally low values of fS_2 calculated at high pressures. Only at low pressure does there exist the possibility that pressures of fluid saturation could be significantly underestimated by neglecting sulphur-bearing species. The Etna example shows that the misfit between the two approaches can be in the order of several hundreds of bars for magmas rich in sulphur.

Our calculations suggest that fS_2 is usually in the range 0.1-1 bar in the most primitive magmas found at Italian volcanoes. Whether this corresponds to environments unusually rich in sulphur relative to other tectonic settings or other arc magmas, remains an open question given the almost total absence of such information for primitive magmas worldwide. We are therefore currently applying our model to mafic magmas erupted in a variety of tectonic environments in an effort to fill this gap. The calculated fS_2 may display dramatic variations during magmatic evolution, however, that seem to depend on open *versus* closed conduit conditions, as illustrated for Vesuvius. Thus, variations in fS_2 can be potentially used as a sensor to infer the behaviour of magma reservoirs that fed past eruptive events. However, the important variations in fS_2 evidenced in this study conflict with models invoking the control of magma redox state by heterogeneous equilibria between sulphur bearing species dissolved in melt and fluid (*e.g.*, Matthews *et al.*, 1994). In our calculations we assumed that the redox states determined in melt inclusions or whole rocks have remained constant throughout the

magmatic evolution of any particular eruptive event. The lack of two-oxide assemblages in most volcanic products prevents us from testing this fundamental assumption and it could be argued that the large variations we calculate in fS_2 correspond in fact to variations in fO_2 due to volatile exsolution. The following lines of evidence suggest they are not, however. First, if we suppose that equilibria involving sulphur-bearing species control fO_2 then we can use eq. (2.1) to evaluate what magnitude of fO_2 change is needed in order to maintain relatively constant fS_2 during sulphur exsolution. Suppose a mafic melt with 3000 ppm dissolved sulphur at NNO+0.95 and 1100°C with 3 wt% H₂O (similar to Stromboli). If it loses 1000 ppm of sulphur while keeping the same fS_2 , then equation 10 predicts that fO_2 should decrease down to NNO-2.19. While such a reduction during sulphur exsolution is conceivable, it is difficult to reconcile, for instance, with the constant redox state inferred from isotope systematics on pumices at Vesuvius (Marini *et al.*, 1998). Similarly, such a three orders of magnitude drop in fO_2 should be accompanied by strong changes in both phase stabilities and compositions, all of which have gone undetected so far. Second, the detailed thermodynamic analysis of Kilauea volcanic gases has shown that fO_2 is unaffected by shallow volatile exsolution (Gerlach, 1993), the collected gases following a cooling trend parallel to NNO, identical to that inferred for the lavas (see also Symonds *et al.*, 1994). Therefore, although we cannot rule out that fO_2 varies during volatile exsolution, we believe that such a change is relatively minor compared to that encompassed by fS_2 . One possible reason for the relative insensitivity of fO_2 to the behaviour of sulphur is that water, present in abundance in Italian basalts, is the main controlling species of the magma redox state (via the water dissociation reaction).

The approach we followed in modeling magma degassing needs to be implemented by performing numerical simulations of magma ascent and volatiles exsolution, considering open and closed system behaviours. However, the comparison between our calculations and observed gas compositions suggests that a fluid phase is present at depth, at amounts in the

order of 4-5 wt%. Although the available data on MI do not allow to prove yet such an hypothesis, it is important to note that if it is true then it will require a significant reevaluation of magma budgets based on volatile emissions (e.g., Allard, 1997; Bruno *et al.*, 2001). Current evaluations have only considered the sulphur or carbon dioxide dissolved in melt as the ultimate source of volcanic degassing. However, consideration of a fluid phase could lead to a significant downsizing of magma volumes required to balance the volatile budget measured at permanently degassing volcanoes such as Etna and Stromboli. To illustrate this point, we take the case of an Etna magma reservoir at 1000 bar with 5 wt% fluid previously discussed, whose SO₂ content is 1.57 wt%. Allard (1997) calculated that the average SO₂ output at Etna for the period 1975-1995 is 1.7·10⁶ tons/year, whereas the average amount of erupted magmas over the same period is 81·10⁶ tons/year. By considering only the melt contribution, Allard (1997) calculated that an additional amount of 678·10⁶ tons/year of magma must be intruded (and not erupted) in order to account for sulphur emissions. In contrast, if a fluid phase is present at 1000 bar, then the 81·10⁶ tons of magma extruded every year could release 1.27·10⁶ tons of sulphur or 75 % of the total amount of sulphur emitted, and the need for endogenous magma degassing at Etna would be much less severe than previously thought.

Acknowledgements

The paper greatly benefited from constructive reviews by H. Shinoara and R. Moretti.

REFERENCES

- ALLARD, P. (1997): Endogeneous magma degassing and storage at Mt. Etna, *Geophys. Res. Lett.*, **24**, 2219-2222.
- ALLARD, P., J. CARBONNELLE, D. DAJLEVIC, J. LE BRONEC, P. MOREL, M.C. ROBE, J.M. MAURENAS, R. FAIVRE-PIERRET, D. MARTIN, J.C. SABROUX and P. ZETTWOOG (1991) : Eruptive and diffuse emissions of CO₂ from Mt. Etna, *Nature*, **351**, 387-390.
- ALLARD, P., J. CARBONNELLE, N. MÉTRICH, H. LOYER and P. ZETTWOOG (1994): Sulphur output and magma degassing budget of Stromboli volcano, *Nature*, **368**, 326-329.
- ANDERSON, A.T., S. NEWMAN, S.N. WILLIAMS, T.H. DRUITT, C. SKIRIUS and E. STOLPER (1989): H₂O, CO₂, Cl gas in Plinian and ash-flow Bishop rhyolite, *Geology*, **17**, 221-225.
- ARMIENTI, P., F. INNOCENTI, R. PETRINI, M. POMPILIO and L. VILLARI (1988): Sub-aphyric alkali basalt from Mt. Etna: inferences on the depth and composition of the source magma, *Rend. Soc. Ital. Mineral. Petrol.*, **43**, 877-891.
- BALLHAUS, C., R.F. BERRY and D.H. GREEN (1991): High pressure experimental calibration of the olivine-orthopyroxene-spinel oxygen barometer: implications for the oxidation state of the upper mantle, *Contrib. Mineral. Petrol.*, **107**, 27-40.
- BELKIN, H.E., B. DE VIVO, K. TÖRÖK and J.D. WEBSTER (1998): Pre-eruptive volatile content, melt inclusion chemistry, and microthermometry of interplinian Vesuvius lavas (pre-A.D. 1631), *J. Volcanol. Geotherm. Res.*, **82**, 79-95.
- BERTAGNINI, A., N. MÉTRICH, P. LANDI and M. ROSI (2003): Stromboli volcano (Aeolian Archipelago, Italy): an open window on the deep-feeding system of a steady state basaltic volcano, *J. Geophys. Res.*, **108** (B7), 2336, doi:10.1029/2002JB002146.
- BONACCORSO, A. (1996): Dynamic inversion of ground deformation data for modelling volcanic sources (Etna 1991-93), *Geophys. Res. Lett.*, **23**, 451-454.
- BONACCORSO, A. (2001): Mt. Etna volcano: modelling of ground deformation patterns of recent eruptions and considerations on the associated precursors, *J. Volcanol. Geotherm. Res.*, **109**, 99-108.
- BRUNO, N., T. CALTABIANO and R. ROMANO (1999): SO₂ emissions at Mt. Etna with particular reference to the period 1993-1995, *Bull. Volcanol.*, **60**, 405-411.
- BRUNO, N., T. CALTABIANO, S. GIAMMANCO and R. ROMANO (2001): Degassing of SO₂ and CO₂ at Mt. Etna (Sicily) as an indicator of pre-eruptive ascent and shallow emplacement of magma, *J. Volcanol. Geotherm. Res.*, **110**, 137-153.
- BUCHANAN, D.L. and J. NOLAN (1979): Solubility of sulfur and sulfide immiscibility in synthetic tholeiitic melts and their relevance to Bushveld-complex rocks, *Can. Mineral.*, **17**, 483-494.
- CAPASSO, G., S. INGUAGGIATO and P.M. NUCCIO (1994): Chemical variations of the fumarolic gases of La Fossa Vulcano crater, *Acta Vulcanol.*, **3**, 173-187.
- CAPASSO, G., W. D'ALESSANDRO, R. FAVARA, S. INGUAGGIATO and F. PARELLO (2001): Interaction between the deep fluids and the shallow groundwaters on Vulcano Island (Italy), *J. Volcanol. Geotherm. Res.*, **108**, 187-198.
- CARACAUSI, A., F. ITALIANO, A. PAONITA, A. RIZZO and P. M. NUCCIO (2003): Evidence of deep magma degassing and ascent by geochemistry of peripheral gas emissions at Mt. Etna (Italy): assessment of the magmatic reservoir pressure, *J. Geophys. Res.*, **108** (O), XXXX, doi:10.1029/2002JB002095.
- CARROLL, M.C. and M.J. RUTHERFORD (1987): The stability of igneous anhydrite: experimental results and implications for sulfur behavior in the 1982 El Chichon trachyandesite and other evolved magmas, *J. Petrol.*, **28**, 781-801.

- CARROLL, M.C. and J.D. WEBSTER (1994): Solubilities of sulfur, noble gases, nitrogen, chlorine and fluorine in magmas, in *Volatiles in Magmas*, edited by M.R. CARROLL and J.R. HOLLOWAY, *Rev. Mineral.*, **30**, 231-279.
- CHIODINI, G., R. CIONI, L. MARINI and C. PANICHI (1995): Origin of the fumarolic fluids of Vulcano Island, Italy and implications for volcanic surveillance, *Bull. Volcanol.*, **57**, 99-110.
- CIONI, R. (2000): Volatile content and degassing processes in the A.D. 79 magma chamber at Vesuvius (Italy), *Contrib. Mineral. Petrol.*, **140**, 40-54.
- CIONI, R., L. CIVETTA, P. MARIANELLI, N. MÉTRICH, R. SANTACROCE and A. SBRANA (1995): Compositional layering and syn-eruptive mixing of a periodically refilled shallow magma chamber: the A.D. 79 Plinian eruption of Vesuvius, *J. Petrol.*, **36**, 739-776.
- CIONI, R., P. MARIANELLI and R. SANTACROCE (1998): Thermal and compositional evolution of the shallow magma chambers of Vesuvius: evidence from pyroxene phenocrysts and melt inclusions, *J. Geophys. Res.*, **103**, 18277-18294.
- CIVETTA, L. and R. SANTACROCE (1992): Steady state magma supply in the last 3400 years of Vesuvius activity, *Acta Vulcanol.*, **2**, 147-159.
- CIVETTA, L., R. GALATI and R. SANTACROCE (1991): Magma mixing and convective compositional layering within the Vesuvius magma chamber, *Bull. Volcanol.*, **53**, 287-300.
- CLEMENTE, B., B. SCAILLET and M. PICHAVANT (2004): The solubility of sulphur in hydrous rhyolitic melts, *J. Petrol.*, **45**, 2171-2196.
- CLOCCHIATTI, R.A. and N. MÉTRICH (1984): Témoignages de la contamination dans les produits des éruptions explosives des M. Silvestri (1892) et M. Rossi (1669) (Mt. Etna), *Bull. Volcanol.*, **47**, 909-928.
- CLOCCHIATTI, R.A., J. WEISZ, M. MOSBAH and J.C. TANGUY (1992): Coexistence de verres alcalins et tholéïtiques saturés en CO₂ dans les olivines des hyaloclastites d'Acì Castello (Etna, Sicile, Italie). Arguments en faveur d'un manteau anormal et d'un réservoir profond, *Acta Vulcanol.*, **2**, 161-173.
- CLOCCHIATTI, R., A. GIONCADA, M. MOSBAH and A. SBRANA (1994a): Possible deep origin of sulfur output at Vulcano (Southern Italy) in the light of melt inclusion studies, *Acta Vulcanol.*, **5**, 49-53.
- CLOCCHIATTI, R., A. DEL MORO, A. GIONCADA, J.L. JORON, M. MOSBAH, L. PINARELLI and A. SBRANA (1994b): Assessment of a shallow magmatic system: the 1888-1890 eruption, Vulcano Island, Italy, *Bull. Volcanol.*, **56**, 466-486.
- DI CARLO, I., M. PICHAVANT, S. ROTOLO and B. SCAILLET (2004): Ascent of Stromboli yellow pumice magmas: experimental simulation at $P \leq 4$ kb, *Geophys. Res. Abstr.*, **6**, 04122.
- DI LIBERTO, V., P.M. NUCCIO and A. PAONITA (2002): Genesis of chlorine and sulphur in fumarole emissions at Vulcano Island (Italy): assessment of pH and redox conditions in the hydrothermal system, *J. Volcanol. Geotherm. Res.*, **116**, 137-150.
- DILIBERTO, I.S., S. GURRIERI and M. VALENZA (2002): Relationships between diffuse CO₂ emissions and volcanic activity on the Island of Vulcano (Aeolian Islands, Italy) during the period 1984-1994, *Bull. Volcanol.*, **64**, 219-228.
- DIXON, J.E. (1997): Degassing of alkalic basalts, *Am. Mineral.*, **82**, 368-378.
- DIXON, J.E., E.M. STOLPER and J.R. HOLLOWAY (1995): An experimental study of water and carbon dioxide solubilities in mid-ocean ridge basaltic liquids, Part I. Calibration and solubility models, *J. Petrol.*, **36**, 1607-1631.
- FERRY, J.M. and L. BAUMGARTNER (1987): Thermodynamic models of molecular fluids at the elevated pressures and temperatures of crustal metamorphism, in *Thermodynamic Modelling of Geological Materials: Minerals, Fluids and Melts*, edited by I.S.E. CARMICHAEL and H.P. EUGSTER, *Rev. Mineral.*, **17**, 323-365.
- FLOWERS, G.C. (1979): Correction of Holloway's (1977) adaptation of the modified Redlich-Kwong equation of state for calculation of the fugacities of molecular species in supercritical fluids of geologic interest, *Contrib. Mineral. Petrol.*, **69**, 315-18.
- FRANCALANCI, L., P. MANETTI and A. PECCERILLO (1989): Volcanological and magmatological evolution of Stromboli volcano (Aeolian Islands): the roles of fractional crystallization, magma mixing, crustal contamination and source heterogeneity, *Bull. Volcanol.*, **51**, 355-378.
- FREDA, C., M. GAETA, D.M. PALLADINO and R. TRIGILA (1997): The Villa Senni eruption (Alban Hills, Central Italy): the role of H₂O and CO₂ on the magma chamber evolution and on the eruptive scenario, *J. Volcanol. Geotherm. Res.*, **78**, 103-120.
- FREZZOTTI, M.L., B. DE VIVO and R. CLOCCHIATTI (1991): Melt-mineral-fluid interactions in ultramafic nodules from alkaline lavas of Mt. Etna (Sicily, Italy): melt and fluid inclusion evidence, *J. Volcanol. Geotherm. Res.*, **47**, 209-219.
- FROESE, E. and A.E. GUNTER (1976): A note on the pyrrhotite-sulfur vapor equilibrium, *Econ. Geol.*, **71**, 1589-1594.
- GERLACH, T.M. (1980): Evaluation and restoration of the 1970 volcanic gas analyses from Mt. Etna, Sicily, *J. Volcanol. Geotherm. Res.*, **6**, 165-178.
- GERLACH, T.M. (1993): Oxygen buffering of Kilauea volcanic gases and the oxygen fugacity of Kilauea basalt, *Geochim. Cosmochim. Acta*, **57**, 795-814.
- GIGGENBACH, W.F., D. TEDESCO, Y. SULISTIYO, A. CAPRAI, R. CIONI, R. FAVARA, T.P. FISCHER, J.I. HIRABAYASHI, M. KORZHINSKY, M. MARTINI, I. MENYAILOV and H. SHINOARA (2001): Evaluation of results from the fourth and fifth IAVCEI field workshops on volcanic gases, Vulcano Island, Italy and Java, Indonesia, *J. Volcanol. Geotherm. Res.*, **108**, 157-172.
- GIONCADA, A., R. CLOCCHIATTI, A. SBRANA, P. BOTTAZZI, D. MASSARE and L. OTTOLINI (1998): A study of melt inclusions at Vulcano (Aeolian Islands, Italy): insights on the primitive magmas and on the volcanic feeding system, *Bull. Volcanol.*, **60**, 286-306.
- HARRIS, A.J.L. and D.S. STEVENSON (1997): Magma budgets and steady-state activity of Vulcano and Stromboli, *Geophys. Res. Lett.*, **24**, 1043-1046.
- HAUGHTON, D.R., P.L. ROEDER and B.J. SKINNER (1974): Solubility of sulfur in mafic magmas, *Econom. Geol.*, **69**, 451-467.
- HOLLOWAY, J.R. (1977): Fugacity and activity of molecular species in supercritical fluids, in *Thermodynamics in Geology*, edited by D. FRASE (Dordrecht: Reidel Publishing Company), 161-81.

- HOLLOWAY, J.R. and J. BLANK (1994): Applications of experimental results to C-O-H species in natural melts, in *Volatiles in Magmas*, edited by M.R. CARROLL and J.R. HOLLOWAY, *Rev. Mineral.*, **30**, 187-230.
- HUNTINGDON, A.T. (1973): The collection of volcanic gases from Mt. Etna, *Phil. Trans. R. Soc. London*, **274**, 119-128.
- KAMENETSKY, V. and R. CLOCCHIATTI (1996): Primitive magmatism of Mt. Etna: insights from mineralogy and melt inclusions, *Earth Planet. Sci. Lett.*, **142**, 553-572.
- KATSURA, T. and S. NAGASHIMA (1974): Solubility of sulfur in some magmas at 1 atmosphere, *Geochim. Cosmochim. Acta*, **38**, 517-531.
- KILINC, A., I.S.E. CARMICHAEL, M.L. RIVERS and R.O. SACKS (1984): The ferric-ferrous ratio of natural silicate liquids equilibrated in air, *Contrib. Mineral. Petrol.*, **83**, 136-140.
- LIMA, A., H.764E. BELKIN and K. TÖRÖK (1999): Understanding Vesuvius magmatic processes: evidence from primitive silicate-melt inclusions in medieval scoria clino-pyroxenes (Terzigno Formation), *Mineral. Petrol.*, **65**, 185-206.
- LUHR, J.F. (1990): Experimental phase relations of water and sulfur-saturated arc magmas and the 1982 eruptions of El Chichón volcano, *J. Petrol.*, **31**, 1071-1114.
- MARIANELLI, P., N. MÉTRICH, R. SANTACROCE and A. SBRANA (1995): Mafic magma batches at Vesuvius: a glass inclusion approach to the modalities of feeding stratovolcanoes, *Contrib. Mineral. Petrol.*, **120**, 159-169.
- MARIANELLI, P., N. MÉTRICH and A. SBRANA (1999): Shallow and deep reservoirs involved in magma supply of the 1944 eruption of Vesuvius, *Bull. Volcanol.*, **61**, 48-63.
- MARINI, L., V. CHIAPPINI, R. CIONI, G. CORTECCI, E. DINELLI, C. PRINCIPE and G. FERRARA (1998): Effect of degassing on sulfur contents of $\delta^{34}\text{S}$ values in Somma-Vesuvius magmas, *Bull. Volcanol.*, **60**, 187-194.
- MATTHEWS, S.J., A.P. JONES and A.D. BEARD (1994): Buffering of melt oxygen fugacity by sulphur redox reactions in calc-alkaline magmas, *J. Geol. Soc. London*, **151**, 815-23.
- MAVROGENES, J. and H.S.C. O'NEILL (1999): The relative effects of pressure, temperature and oxygen fugacity on the solubility of sulfide in mafic magmas, *Geochim. Cosmochim. Acta*, **63**, 1173-1180.
- MÉTRICH, N. (1990): Chlorine and fluorine in tholeiitic and alkaline lavas of Etna (Sicily), *J. Volcanol. Geotherm. Res.*, **40**, 133-148.
- MÉTRICH, N. and R. CLOCCHIATTI (1989): Melt inclusion investigation of the volatile behaviour in historic alkali basaltic magmas of Etna, *Bull. Volcanol.*, **51**, 185-198.
- MÉTRICH, N. and R. CLOCCHIATTI (1996): Sulfur abundance and its speciation in oxidized alkaline melts, *Geochim. Cosmochim. Acta*, **60**, 4151-4160.
- MÉTRICH, N. and M. MOSBAH (1988): Détermination des teneurs en carbone de quelques verres basaltiques; analyses par réactions nucléaires, *Bull. Mineral.*, **111**, 511-522.
- MÉTRICH, N. and M.J. RUTHERFORD (1998): Low pressure crystallization paths of H₂O-saturated basaltic-hawaiitic melts from Mt. Etna: implications for open-system degassing of basaltic volcanoes, *Geochim. Cosmochim. Acta*, **62**, 1195-1205.
- MÉTRICH, N., R. CLOCCHIATTI, M. MOSBAH and M. CHAUDSIDON (1993): The 1989-1990 activity of Etna magma mingling and ascent of H₂O-Cl-S-rich basaltic magma. Evidence from melt inclusions, *J. Volcanol. Geotherm. Res.*, **59**, 131-144.
- MÉTRICH, N., A. BERTAGNINI, P. LANDI and M. ROSI (2001): Crystallisation driven by decompression and water loss at Stromboli volcano (Aeolian Islands, Italy), *J. Petrol.*, **42**, 1471-1490.
- MÉTRICH, N., M. BONNIN-MOSBAH, J. SUSINI, B. MENEZ and L. GALOISY (2002): Presence of sulfite (S^{IV}) in arc magma: implications for volcanic sulfur emissions, *Geophys. Res. Lett.*, **29**, 10.1029/2001GL014607.
- MORETTI, R., P. PAPALE and G. OTTONELLO (2003): A model for the saturation of C-O-H-S fluids in silicate melts, in *Volcanic Degassing*, edited by C. OPPENHEIMER, D.M. PYLE and J. BARCLAY, *Geol. Soc. Spec. Publ.* **213**, 81-101.
- MURRU, M., C. MONTUORI, M. WYSS and E. PRIVITERA (1999): The location of magma chambers at Mt. Etna, Italy, mapped by *b*-values, *Geophys. Res. Lett.*, **26**, 2553-2556.
- NUCCIO, P.M. and A. PAONITA (2001): Magmatic degassing of multicomponent vapors and assessment of magma depth: application to Vulcano Island (Italy), *Earth Planet. Sci. Lett.*, **193**, 467-481.
- NUCCIO, P.M., A. PAONITA and F. SORTINO (1999): Geochemical modelling of mixing between magmatic and hydrothermal gases: the case of Vulcano Island, Italy, *Earth Planet. Sci. Lett.*, **167**, 321-333.
- O'NEILL, H.St. and J.A. MAVROGENES (2002): The sulfide capacity and the sulfur content at sulfide saturation of silicate melts at 1400°C and 1 bar, *J. Petrol.*, **43**, 1049-1087.
- PAONITA, A., R. FAVARA, P.M. NUCCIO and F. SORTINO (2002): Genesis of fumarolic emissions as inferred by isotope mass balances: CO₂ and water at Vulcano Island, Italy, *Geochim. Cosmochim. Acta*, **66**, 759-772.
- PAPALE, P. (1997): Thermodynamic modelling of the solubility of a one-component H₂O or CO₂ fluid in silicate liquids, *Contrib. Mineral. Petrol.*, **126**, 237-251.
- RAIA, F., J.D. WEBSTER and B. DE VIVO (2000): Pre-eruptive volatile contents of Vesuvius magmas: constraints on eruptive history and behavior, I. The medieval and modern interplinian activities, *Eur. J. Mineral.*, **12**, 179-193.
- ROGGENSACK, K., R.L. HERVIG, S.B. MCKNIGHT and S.N. WILLIAMS (1997): Explosive basaltic volcanism from Cerro Negro volcano: influence of volatiles on eruptive style, *Science*, **277**, 1639-1642.
- SATO, M. and J.G. MOORE (1973): Oxygen and sulfur fugacities of magmatic gases directly measured in active vents of Mt. Etna, *Philos. Trans. R. Soc. London, Ser. A*, **274**, 137-146.
- SAXENA, S.K. and Y. FEI (1987): High pressure and high temperature fluid fugacities, *Geochim. Cosmochim. Acta*, **51**, 783-791.
- SCAILLET, B. and B.W. EVANS (1999): The June 15, 1991 eruption of Mt. Pinatubo. I. Phase equilibria and pre-eruption *P-T-fO₂-fH₂O* conditions of the dacite magma, *J. Petrol.*, **40**, 381-411.
- SCAILLET, B. and M. PICHAVANT (2003): Experimental constraints on volatile abundances in arc magmas and their implications for degassing processes, in *Volcanic Degassing*, edited by C. OPPENHEIMER, D.M. PYLE and J. BARCLAY, *Geol. Soc. Spec. Publ.* **213**, 23-52.
- SCAILLET, B., CLEMENTE, B.W. EVANS and M. PICHAVANT

- (1998): Redox control of sulfur degassing in silicic magmas, *J. Geophys. Res.*, **103**, 23937-23949.
- SHI, P. and S.K. SAXENA (1992): Thermodynamic modeling of the C-O-H-S fluid system, *Am. Mineral.*, **77**, 1038-1049.
- SIGNORELLI, S. and B. CAPACCIONI (1999): Behaviour of chlorine prior and during the 79 A.D. Plinian eruption of Vesuvius (Southern Italy) as inferred from the present distribution in glassy mesostases and whole-pumices, *Lithos*, **46**, 715-730.
- SIGNORELLI, S., G. VAGELLI and C. ROMANO (1999): Pre-eruptive volatile (H₂O, F, Cl and S) contents of phonolitic magmas feeding the 3550-year old Avellino eruption from Vesuvius, Southern Italy, *J. Volcanol. Geotherm. Res.*, **93**, 237-256.
- SOBOLEV, A.V., V.S. KAMENETSKY, N. MÉTRICH, R. CLOCCHIATTI, N.N. KONONKOVA, A.L. DEVIRTS and V.I. USTINOV (1991): Volatile regime and crystallisation conditions in Etna Hawaiiite lavas, *Geochimica*, **9**, 1277-1290.
- SYMONDS, R.B., W.I. ROSE, G.J.S. BLUTH and T.M. GERLACH (1994): Volcanic gas studies: methods, results, and applications, in *Volatiles in Magmas*, edited by M.R. CARROLL and J.R. HOLLOWAY, *Rev. Mineral.*, **30**, 1-60.
- SYMONDS, R.B., T.M. GERLACH and M.H. REED (2001): Magmatic gas scrubbing: implications for volcano monitoring, *J. Volcanol. Geotherm. Res.*, **108**, 303-341.
- TANGUY, J.C., M. CONDOMINES and G. KIEFFER (1997): Evolution of the Mt. Etna magma: constraints on the present feeding system and eruptive mechanism, *J. Volcanol. Geotherm. Res.*, **75**, 221-250.
- TODESCO, M. (1997): Origin of fumarolic fluids at Vulcano (Italy). Insights from isotope data and numerical modeling of hydrothermal circulation, *J. Volcanol. Geotherm. Res.*, **79**, 63-85.
- TRIGILA, R. and A.A. DE BENEDETTI (1993): Petrogenesis of Vesuvius historical lavas constrained by Pearce element ratios analysis and experimental phase equilibria, *J. Volcanol. Geotherm. Res.*, **58**, 315-343.
- TRIGILA, R., F.J. SPERA and C. AURISICCHIO (1990): The 1983 Mt. Etna eruption: thermochemical and dynamical inferences, *Contrib. Mineral. Petrol.*, **104**, 594-608.
- VAGELLI, G., H.E. BELKIN, B. DE VIVO and R. TRIGILA (1992): Silicate-melt inclusions in recent Vesuvius lavas (A.D. 1631-1944), I. Petrography and microthermometry, *Eur. J. Mineral.*, **4**, 1113-1124.
- VAGELLI, G., B. DE VIVO and R. TRIGILA (1993): Silicate-melt inclusions in recent Vesuvius lavas (A.D. 1631-1944), II. Analytical chemistry, *J. Volcanol. Geotherm. Res.*, **58**, 367-376.
- VERGNIOLE, S. and C. JAUPART (1986): Separated two-phase flow and basaltic eruptions, *J. Geophys. Res.*, **91**, 12842-12860.
- WALLACE, P.J. and I.S.E. CARMICHAEL (1992): Sulfur in basaltic magmas, *Geochim. Cosmochim. Acta*, **56**, 1683-1874.
- WEBSTER, J.D. and B. DE VIVO (2002): Experimental and modeled solubilities of chlorine in aluminosilicate melts, consequences of magma evolution, and implications for exsolution of hydrous chloride melt at Mt. Somma-Vesuvius, *Am. Mineral.*, **87**, 1046-1061.
- WEBSTER, J.D., F. RAIA, B. DE VIVO and G. ROLANDI (2001): The behavior of chlorine and sulfur during differentiation of the Mt. Somma-Vesuvius magmatic system, *Mineral. Petrol.*, **73**, 177-200.
- ZHANG, Y. (1999): H₂O in rhyolitic glasses and melts: measurement, speciation, solubility, and diffusion, *Rev. Geophys.*, **37**, 493-516.