

# Cardinal invariants and special maps of quasicontinuous functions with the topology of pointwise convergence

MANDEEP KUMAR <sup>a</sup>  AND BRIJ KISHORE TYAGI <sup>b</sup> 

<sup>a</sup> Department of Mathematics, University of Delhi, Delhi - 110007, India. ([mjakhar5@gmail.com](mailto:mjakhar5@gmail.com))

<sup>b</sup> Department of Mathematics, Atma Ram Sanatan Dharma College, University of Delhi, New Delhi - 110021, India. ([brijkishore.tyagi@gmail.com](mailto:brijkishore.tyagi@gmail.com))

Communicated by D. N. Georgiou

## ABSTRACT

---

For topological spaces  $X$  and  $Y$ , let  $Q_p(X, Y)$  be the space of all quasicontinuous functions from  $X$  to  $Y$  with the topology of pointwise convergence. In this paper, we study the cardinal invariants such as character, weight, density, pseudocharacter, spread and cellularity of the space  $Q_p(X, Y)$ . We also discuss the properties of the restriction and induced maps related to the space  $Q_p(X, Y)$ .

---

2020 MSC: 54C35; 54C08; 54C30.

KEYWORDS: quasicontinuous functions; topology of pointwise convergence; character; weight; density; spread; cellularity; induced map; restriction map.

## 1. INTRODUCTION

Kempisty [10] introduced a weaker form of continuity for real-valued functions, named as quasicontinuity. The properties of quasicontinuous functions are discussed in many papers, for example see [2, 13, 15, 16].

The quasicontinuous functions have various applications in different areas of mathematics; for instance topological groups [11], dynamical systems [4] and the study of minimal usco and minimal cusco maps [6]. Some examples [3] of

quasicontinuous functions are the doubling function

$$D : [0, 1) \rightarrow [0, 1) \text{ defined by } D(x) = 2x \pmod{1},$$

the extended  $\sin(1/x)$  function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} \sin(\frac{1}{x}) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0, \end{cases}$$

the floor function from  $\mathbb{R}$  to  $\mathbb{R}$  defined by

$$\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\},$$

and any monotonic left or right continuous function from  $\mathbb{R}$  to  $\mathbb{R}$  [15].

The set of all real-valued quasicontinuous maps on a topological space  $X$  with the topology of pointwise convergence, denoted by  $Q_p(X, \mathbb{R})$ , is studied in [7, 8, 9]. The pointwise convergence of real-valued quasicontinuous maps defined on a Baire space is examined in [7]. In [9] metrizable, first countability, closed and compact subsets of the space  $Q_p(X, \mathbb{R})$  are discussed. The cardinal functions of the space  $Q_p(X, \mathbb{R})$  are studied in [8].

In this paper, we study the results about metrizable, first countability and cardinal functions of the space  $Q_p(X, Y)$  along with the concept of induced and the restriction maps.

In a more detail, this paper is organized as follows: In Section 3, we define the topology of pointwise convergence on  $Q(X, Y)$ , the set of all quasicontinuous functions from a topological space  $X$  to a topological space  $Y$ . In Section 4, when  $X$  is Hausdorff and  $Y$  is a nontrivial  $T_1$ -space, we compare the cardinal functions  $\pi$ -character, character and weight of the space  $Q_p(X, Y)$ . Moreover, if  $Y$  is second countable, we characterize these cardinal functions. For a regular space  $X$  and a nontrivial  $T_1$ -space  $Y$ , we discuss pseudocharacter and spread of the space  $Q_p(X, Y)$ . We also show that  $Q_p(X, Y)$  is dense in the space  $Y^X$ . In Section 5, we discuss the topological properties of induced maps and the restriction map related to the space  $Q_p(X, Y)$ .

## 2. PRELIMINARIES

Throughout this paper, the symbols  $X, Y, Z$  are topological spaces unless otherwise stated,  $\mathbb{R}$  is the space of real numbers with the usual topology,  $\mathbb{N}$  is the set of positive integers, and  $\mathbb{I}$  is the closed interval  $[-1, 1]$ . The topology of a space  $X$  is denoted by  $\tau(X)$ . By a nontrivial space we mean a topological space with at least two different points. The symbol  $A^\circ$  denotes the interior of  $A$  in  $X$  and the symbol  $\bar{A}$  denotes the closure of  $A$  in  $X$ .

**Definition 2.1.** A map  $f : X \rightarrow Y$  is quasicontinuous [15] at  $x \in X$  if for every open set  $U$  containing  $x$  and every open set  $V$  containing  $f(x)$ , there exists a nonempty open set  $G \subseteq U$  such that  $f(G) \subseteq V$ . If  $f$  is quasicontinuous at every point of  $X$ , we say that  $f$  is quasicontinuous.

Note that every continuous map is quasicontinuous. Conversely, for  $X = [0, 1)$  with the usual topology and  $Y = [0, 1)$  with the Sorgenfrey topology, the identity map from  $X$  to  $Y$  is quasicontinuous but nowhere continuous [12].

Levine [13] studied quasicontinuous maps under the name of semi-continuity using the terminology of semi-open sets. A subset  $A$  of a space  $X$  is said to be semi-open (or quasi-open [15]) if  $A \subseteq \overline{A^\circ}$ . A map  $f : X \rightarrow Y$  is quasicontinuous if and only if for every open set  $V$  in  $Y$ ,  $f^{-1}(V)$  is semi-open in  $X$ .

### 3. QUASICONTINUOUS FUNCTIONS AND THE TOPOLOGY OF POINTWISE CONVERGENCE

Let  $F(X, Y)$  be the set of all functions and  $C(X, Y)$  be the set of all continuous functions from  $X$  to  $Y$ . The function spaces  $F(X, Y)$  and  $C(X, Y)$  with the topology of pointwise convergence denoted by  $F_p(X, Y)$  and  $C_p(X, Y)$ , respectively, are widely studied in the literature, for example [1, 5, 14, 17].

For  $x \in X$  and  $V \in \tau(Y)$ , let  $S(x, V) = \{f \in F(X, Y) : f(x) \in V\}$ . Then  $F_p(X, Y)$  has a subbase

$$\mathcal{S} = \{S(x, V) : x \in X, V \in \tau(Y)\}.$$

Note that  $F(X, Y) = Y^X$  and the topology of pointwise convergence on  $F(X, Y)$  is just the product topology on  $Y^X$ .

Let  $Q(X, Y)$  be the set of all quasicontinuous functions in  $F(X, Y)$ . The space  $Q(X, Y)$  with the topology of pointwise convergence is the subspace  $Q(X, Y)$  of the space  $F_p(X, Y)$  and is denoted by  $Q_p(X, Y)$ .

For  $x \in X$  and  $V \in \tau(Y)$ , denote  $[x, V] = \{f \in Q(X, Y) : f(x) \in V\}$ . Then  $\mathcal{S}' = \{[x, V] : x \in X, V \in \tau(Y)\}$  is a subbase for the space  $Q_p(X, Y)$ . Observe that for  $x \in X$  and  $V_1, V_2 \in \tau(Y)$ , we have  $[x, V_1] \cap [x, V_2] = [x, V_1 \cap V_2]$ . For  $x_1, \dots, x_n \in X$  and  $V_1, \dots, V_n \in \tau(Y)$ , denote

$$[x_1, \dots, x_n; V_1, \dots, V_n] = \{f \in Q(X, Y) : f(x_i) \in V_i, 1 \leq i \leq n\}.$$

Clearly the family

$$\mathcal{B} = \{[x_1, \dots, x_n; V_1, \dots, V_n] : x_i \in X, V_i \in \tau(Y), 1 \leq i \leq n, n \in \mathbb{N}\}$$

is a base for the space  $Q_p(X, Y)$ . If  $\mathcal{V}$  is a basis for  $Y$  then the family

$$\mathcal{B}' = \{[x_1, \dots, x_n; V_1, \dots, V_n] : x_i \in X, V_i \in \mathcal{V}, 1 \leq i \leq n, n \in \mathbb{N}\}$$

is also a basis for the space  $Q_p(X, Y)$ . If  $(Y, d)$  is a metric space then for  $f \in Q(X, Y); x_1, \dots, x_n \in X$  and  $\epsilon > 0$ , denote

$$O(f, x_1, \dots, x_n, \epsilon) = \{g \in Q(X, Y) : d(g(x_i), f(x_i)) < \epsilon, 1 \leq i \leq n\}.$$

It is easy to see that the family

$$\mathcal{B}_f = \{O(f, x_1, \dots, x_n, \epsilon) : x_1, \dots, x_n \in X, n \in \mathbb{N}, \epsilon > 0\}$$

is a local base at  $f \in Q_p(X, Y)$ .

### 4. CARDINAL FUNCTIONS AND THE SPACE $Q_p(X, Y)$

In this section, we discuss first countability, metrizability and cardinal functions of the space  $Q_p(X, Y)$ . Before generalizing some results obtained in [8, 9], first we recall definitions of the cardinal functions for a topological space [8, 14].

A collection  $\mathcal{V}$  of nonempty open subsets of  $X$  is called a local  $\pi$ -base at  $x \in X$  if for each open set  $U$  containing  $x$ , there exists  $V \in \mathcal{V}$  such that  $V \subseteq U$ . The  $\pi$ -character of a point  $x \in X$  is  $\pi_\chi(x, X) = \aleph_0 + \min\{|\mathcal{V}| : \mathcal{V} \text{ is a local } \pi\text{-base at } x\}$ . The  $\pi$ -character of a space  $X$  is defined as  $\pi_\chi(X) = \sup\{\pi_\chi(x, X) :$

$x \in X$ . The character of a space  $X$  is  $\chi(X) = \sup\{\chi(x, X) : x \in X\}$ , where  $\chi(x, X) = \aleph_0 + \min\{|\mathcal{B}_x| : \mathcal{B}_x \text{ is a base at } x\}$ . The weight of a space  $X$  is defined by  $\omega(X) = \aleph_0 + \min\{|\mathcal{B}| : \mathcal{B} \text{ is a base for } X\}$ . A collection  $\beta$  of nonempty subsets of a space  $X$  is called a  $\pi$ -base for  $X$  provided that every nonempty open subset of  $X$  contains some member of  $\beta$ . The  $\pi$ -weight of a space  $X$  is defined by  $\pi\omega(X) = \aleph_0 + \min\{|\beta| : \beta \text{ is a } \pi\text{-base for } X\}$ . The density of a space  $X$  is  $d(X) = \aleph_0 + \min\{|D| : D \text{ is a dense subset of } X\}$ . The pseudocharacter of a point  $x \in X$  is  $\psi(x, X) = \aleph_0 + \min\{|\gamma| : \gamma \text{ is a family of open sets in } X \text{ such that } \bigcap \gamma = \{x\}\}$ . The pseudocharacter of a space  $X$  is defined as  $\psi(X) = \sup\{\psi(x, X) : x \in X\}$ . The spread of a space  $X$  is defined as  $s(X) = \aleph_0 + \sup\{|D| : D \subseteq X \text{ is discrete}\}$ . The cellularity or Souslin number of a space  $X$  is defined by  $c(X) = \aleph_0 + \sup\{|\mathcal{U}| : \mathcal{U} \text{ is a family of pairwise disjoint nonempty open subsets of } X\}$ . A space  $X$  is said to have Souslin property if  $c(X) = \aleph_0$ .

**Lemma 4.1** ([8, Lemma 4.2]). *Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$  be a map such that for any  $x \in X$ , there exists an open set  $G$  in  $X$  such that  $x \in \overline{G}$  and  $f(y) = f(x)$  for all  $y \in G$ . Then  $f$  is quasicontinuous.*

**Lemma 4.2** ([8, Lemma 4.3]). *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff. For given  $x_1, \dots, x_n \in X$  and (not necessarily distinct)  $y_1, \dots, y_n \in Y$ , there exists a quasicontinuous map  $f : X \rightarrow Y$  such that  $f(x_i) = y_i$  for each  $i \in \{1, \dots, n\}$ .*

**Theorem 4.3.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is uncountable Hausdorff and  $Y$  is a nontrivial  $T_1$ -space. Then for any  $f \in Q_p(X, Y)$ ,  $f$  does not have a countable local  $\pi$ -base.*

*Proof.* Suppose  $\{U_n : n \in \mathbb{N}\}$  be a countable local  $\pi$ -base at some  $f \in Q_p(X, Y)$ . So for each  $n$ , there is a basic open set  $W_n$  such that  $W_n \subseteq U_n$ . Then  $\{W_n : n \in \mathbb{N}\}$  is also a countable local  $\pi$ -base at  $f$ . Let  $W_n = [x_1^n, \dots, x_{k_n}^n; V_1^n, \dots, V_{k_n}^n]$  for each  $n \in \mathbb{N}$ . The set  $A = \{x_j^i : 1 \leq j \leq k_i, i \in \mathbb{N}\}$  is countable.

Since  $X$  is uncountable, choose  $x \in X \setminus A$ . Because  $Y$  is a nontrivial  $T_1$ -space, choose  $y \in Y$  such that  $y \notin V$  for some open set  $V$  containing  $f(x)$ . Then  $W = [x, V]$  is an open set containing  $f$ . Suppose  $W_n \subseteq W$  for some  $n \in \mathbb{N}$ . By Lemma 4.2, let  $g : X \rightarrow Y$  be a quasicontinuous function such that  $g(x_i^n) \in V_i^n$  for each  $i \in \{1, \dots, k_n\}$  and  $g(x) = y$ . Then  $g \in W_n \setminus W$ , a contradiction. □

**Corollary 4.4.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff and  $Y$  is a nontrivial  $T_1$ -space. If the space  $Q_p(X, Y)$  has a countable local  $\pi$ -base at some  $f \in Q_p(X, Y)$ , then  $X$  is countable.*

**Corollary 4.5.** *Let  $X$  be an uncountable Hausdorff space and  $Y$  be a nontrivial  $T_1$ -space. Then for any  $f \in Q_p(X, Y)$ ,  $f$  does not have a countable local base.*

Using Corollary 4.4, a more general result than [9, Theorem 3.2] is the following.

**Theorem 4.6.** *Let  $X$  and  $Y$  be spaces such that  $X$  is Hausdorff and  $Y$  is a nontrivial metrizable space. Then the following are equivalent:*

- (a)  $F_p(X, Y)$  is metrizable.
- (b)  $F_p(X, Y)$  is first countable.
- (c)  $Q_p(X, Y)$  is metrizable.
- (d)  $Q_p(X, Y)$  is first countable.
- (e) For any  $f \in Q_p(X, Y)$ ,  $f$  has a countable local  $\pi$ -base.
- (f)  $X$  is countable.

*Proof.* Clearly (d) implies (e) holds. The assertion (e) implies (f) follows from Corollary 4.4. Using the facts that  $Q_p(X, Y)$  is a subspace of  $F_p(X, Y)$  and a countable product of metrizable spaces is metrizable, the rest of the implications can be verified easily.  $\square$

The result obtained in Theorem 4.3 can also be deduced from the following result about the cardinal functions related to the space  $Q_p(X, Y)$ .

**Theorem 4.7.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff and  $Y$  is a nontrivial  $T_1$ -space. Then  $|X| \leq \pi_\chi(Q_p(X, Y)) \leq \chi(Q_p(X, Y)) \leq \omega(Q_p(X, Y))$ . Moreover, if  $X$  is infinite and  $Y$  is second countable, we have  $|X| = \pi_\chi(Q_p(X, Y)) = \chi(Q_p(X, Y)) = \pi\omega(Q_p(X, Y)) = \omega(Q_p(X, Y))$ .*

*Proof.* To show  $|X| \leq \pi_\chi(Q_p(X, Y))$ , let  $y_1 \in Y$  and  $f \in Q_p(X, Y)$  be the constant function such that  $f(x) = y_1$  for each  $x \in X$ . Let  $\{U_t : t \in T\}$  be a local  $\pi$ -base at  $f$  with  $|T| \leq \pi_\chi(Q_p(X, Y))$ . Since each  $U_t$  is a nonempty open subset of  $Q_p(X, Y)$ , there exists a basic open set  $B_t = [x_1^t, \dots, x_{n_t}^t; V_1^t, \dots, V_{n_t}^t] \subseteq U_t$  for each  $t \in T$ . Then the collection  $\mathcal{B}_f = \{B_t : t \in T\}$  is also a local  $\pi$ -base at  $f$ . For each  $t \in T$ , let  $A_t = \{x_1^t, \dots, x_{n_t}^t\}$ . We claim that  $\bigcup_{t \in T} A_t = X$ .

Let  $x \in X$ . Since  $Y$  is a nontrivial  $T_1$ -space, choose  $y_2 \in Y$  such that  $y_2 \notin V_1$  for some open set  $V_1$  in  $Y$  containing  $y_1$ . Because  $[x, V_1]$  is an open set containing  $f$  and  $\mathcal{B}_f$  is a local  $\pi$ -base at  $f$ , there exists  $t \in T$  such that  $B_t = [x_1^t, \dots, x_{n_t}^t; V_1^t, \dots, V_{n_t}^t] \subseteq [x, V_1]$ . We claim that  $x \in A_t = \{x_1^t, \dots, x_{n_t}^t\}$ .

Suppose  $x \notin A_t$ . Since  $X$  is Hausdorff, there exists an open set  $U$  such that  $x \in U$  and  $\overline{U} \cap A_t = \emptyset$ . Because  $B_t$  is nonempty, let  $s_t \in B_t$ . Then  $s_t(x_i^t) \in V_i^t$  for each  $i \in \{1, \dots, n_t\}$ . By Lemma 4.2, there is a quasicontinuous map  $h_t : X \rightarrow Y$  such that  $h_t(x_i^t) = s_t(x_i^t)$  for each  $i \in \{1, \dots, n_t\}$ . Let us define  $g : X \rightarrow Y$  such that

$$g(z) = \begin{cases} y_2 & \text{if } z \in \overline{U} \\ h_t(z) & \text{if } z \in X \setminus \overline{U} \end{cases}$$

By Lemma 4.1,  $g$  is a quasicontinuous map such that  $g \in B_t$ , but  $g \notin [x, V_1]$ , which contradicts  $B_t \subseteq [x, V_1]$ . So  $\bigcup_{t \in T} A_t = X$ . Hence  $|X| \leq \pi_\chi(Q_p(X, Y)) \leq \chi(Q_p(X, Y)) \leq \omega(Q_p(X, Y))$ .

If  $\mathcal{B}_Y$  is a countable base for  $Y$  then  $\{[x_1, \dots, x_n; V_1, \dots, V_n] : x_i \in X, V_i \in \mathcal{B}_Y, 1 \leq i \leq n\}$  is a base for the space  $Q_p(X, Y)$ . Thus  $\omega(Q_p(X, Y)) \leq |X|$ .  $\square$

**Theorem 4.8.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is infinite Hausdorff space and  $Y$  is a nontrivial metrizable space. Then  $|X| = \pi_\chi(Q_p(X, Y)) = \chi(Q_p(X, Y))$ .*

*Proof.* By Theorem 4.7, we have  $|X| \leq \pi_\chi(Q_p(X, Y)) \leq \chi(Q_p(X, Y))$ . To show  $\chi(Q_p(X, Y)) \leq |X|$ , let  $f \in Q_p(X, Y)$ . If  $(Y, d)$  is a metric space then the collection  $\mathcal{B}_f = \{O(f, x_1, \dots, x_k, \frac{1}{n}) : x_1, \dots, x_k \in X, k, n \in \mathbb{N}\}$  is a local base at  $f$ . Thus  $|X| = \pi_\chi(Q_p(X, Y)) = \chi(Q_p(X, Y))$ .  $\square$

**Lemma 4.9.** *Let  $X$  and  $Y$  be topological spaces such that  $U_1, \dots, U_n$  are nonempty pairwise disjoint open subsets of  $X$  and  $y_1, \dots, y_n \in Y$ . Then there exists a quasicontinuous map  $g : X \rightarrow Y$  such that  $g(U_i) = \{y_i\}$  for each  $i \in \{1, \dots, n\}$ .*

*Proof.* Let  $H = \overline{U_1} \cup \dots \cup \overline{U_n}$  and  $y_0 \in Y$ . For  $x \in H$ , let  $k = \min\{i \in \{1, \dots, n\} : x \in \overline{U_i}\}$ . Let us define  $g : X \rightarrow Y$  such that

$$g(x) = \begin{cases} y_k & \text{if } x \in H \\ y_0 & \text{if } x \in X \setminus H \end{cases}$$

By Lemma 4.1, the map  $g$  is quasicontinuous and  $g(U_i) = \{y_i\}$  for each  $i \in \{1, \dots, n\}$ .  $\square$

**Theorem 4.10.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff. Then  $d(Q_p(X, Y)) \leq \omega(X) \cdot d(Y)$ .*

*Proof.* Let  $\mathcal{B}$  be a base for  $X$  such that  $|\mathcal{B}| \leq \omega(X)$  and  $\mathcal{U}$  be the family of all finite pairwise disjoint nonempty members of  $\mathcal{B}$ . Let  $D$  be a dense set in  $Y$  such that  $|D| \leq d(Y)$  and  $\mathcal{V}$  be the family of all nonempty finite subsets of  $D$ . For each  $U = \{U_1, \dots, U_n\} \in \mathcal{U}$  and  $y = \{y_1, \dots, y_n\} \in \mathcal{V}$ , by Lemma 4.9, there exists a quasicontinuous function  $g_{U,y} : X \rightarrow Y$  such that  $g_{U,y}(U_i) = y_i$  for each  $i \in \{1, \dots, n\}$ . Then  $G = \{g_{U,y} : U \in \mathcal{U}, y \in \mathcal{V}\}$  is dense set in  $Q_p(X, Y)$  such that  $|G| \leq \omega(X) \cdot d(Y)$ .

Indeed, for any nonempty basic open set  $H = [x_1, \dots, x_n; V_1, \dots, V_n]$  in  $Q_p(X, Y)$ , there exist  $U = \{U_1, \dots, U_n\} \in \mathcal{U}$  such that  $x_i \in U_i$  and  $y = \{y_1, \dots, y_n\} \in \mathcal{V}$  such that  $y_i \in V_i$  for each  $i \in \{1, \dots, n\}$ . Thus there is  $g_{U,y} \in G$  such that  $g_{U,y}(x_i) \in V_i$  for each  $i \in \{1, \dots, n\}$  and hence  $g_{U,y} \in H \cap G$ .  $\square$

**Corollary 4.11.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is second countable Hausdorff and  $Y$  is separable. Then the space  $Q_p(X, Y)$  is separable.*

**Lemma 4.12.** *Let  $X$  and  $Y$  be spaces such that  $X$  is regular. Then for any  $x \in X$ , any nonempty closed set  $F \subseteq X$  such that  $x \notin F$  and  $y_1, y_2 \in Y$ , there exists a quasicontinuous function  $f : X \rightarrow Y$  such that  $f(x) = y_1$  and  $f(F) = \{y_2\}$ .*

*Proof.* Since  $x \notin F$  and  $X$  is regular, there exist open sets  $U$  and  $V$  such that  $x \in U$ ,  $F \subseteq V$  and  $U \cap V = \emptyset$ . Note that  $x \notin \overline{V}$ . Let us define  $f : X \rightarrow Y$

such that

$$f(z) = \begin{cases} y_1 & \text{if } z \in X \setminus \bar{V} \\ y_2 & \text{if } z \in \bar{V} \end{cases}$$

By Lemma 4.1,  $f$  is a quasicontinuous map such that  $f(x) = y_1$  and  $f(F) = \{y_2\}$ .  $\square$

**Theorem 4.13.** *Let  $X$  be a regular space and  $Y$  be any nontrivial space. Then  $d(X) \leq \psi(Q_p(X, Y))$ .*

*Proof.* Given a basic open set  $U = [x_1, \dots, x_n; V_1, \dots, V_n]$  in  $Q_p(X, Y)$ , let  $A_U = \{x_1, \dots, x_n\}$ . Let  $f_0 \in Q_p(X, Y)$  be the constant function such that  $f_0(x) = y_0$  for all  $x \in X$  and  $\gamma$  be a family of open sets with  $|\gamma| \leq \psi(Q_p(X, Y))$  such that  $\cap \gamma = \{f_0\}$ . For each  $G \in \gamma$ , there exists a basic open set  $U_G = [x_1^G, \dots, x_{n_G}^G; V_1^G, \dots, V_{n_G}^G]$  such that  $f_0 \in U_G \subseteq G$ . We claim that the set  $D = \bigcup \{A_{U_G} : G \in \gamma\}$  is dense in  $X$ .

Suppose that  $x \in X \setminus \bar{D}$  and  $y_1 \in Y$  such that  $y_1 \neq y_0$ . By Lemma 4.12, there exists  $f \in Q_p(X, Y)$  such that  $f(x) = y_1$  and  $f(\bar{D}) = \{y_0\}$ . Then  $f \in \cap \gamma$  and  $f \neq f_0$ , which is a contradiction. Thus  $D$  is dense in  $X$ .  $\square$

Note that if  $X$  is a Tychonoff space, then the result obtained in Theorem 4.13 for  $Y = \mathbb{R}$  can be concluded from the results  $d(X) = \psi(C_p(X, \mathbb{R}))$  [17, Problem 173] and  $\psi(C_p(X, \mathbb{R})) \leq \psi(Q_p(X, \mathbb{R}))$  [17, Problem 159]. We cannot expect the equality in between  $d(X)$  and  $\psi(Q_p(X, \mathbb{R}))$  even for  $X = \mathbb{R}$ , because  $d(\mathbb{R}) = \aleph_0$ , while [8, Example 5.1] shows that  $\psi(Q_p(\mathbb{R}, \mathbb{R})) = 2^{\aleph_0}$ .

**Theorem 4.14.** *Let  $X$  be a regular space and  $Y$  be a nontrivial  $T_1$ -space. Then  $s(X) \leq s(Q_p(X, Y))$ .*

*Proof.* Let  $D$  be a discrete subspace of  $X$  and  $\{V_d : d \in D\}$  be a family of open subsets of  $X$  such that  $V_d \cap D = \{d\}$  for each  $d \in D$ . Choose  $y_1, y_2 \in Y$  such that  $y_1 \neq y_2$ , by Lemma 4.12, there exists a quasicontinuous function  $f_d : X \rightarrow Y$  such that  $f_d(d) = y_1$  and  $f_d(X \setminus V_d) = \{y_2\}$ . Then the set  $A = \{f_d : d \in D\}$  is discrete in  $Q_p(X, Y)$ . To see this, choose an open set  $G$  in  $Y$  such that  $y_1 \in G$  but  $y_2 \notin G$ . Then  $U_d = [d, G]$  is open in  $Q_p(X, Y)$  and  $U_d \cap A = \{f_d\}$ . Thus  $s(X) \leq s(Q_p(X, Y))$ .  $\square$

If  $X$  is a Tychonoff space and  $Y = \mathbb{R}$ , then the result obtained in Theorem 4.14 can be obtained from the results  $s(X) \leq s(C_p(X, \mathbb{R}))$  [17, Problem 176] and  $s(C_p(X, \mathbb{R})) \leq s(Q_p(X, \mathbb{R}))$  [17, Problem 159].

**Theorem 4.15.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff. Then  $Q(X, Y)$  is dense in  $F_p(X, Y)$ .*

*Proof.* Let  $W = [x_1, \dots, x_n; V_1, \dots, V_n]$  be any nonempty basic open set in  $F_p(X, Y)$  and  $f \in W$ . Then  $f(x_i) \in V_i$  for each  $i \in \{1, \dots, n\}$ . Since  $X$  is Hausdorff and  $x_1, \dots, x_n \in X$ , by Lemma 4.2, there exists a quasicontinuous function  $g : X \rightarrow Y$  such that  $g(x_i) = f(x_i)$  for each  $i \in \{1, \dots, n\}$ . Then  $g \in W \cap Q(X, Y)$ .  $\square$

**Corollary 4.16.** *Let  $X$  be a Hausdorff space and  $Y$  be a separable space. Then the space  $Q_p(X, Y)$  has the Souslin property, that is,  $c(Q_p(X, Y)) = \aleph_0$ .*

*Proof.* It is known that if  $Y$  is separable then the space  $F_p(X, Y)$  has the Souslin property [17, Problem 109]. Since  $Q(X, Y)$  is dense in  $F_p(X, Y)$ ,  $c(Q_p(X, Y)) = c(F_p(X, Y))$  [17, Problem 110]. Thus the space  $Q_p(X, Y)$  has the Souslin property.  $\square$

Note that if  $X$  is a Tychonoff space then  $C(X, \mathbb{R})$  is a dense subset of the space  $Q_p(X, \mathbb{R})$  [17, Problem 034]. Also  $P(\mathbb{R}, \mathbb{R})$ , the set of all polynomials from  $\mathbb{R}$  to  $\mathbb{R}$  and  $U(\mathbb{R}, \mathbb{R})$ , the set of all uniformly continuous functions from  $\mathbb{R}$  to  $\mathbb{R}$  are dense subsets of the space  $Q_p(\mathbb{R}, \mathbb{R})$  [17, Problem 041,043].

**Proposition 4.17.** *Let  $X$  and  $Y$  be topological spaces such that  $X$  is Hausdorff and  $Y$  is separable. If  $\mathcal{F}$  is any locally finite family of nonempty open subsets of  $Q_p(X, Y)$ , then  $\mathcal{F}$  is countable.*

*Proof.* If possible, suppose  $\mathcal{F}$  is uncountable. Let  $\mathcal{A}$  be a maximal disjoint family of nonempty open subsets of  $Q_p(X, Y)$  such that each member of  $\mathcal{A}$  meets at most finitely many members of  $\mathcal{F}$ . Because  $\mathcal{F}$  is locally finite, the set  $\bigcup \mathcal{A}$  is dense in  $Q_p(X, Y)$ . By Corollary 4.16,  $c(Q_p(X, Y)) \leq \aleph_0$ , which implies  $\mathcal{A}$  is countable. Since  $\bigcup \mathcal{A}$  is dense in  $Q_p(X, Y)$ , each  $U \in \mathcal{F}$  intersect some  $V \in \mathcal{A}$ . But every member of  $\mathcal{A}$  can intersect only finitely many members of  $\mathcal{F}$ . Since  $\mathcal{A}$  is countable, this implies  $\mathcal{F}$  is countable, a contradiction.  $\square$

### 5. SPECIAL MAPS AND THE SPACE $Q_p(X, Y)$

The properties of induced maps related to the space  $C(X, Y)$  with the topology of pointwise convergence and others are discussed in [14, Chapter II]. Before discussing the properties of induced maps related to the space  $Q_p(X, Y)$ , let us first define these maps in view of quasicontinuous maps.

Note that the composition of two quasicontinuous maps need not be quasicontinuous [16]. However, if  $f : X \rightarrow Y$  is quasicontinuous and  $g : Y \rightarrow Z$  is continuous, then the composition map  $g \circ f : X \rightarrow Z$  is quasicontinuous. If  $g : Y \rightarrow Z$  is continuous, then the induced map  $g_* : Q(X, Y) \rightarrow Q(X, Z)$  is defined by  $g_*(f) = g \circ f$  for all  $f \in Q(X, Y)$ . Also if  $g \in Q(X, Y)$ , then the induced map  $g^* : C(Y, Z) \rightarrow Q(X, Z)$  is defined as  $g^*(h) = h \circ g$  for all  $h \in C(Y, Z)$ .

**Theorem 5.1.** *For a given continuous map  $g : Y \rightarrow Z$ , the induced map  $g_* : Q_p(X, Y) \rightarrow Q_p(X, Z)$  such that  $g_*(f) = g \circ f$  is continuous. Moreover, if  $g$  is an embedding, then  $g_*$  is also an embedding.*

*Proof.* Let  $f \in Q_p(X, Y)$  and  $U$  be any open set in  $Q_p(X, Z)$  containing  $g_*(f)$ . There is a basic open set  $V = [x_1, \dots, x_n; V_1, \dots, V_n]$  in  $Q_p(X, Z)$  such that  $g_*(f) \in V \subseteq U$ . Now  $W = [x_1, \dots, x_n; g^{-1}(V_1), \dots, g^{-1}(V_n)]$  is an open set in  $Q_p(X, Y)$  such that  $f \in W$  and  $g_*(W) \subseteq V \subseteq U$ . Thus the map  $g_*$  is continuous.



Note that if  $g$  is injective then  $g_*$  is also injective. Now to show  $g_* : Q_p(X, Y) \rightarrow g_*(Q_p(X, Y))$  is an open map, let  $[x, V]$  be any subbasic open set in  $Q_p(X, Y)$ . Since  $g$  is an embedding and  $V$  is open in  $Y$ , there exists an open set  $W$  in  $Z$  such that  $g(V) = W \cap g(Y)$ . We have  $[x, V] = [x, g^{-1}(W)] = g_*^{-1}([x, W])$ . Then  $g_*([x, V]) = [x, W] \cap g_*(Q_p(X, Y))$  is open in  $g_*(Q_p(X, Y))$ .  $\square$

**Proposition 5.2.** *For any space  $X$ , there is a continuous map  $h : Q_p(X, \mathbb{R}) \rightarrow Q_p(X, \mathbb{I})$  such that  $h(f) = f$  for each  $f \in Q_p(X, \mathbb{I})$ .*

*Proof.* Consider the map  $g : \mathbb{R} \rightarrow \mathbb{I}$  such that  $g(t) = -1$  if  $t < -1$ ,  $g(t) = t$  if  $t \in \mathbb{I} = [-1, 1]$  and  $g(t) = 1$  if  $t > 1$ . Clearly  $g$  is continuous. By Theorem 5.1, the map  $h = g_* : Q_p(X, \mathbb{R}) \rightarrow Q_p(X, \mathbb{I})$  defined by  $h(f) = gof$  is continuous. Also  $h(f) = f$  for each  $f \in Q_p(X, \mathbb{I})$ .  $\square$

**Theorem 5.3.** *For a given quasicontinuous map  $g : X \rightarrow Y$ , the map  $g^* : C_p(Y, Z) \rightarrow Q_p(X, Z)$  such that  $g^*(h) = h \circ g$  is continuous. Moreover, if  $g(X) = Y$ , then  $g^*$  is an embedding.*

*Proof.* Let  $h_0 \in C_p(Y, Z)$  and  $V = [x_1, \dots, x_n; V_1, \dots, V_n]$  be any basic open set in  $Q_p(X, Z)$  containing  $g^*(h_0)$ . Consider  $U = [g(x_1), \dots, g(x_n); V_1, \dots, V_n]$  open in  $C_p(Y, Z)$ . Then  $h_0 \in U$  and for any  $h \in U$ , we have  $g^*(h) \in V$ . Thus  $g^*(U) \subseteq V$  and hence  $g^*$  is continuous.

Now suppose that  $g(X) = Y$ . To see  $g^*$  is an injection, let  $h, h' \in C_p(Y, Z)$  such that  $h \neq h'$ . Then  $h(y) \neq h'(y)$  for some  $y \in Y$ . Because  $g(X) = Y$ , let  $x \in g^{-1}(y)$ . Then  $g^*(h)(x) = h(y) \neq h'(y) = g^*(h')(x)$ . Hence  $g^*(h) \neq g^*(h')$ . To prove  $g^*$  is an embedding, it suffices to show that  $(g^*)^{-1} : g^*(C_p(Y, Z)) \rightarrow C_p(Y, Z)$  is continuous. Let  $g^*(f) \in g^*(C_p(Y, Z))$  and  $U = [y_1, \dots, y_n; V_1, \dots, V_n]$  be any basic open set in  $C_p(Y, Z)$  containing  $f$ . Choose  $x_i \in g^{-1}(y_i)$  for each  $i \in \{1, \dots, n\}$ . Then  $V = [x_1, \dots, x_n; V_1, \dots, V_n] \cap g^*(C_p(Y, Z))$  is open in  $g^*(C_p(Y, Z))$  containing  $g^*(f)$ . To verify  $(g^*)^{-1}(V) \subseteq U$ , let  $h \in V$ . Then  $h = g^*(h')$  for some  $h' \in C_p(Y, Z)$ . Since  $h = g^*(h') = h' \circ g \in V$ , we have  $h' \circ g(x_i) \in V_i$  for each  $i \in \{1, \dots, n\}$ . This implies  $h'(y_i) \in V_i$  for each  $i \in \{1, \dots, n\}$  so that  $h' \in U$ . Hence  $h' = (g^*)^{-1}(h) \in U$  and we have  $(g^*)^{-1}(V) \subseteq U$ .  $\square$

For any space  $X$  and maps  $f, g : X \rightarrow \mathbb{R}$  such that  $f$  is continuous and  $g$  is quasicontinuous, it is easy to see that the map  $f + g : X \rightarrow \mathbb{R}$  defined by  $(f + g)(x) = f(x) + g(x)$  is quasicontinuous.

**Proposition 5.4.** *For any space  $X$ , the map  $s : C_p(X, \mathbb{R}) \times Q_p(X, \mathbb{R}) \rightarrow Q_p(X, \mathbb{R})$  defined by  $s(f, g) = f + g$  is continuous.*

*Proof.* Let  $(f_0, g_0) \in C_p(X, \mathbb{R}) \times Q_p(X, \mathbb{R})$  and  $U$  be any open set in  $Q_p(X, \mathbb{R})$  containing  $h_0 = f_0 + g_0$ . There exist  $x_1, \dots, x_n \in X$  and  $\epsilon > 0$  such that  $h_0 \in O(h_0, x_1, \dots, x_n, \epsilon) \subseteq U$ . Then  $V = O(f_0, x_1, \dots, x_n, \frac{\epsilon}{2})$  and  $W = O(g_0, x_1, \dots, x_n, \frac{\epsilon}{2})$  are open in  $C_p(X, \mathbb{R})$  and  $Q_p(X, \mathbb{R})$ , respectively. Therefore  $V \times W$  is open in  $C_p(X, \mathbb{R}) \times Q_p(X, \mathbb{R})$  containing  $(f_0, g_0)$ . We claim that

$s(V \times W) \subseteq U$ . For this, let  $s(f, g) = f + g \in s(V \times W)$ , then  $|f(x_i) + g(x_i) - h_0(x_i)| \leq |f(x_i) - f_0(x_i)| + |g(x_i) - g_0(x_i)| < \epsilon$  for all  $i \in \{1, \dots, n\}$ . Thus  $f + g \in O(h_0, x_1, \dots, x_n, \epsilon) \subseteq U$ .  $\square$

**Lemma 5.5.** *For any  $x \in X$ , the evaluation map at  $x$ ,  $e_x : Q_p(X, Y) \rightarrow Y$  defined by  $e_x(f) = f(x)$  is continuous.*

*Proof.* Let  $f \in Q_p(X, Y)$  and  $V$  be any open set in  $Y$  containing  $f(x)$ . Then  $U = [x, V]$  is an open set containing  $f$  such that  $e_x(U) \subseteq V$ . Thus  $e_x$  is continuous.  $\square$

For any space  $X$  and  $A \subseteq X$ , a family  $\mathcal{B}_A$  of open subsets of  $X$  is called a base at  $A$  [5] if each member of  $\mathcal{B}_A$  contains  $A$  and for any open set  $U$  containing  $A$ , there exists  $B \in \mathcal{B}_A$  such that  $B \subseteq U$ . The character of  $A$  in  $X$  is defined as  $\chi(A, X) = \aleph_0 + \min\{|\mathcal{B}_A| : \mathcal{B}_A \text{ is a base at } A\}$ . Note that  $\chi(\{x\}, X) = \chi(x, X)$ .

**Proposition 5.6.** *Let  $X$  be a Hausdorff space. If there exists a compact subspace  $K$  of the space  $Q_p(X, \mathbb{R})$  such that  $\chi(K, Q_p(X, \mathbb{R})) \leq \aleph_0$ , then  $X$  is countable.*

*Proof.* Given a basic open set  $U = [x_1, \dots, x_n; V_1, \dots, V_n]$  in  $Q_p(X, \mathbb{R})$ , let  $A_U = \{x_1, \dots, x_n\}$ . Suppose that  $\{B_n : n \in \mathbb{N}\}$  is a countable base at  $K$  in  $Q_p(X, \mathbb{R})$ . Fix  $n \in \mathbb{N}$ , for each  $f \in K$ , choose a basic open set  $U_f^n$  such that  $f \in U_f^n \subseteq B_n$ . For open cover  $\{U_f^n : f \in K\}$  of  $K$ , choose a finite subcover  $\{U_{f_1}^n, \dots, U_{f_{m_n}}^n\}$  for some  $m_n \in \mathbb{N}$ . Let  $W_n = U_{f_1}^n \cup \dots \cup U_{f_{m_n}}^n$  and  $A_n = A_{U_{f_1}^n} \cup \dots \cup A_{U_{f_{m_n}}^n}$ , then  $K \subseteq W_n \subseteq B_n$ . Clearly  $A = \bigcup\{A_n : n \in \mathbb{N}\}$  is countable. We claim that  $A = X$ .

Suppose that  $x \in X \setminus A$ . By Lemma 5.5, the map  $e_x : Q_p(X, \mathbb{R}) \rightarrow \mathbb{R}$  defined by  $e_x(f) = f(x)$  is continuous. Therefore the set  $e_x(K)$  is bounded in  $\mathbb{R}$ . Choose  $M > 0$  such that  $|f(x)| < M$  for all  $f \in K$ . Since  $W = [x, (-M, M)]$  is an open set containing  $K$ , there exists  $k \in \mathbb{N}$  such that  $K \subseteq B_k \subseteq W$  and hence  $W_k = U_{f_1}^k \cup \dots \cup U_{f_{m_k}}^k \subseteq W$ . Thus  $U_{f_1}^k = [x_1, \dots, x_n; V_1, \dots, V_n] \subseteq W$  such that  $x \notin \{x_1, \dots, x_n\}$ . Since  $X$  is Hausdorff, by Lemma 4.2, choose  $g \in Q_p(X, \mathbb{R})$  such that  $g(x_i) \in V_i$  for each  $i \in \{1, \dots, n\}$  and  $g(x) = M$ . Then  $g \in W_k \setminus W$ , which is a contradiction.  $\square$

The properties of the restriction map related to the space  $C(X, \mathbb{R})$  with the topology of pointwise convergence are discussed in [1]. For  $Y \subseteq X$ , the restriction map is defined as  $\pi_Y : F(X, Z) \rightarrow F(Y, Z)$  such that  $\pi_Y(f) = f|_Y$  for all  $f \in F(X, Z)$ . Note that the restriction of a quasicontinuous map on an open or a dense subset is quasicontinuous [16]. A map  $f : X \rightarrow Y$  is called almost onto if  $f(X)$  is dense in  $Y$ .

**Proposition 5.7.** *Let  $X$  be a regular space and  $Y$  be an open subset of  $X$ . If the map  $\pi_Y : Q(X, \mathbb{R}) \rightarrow Q(Y, \mathbb{R})$  such that  $\pi_Y(f) = f|_Y$  is injective, then  $Y$  is dense in  $X$ .*

*Proof.* Let  $h_0 \in Q(X, \mathbb{R})$  such that  $h_0(x) = 0$  for all  $x \in X$ . Suppose that  $\pi_Y$  is injective but  $Y$  is not dense in  $X$  so that  $z \in X \setminus \overline{Y}$ . By Lemma 4.12, there exists

$h \in Q(X, \mathbb{R})$  such that  $h(z) = 1$  and  $h(\overline{Y}) = \{0\}$ . We have  $\pi_Y(h) = \pi_Y(h_0)$  but  $h \neq h_0$ , which is a contradiction. Hence  $Y$  is dense in  $X$ .  $\square$

**Theorem 5.8.** *Let  $X$  be a Hausdorff space and  $Y \subseteq X$  be open or dense in  $X$ . Then the restriction map  $\pi_Y : Q_p(X, \mathbb{R}) \rightarrow Q_p(Y, \mathbb{R})$  such that  $\pi_Y(f) = f|_Y$  is continuous and almost onto. Moreover,  $\pi_Y$  is a homeomorphism if and only if  $Y = X$ .*

*Proof.* Consider the natural projection  $p_Y : \mathbb{R}^X \rightarrow \mathbb{R}^Y$  such that  $p_Y(x) = x|_Y$ . Then  $p_Y$  is a continuous map [17, Problem 107] and  $\pi_Y = p_Y|_{Q_p(X, \mathbb{R})}$ . Therefore  $\pi_Y$  is continuous. By Theorem 4.15,  $Q(X, \mathbb{R})$  is dense in  $\mathbb{R}^X$ . Since  $p_Y$  is continuous,  $\mathbb{R}^Y = p_Y(\mathbb{R}^X) = p_Y(\overline{Q_p(X, \mathbb{R})}) \subseteq \overline{p_Y(Q_p(X, \mathbb{R}))}$ . Thus  $\pi_Y(Q_p(X, \mathbb{R})) = p_Y(Q_p(X, \mathbb{R}))$  is dense in  $\mathbb{R}^Y$  and hence also dense in  $Q_p(Y, \mathbb{R})$ .

Now if  $\pi_Y$  is a homeomorphism and  $Y \neq X$ . For  $x \in X \setminus Y$ , the set  $D = \{f \in Q_p(X, \mathbb{R}) : f(x) = 0\}$  is not dense in  $Q_p(X, \mathbb{R})$ , because  $D \cap [x, (0, 1)] = \emptyset$ . But  $\pi_Y(D)$  is dense in  $Q_p(Y, \mathbb{R})$ . Let  $G = [y_1, \dots, y_n; V_1, \dots, V_n]$  be any basic open set in  $Q_p(Y, \mathbb{R})$  containing some  $g$ . By Lemma 4.2, there exists  $f \in Q_p(X, \mathbb{R})$  such that  $f(y_i) = g(y_i)$  and  $f(x) = 0$ . Then  $f \in D$  such that  $\pi_Y(f) \in G$ . Hence  $\pi_Y(D) \cap G \neq \emptyset$ . Because the image of a dense set  $\pi_Y(D)$  under the map  $(\pi_Y)^{-1}$  is  $D$ , which is not dense. This implies that  $(\pi_Y)^{-1}$  is not continuous, which is a contradiction. Finally, if  $Y = X$  then  $\pi_Y$  is the identity map, and hence a homeomorphism.  $\square$

ACKNOWLEDGEMENTS. *The first author acknowledges the fellowship grant of University Grant Commission, India with Student-ID DEC18-414765. The authors are thankful to the anonymous referee for his valuable comments and suggestions.*

## REFERENCES

- [1] A. V. Arhangel'skii, Topological Function Spaces, Mathematics and its Applications (Soviet Series), vol. 78, Kluwer Academic Publishers Group, Dordrecht, 1992.
- [2] J. Borsík, Points of continuity, quasicontinuity and cliquishness, Rend. Istit. Mat. Univ. Trieste 26 (1994), 5–20.
- [3] R. Cazacu and J. D. Lawson, Quasicontinuous functions, domains, and extended calculus, Appl. Gen. Topol. 8 (2007), 1–33.
- [4] A. Crannell, M. Frantz and M. LeMasurier, Closed relations and equivalence classes of quasicontinuous functions, Real Anal. Exchange 31 (2005/06), 409–424.
- [5] R. Engelking, General Topology, Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989.
- [6] Ľ. Holá and D. Holý, Minimal USCO maps, densely continuous forms and upper semi-continuous functions, Rocky Mountain J. Math. 39 (2009), 545–562.
- [7] Ľ. Holá and D. Holý, Pointwise convergence of quasicontinuous mappings and Baire spaces, Rocky Mountain J. Math. 41 (2011), 1883–1894.

- [8] Ľ. Holá and D. Holý, Quasicontinuous functions and the topology of pointwise convergence, *Topology Appl.* 282 (2020), Article No. 107301.
- [9] D. Holý and L. Matejíčka, Quasicontinuous functions, minimal USCO maps and topology of pointwise convergence, *Math. Slovaca* 60 (2010), 507–520.
- [10] S. Kempisty, Sur les fonctions quasicontinues, *Fundamenta Mathematicae* 19 (1932), 184–197.
- [11] P. S. Kenderov, I. S. Kortezov and W. B. Moors, Topological games and topological groups, *Topology Appl.* 109 (2001), 157–165.
- [12] P. S. Kenderov, I. S. Kortezov and W. B. Moors, Continuity points of quasi-continuous mappings, *Topology Appl.* 109 (2001), 321–346.
- [13] N. Levine, Semi-open sets and semi-continuity in topological spaces, *Amer. Math. Monthly* 70 (1963), 36–41.
- [14] R. A. McCoy and I. Ntantu, *Topological Properties of Spaces of Continuous Functions*, Lecture Notes in Mathematics, vol. 1315, Springer-Verlag, Berlin, 1988.
- [15] T. Neubrunn, Quasi-continuity, *Real Anal. Exchange* 14 (1988/89), 259–306.
- [16] Z. Piotrowski, A survey of results concerning generalized continuity of topological spaces, *Acta Math. Univ. Comenian.* 52/53 (1987), 91–110.
- [17] V. V. Tkachuk, *A  $C_p$ -Theory Problem Book*, Topological and function spaces, Problem Books in Mathematics, Springer, New York, 2011.