

ORNITHOPHAGY IN *Rhinella icterica* (SPIX, 1824)ORNITOFAGIA EM *Rhinella icterica* (SPIX, 1824)

**Vagner Luis CAMILOTTI<sup>1</sup>; André Felipe BARRETO-LIMA<sup>2</sup>**

1. Doutorando, Centro de Ciéncia do Sistema Terrestre, Instituto Nacional de Pesquisas Espaciais – INPE, São José dos Campos, SP. [vagner.camilotti@gmail.com](mailto:vagner.camilotti@gmail.com); 2. Doutorando, Centro de Ecologia, Universidade Federal do Rio Grande do Sul – UFRGS, Porto Alegre, RS. [afblima@hotmail.com](mailto:afblima@hotmail.com)

**ABSTRACT:** In this work we report an observation of an ornitophagy event on *Certhiaxis cinnamomeus* by *Rhinella icterica* occurred in the central region of Rio Grande do Sul, Brazil.

**KEYWORDS:** Ornithophagy. *Rhinella icterica*. *Certhiaxis cinnamomeus*.

The yellow cururu toad, *Rhinella icterica* (SPIX, 1824) (Anura, Bufonidae, *Rhinella marina* group), occurs in central, southeastern and southern Brazil, Misiones and Corrientes, Northeastern Argentina, and Eastern Paraguay (KWET; DI-BERNARDO, 1999; SILVANO et al., 2010; FROST et al., 2011). The body length of *R. icterica* changes from 100 to 166 mm in males, and 135 to 190 mm in females (KWET; DI-BERNARDO, 1999). This species uses several kinds of habitats, from forested to open areas and it breeds in permanent or temporary ponds and streams, mainly in substantially disturbed habitats (SILVANO et al., 2010).

Bufonidae has been classified as ant-specialist family by some authors (TOFT, 1980; FLOWERS; GRAVES, 1995; ISACCH; BARG, 2002; ROSA et al., 2002) and as generalists by others (SMITH; BRAGG, 1949; EVANS; LAMPO, 1996; GRANT, 1996; FERREIRA; TEIXEIRA, 2009; BATISTA et al., 2011). Although some *Rhinella* species are opportunistic predators, including small vertebrates in their diet (ZUG; ZUG, 1979; MATTISON, 1994), Sabagh and Carvalho-e-Silva (2008) found only arthropods in the diet of *R. icterica* in southeastern Brazil.

During a field expedition on March 2010 in the central region (29°41'S, 53°48'W) of the State

of Rio Grande do Sul, Brazil, we observed an adult of *R. icterica* predating an adult of yellow-chinned Spinetail, *Certhiaxis cinnamomeus* (GMELIN, 1778) (posicionamento taxonómico). The size of *C. cinnamomeus* changes from 150 to 168 mm, and weigh from 15 to 16.5 g (BELTON, 1994; SICK, 1997). This bird is a common resident breeder of marches and edges of mangrove swamps (BELTON, 1994; SICK, 1997).

The predation event occurred at the margin of a pond covered by high grasses and shrubs. An adult bird with a broken wing was walking on the ground when, opportunistically, *R. icterica* jumped on and swallowed it. Although the bird injury could be allowed the predation event, this species also forage on or next to the ground, creating opportunity for terrestrial and opportunistic predators as *R. icterica*. There are records of *Rhinella marina* predation on eggs and nestlings of ground-nesting birds (BOLAND, 2004). Finally, we do not know any similar case reported before in literature about ornitophagy in *R. icterica*.

**ACKNOWLEDGMENTS**

We thank to Célio F. B. Haddad and Reuber A. Brandão who considered relevant this report and encouraged us to publish it.

**RESUMO:** Reportamos um evento de ornitofagia por *Rhinella icterica* sobre *Certiaxis cinnamomeus* ocorrido em 2010, na região central do Estado do Rio Grande do Sul (29°41'S, 53°48'W), Brasil.

**PALAVRAS-CHAVE:** Ornitofagia. *Rhinella icterica*. *Certiaxis cinnamomeus*.

**REFERENCES**

BATISTA, R. C.; DE-CARVALHO, C. B.; FREITAS, E. B.; FRANCO, S. C.; BATISTA, C. C.; COELHO, W. A.; FARIA, R. G. Diet of *Rhinella schneideri* (Werner, 1894) (Anura: Bufonidae) in the Cerrado, Central Brazil. *Herpetol. Notes*, Berlim, v. 4, p. 17-21, 2011.

- BELTON, W. **Aves do Rio Grande do Sul: Distribuição e biologia.** São Leopoldo: Ed. Unisinos, 1994. 584p.
- BOLAND, C. R. J. Introduced cane toads are active nest predators and competitors of rainbow bee-eaters: observational and experimental evidence. **Biol. Conserv.**, Amsterdam, v. 120, p. 53–62, 2004.
- EVANS, M.; LAMPO, M. Diet of *Bufo marinus* in Venezuela. **J. Herpetol.**, Salt Lake City, v. 30, p. 73-76, 1996.
- FERREIRA, R. B.; TEIXEIRA, R. L. Feeding pattern and use of reproductive habitat of the Striped toad *Rhinella crucifer* (Anura: Bufonidae) from Southeastern Brazil. **Acta Herpetol.**, Firenze, v. 4, p. 125-134, 2009
- FLOWERS, M. A.; GRAVES, B.M. Prey selectivity and size-specific diet changes in *Bufo cognatus* and *B. woodhousii* during early postmetamorphic ontogeny. **J. Herpetol.**, Salt Lake City, v. 29, p. 608-612, 1995.
- FROST, D. **Amphibian Species of the World: an Online Reference.** Version 5.5 (31 January, 2011). American Museum of Natural History, New York, USA, 2011. Available at: <http://research.amnh.org/vz/herpetology/amphibia>. Access on 25/02/2011.
- GRANT, G. S. Prey of the introduced *Bufo marinus* on American Samoa. **Herp. Rev.**, Salt Lake City, v. 27, p. 67-69, 1996.
- ISACCH, J. P.; BARG, M. Are bufonid toads specialized ant-feeders? A case test from the Argentinian flooding pampa. **J. Nat. Hist.**, London, v. 36, p. 2005-2012, 2002.
- KWET, A.; DI-BERNARDO, M. **Pro-Mata - Anfibios. Amphibien. Amphibians.** Porto Alegre: EDIPUCRS, 1999. 107p.
- MATTISON, C. **Frogs and toads of the word.** New York: Facts on File Inc., 1994. 192p.
- ROSA, I.; CANAVERO, A.; MANEYRO, R.; NAYA, D. E.; CAMARGO, A. Diet of four sympatric anuran species in a temperate environment. **Bol. Soc. Zool. Uruguay**, Montevideo, v. 13, p. 12-20, 2002.
- SABAGH, L. T.; CARVALHO-E-SILVA, A. M. P. T. Feeding overlap in two sympatric species of *Rhinella* (Anura: Bufonidae) of the Atlantic Rain Forest. **Rev. Bras. Zool.**, Curitiba, v. 25, p. 247-253, 2008.
- SICK, H. **Ornitologia brasileira.** Rio de Janeiro: Nova Fronteira, 1997. 914p.
- SILVANO, D.; SCOTT, N.; AQUINO, L.; KWET, A.; BALDO, D. *Rhinella icterica*. In: IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4, 2008. Available at: [www.iucnredlist.org](http://www.iucnredlist.org). Access on 25/02/2011.
- SMITH, C. C.; BRAGG, A. N. Observations on the ecology and natural history of anura, VII. Food and feeding habitats of the common species of toads in Oklahoma. **Ecology**, New York, v. 30, p. 333-349, 1949.
- TOFT, C. A. Feeding ecology of Panamanian litter anurans: patterns in diet and forage mode. **J. Herpetol.**, Salt Lake City, v. 15, p. 139-144, 1981.
- ZUG, G. R.; ZUG, P. B. The marine toad, *Bufo marinus*: a natural history resumé of native populations. **Smiths. Contrib. Zool.**, Washington, v. 284, p. 1-58, 1979.