

---

# Biological action of *Piper nigrum* - the king of spices

Arun Kumar Srivastava\*, Vinay Kumar Singh

Malacology Laboratory, Department of Zoology, DDU Gorakhpur University, Gorakhpur - 273009 U.P. India

\* Corresponding author: Dr. Arun Kumar Srivastava; Phone: +91-9792250710 (Mobile); E-mail: aksgkp5@gmail.com

---

Received: 17 May 2017; Revised submission: 25 July 2017; Accepted: 03 August 2017

Copyright: © The Author(s) 2017. European Journal of Biological Research © T.M.Karpiński 2017. This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial 4.0 International License, which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.

DOI: <http://dx.doi.org/10.5281/zenodo.839039>

---

## ABSTRACT

*Piper nigrum* - the king of spices is originated in the Western Ghats of India. It has gained a global consideration because of its volume in the spice industry. It contains major pungent alkaloid piperine which is known to possess many interesting pharmacological actions. Medicinally black pepper can be used digestive disorder like large intestine toxins, different gastric problems, diarrhoea and indigestion and also can be used against respiratory disorder including cold fever, asthma. Piperine exhibits diverse pharmacological activities like antihypertensive, antiplatelets, antioxidant, antitumor, antipyretic, analgesic, anti-inflammatory, anti-diarrheal, antibacterial, antifungal, anti-reproductive, insecticidal activities. *Piper nigrum* also found to decrease lipid peroxidation *in vivo*. It has reported to possess antioxidant activity that might be due to the presence of flavonoids and phenolic contents.

**Keywords:** *Piper nigrum*; Alkaloids; Antireproductive; Antioxidant; Flavonoids.

## 1. INTRODUCTION

*Piper nigrum* (Black pepper) is one of the most commonly used spices and considered as "The King of Spices" due to its trade in the international market [1, 2]. It is commonly known as Kali Mirch in Urdu and Hindi, Pippali in Sanskrit, Milagu in

Tamil and Peppercorn, White pepper, Green pepper, Black pepper, Madagascar pepper in English [3]. Black pepper is used as medicinal agent, a preservative, and in perfumery [4]. The genus *Piper* has more than 1000 species but the most well known species are *Piper nigrum*, *Piper longum* and *Piper betli* [5]. Black pepper can be used for many different purposes such as human dietaries, as medicine, as preservative, as biocontrol agents [2, 6, 7]. Pepper is used worldwide in different types of sauces and dishes like meat dishes. It contains major pungent alkaloid piperine (1-peperoyl piperidine) which is known to possess many interesting pharmacological actions [8]. Tiwari and Singh [9] reported that this plant and its active components piperine can stimulate the digestive enzymes of pancreas and intestine and also increases biliary bile acid secretion when orally administered. Black pepper is important for its medicinal values [10]. Medicinally black pepper can be used digestive disorder like large intestine toxins, different gastric problems, diarrhoea and indigestion and also can be used against respiratory disorder including cold fever, asthma [11-13]. Piperine exhibits diverse pharmacological activities like antihypertensive and anti-platelets [14], antioxidant, antitumor [15], antipyretic, analgesic, anti-inflammatory, anti-diarrheal, antispasmodic, hepato-protective [16], antibacterial, antifungal, anti-thyroids, anti-apoptotic, anti-spermatogenic, insecticidal and larvicidal activities etc. Piperine has been found to enhance the therapeutic efficacy of many drugs, vaccines and

nutrients by increasing oral bioavailability by inhibiting various metabolising enzymes [17]. In recent pasts, different therapeutic potentials of *Piper nigrum*, its extracts, or its important active chemical constituent "piperine" have been published in different international research journals. The current review is aimed to provide an updated literature review on recent research advancement of pharmacognosy, chemistry and pharmacological activities of *Piper nigrum* L. We have compiled a review on therapeutic potential of *Piper nigrum* by collecting updated scientific research information's from internet using Google search engine.

## 2. TAXONOMICAL CLASSIFICATION OF *PIPER NIGRUM*

Kingdom: Plantae  
 Class: Equisetopsida  
 Sub class: Magnoliidae  
 Super order: Magnolianae  
 Order: Piperales  
 Family: Piperaceae  
 Genus: *Piper*  
 Species: *nigrum*

## 3. GEOGRAPHICAL DISTRIBUTION

Black pepper is grown in many tropical regions like Brazil, Indonesia and India [3]. Geographically, it is confined to Western-Ghats of South India [18]. However, some reports of cultivation from Malaysia, Indonesia, Brazil, Sri-Lanka and West Indies are also available [19]. *P. nigrum* had been found in vast altitudinal regions and showed great adaptability to a wide range of environmental conditions which led to inter-species diversity [20]. "Black-pepper" as its generalized name is due to the color of the peppercorn. It is considered as the "king of spices" due to its trade in the international market [1, 2].

## 4. PHYTOCHEMISTRY


There are many biologically important phytochemicals are extracted from *P. nigrum* plants. They contain alkaloids, amides, propenylphenols, lignans, neolignans, terpenes, steroid, kawapyrones, piperolides, chalcones, dihydrochalcones, brachyamide,

dihydropipericide, 3,4-dihydroxy-6 (N-ethylamine), benzamide, (2E, 4E)-N-eicosadienoyl piperidine, N-trans-feruloyltryamine, N-formyl piperidine, guineensine, (2E, 4E)-N-5[(4-Hydroxyphenyl)-pentadienoyl] piperidine, (2E,4E)-N-isobutyldeca-dienamide), (2E,4E)-N-isobutyl-eicosadienamide, (2E,4E,8Z)-N-isobutyl-eicosatrienamide, (2E,4E)-N-isobutyloctadienamide, piperamide, piperamine, piperettine, pipericide, piperine, piperolein, trichostachine, sarmentine, sarmentosine, tricholein, retrofractamide [21-27] (Figure 1). Pino et al. [27] observed that the major components of the essential oil obtained from the aerial parts of *P. nigrum* were gluulol,  $\alpha$ -pinene,  $\beta$ -caryophyllene and  $\alpha$ -terpinene. Piperine was the first amide to be isolated from piper species and it is the major active principle of black pepper, is closely related in structure to the known natural carcinogens-safrole, estragole and methylengenol which are also widely distributed in spices and plant oils [5]. Zheng et al. [28] studied that the fruits contain 1.0-2.5% volatile oil, 5-9% alkaloids, of which the major ones are piperine, chavicine, piperidine, and piperetine, and a resin. The terpenes, steroids, lignans, flavones, and alkaloids/alkamides have been identified as the primary constituents of the peppers [29]. Khan et al. [30] reported that most of the pharmacological properties of *P. nigrum* fruits are attributed to a piperidine alkaloid, piperine, which is present in the fruits in amounts of 1.7-7.4%. Piperine has also been shown to enhance the bioavailability of several drugs, for example sulfadiazine, tetracycline, streptomycin [31], rifampicin, pyrazinamide, isoniazid, ethambutol and phenytoin [30]. Vasavirama and Upender [5] concluded that due to its diverse pharmacological properties, piperine is important as a biomarker for standardization of fruit of *P. nigrum* and *P. longum* and of polyherbal formulations containing these raw materials.

## 5. ANTI-BACTERIAL ACTIVITY

Karsha and Laxmi [32] reported that anti-bacterial activity of black pepper (*Piper nigrum* Linn.) with special reference to its mode of action on bacteria and found that excellent inhibition on the growth of Gram positive bacteria like *Staphylococcus aureus*, followed by *Bacillus cereus* and *Streptococcus faecalis*. Among the Gram

negative bacteria *Pseudomonas aeruginosa* was more susceptible followed by *Salmonella typhi* and *Escherichia coli*.



**Figure 1.** Structure of important chemical constituent of *Piper nigrum*.

Gram positive bacteria are more susceptible to the extracts due to antibacterial action appears to be loss of control over cell membrane permeability [33]. Khan and Siddiqui, [34] evaluated the antibacterial potential of aqueous decoction of *Piper nigrum* L. (black pepper), *Laurus nobilis* L. (bay leaf), *Pimpinella anisum* L. (aniseed), and *Coriandrum sativum* L. (coriander) against different bacterial isolates from oral cavity of two hundred individual volunteers. Black pepper (aqueous decoction) showed strongest antibacterial activity comparable to aqueous decoction of *Laurus nobilis* and *Pimpinella anisum* at the concentration of 10  $\mu$ l/disc. In a recent study, Palkumar et al. [35]

experimented on *Piper nigrum* leaf and stem assisted green synthesis of silver nano-particles and evaluated its antibacterial activity against agricultural plant pathogens and observe that these silver nano-particles showed the excellent antibacterial activity against plant pathogens. Ganesh et al. [25] experimented photochemical analysis and antibacterial activity of *Piper nigrum* against human pathogenic bacteria and noted that presence of alkaloids, tannins, flavonoids, cardiac and cardiac glycosides shows antibacterial properties against the *Staphylococcus aureus*, *Salmonella typhi*, *Escherichia coli* and *Proteus sp.*

## 6. ANTIOXIDANT ACTIVITY OF BLACK PEPPER

Plants are important source of antioxidants [36]. Antioxidants completely stop or delay the process of oxidation [37]. Some *in vitro* studies revealed that piperine inhibited free radicals and reactive oxygen species, therefore known to possess protective effects against oxidative damage [3]. Free radicals cause many diseases [38]. Different free radicals attack on membranes causing oxidation of lipids, loss of different enzyme activities and may cause cancer [39]. *Piper nigrum* or piperine also found to decrease lipid peroxidation *in vivo* [40]. *Piper nigrum* reported to possess antioxidant activity that might be due to the presence of flavonoids and phenolic contents [41]. *Piper nigrum* was found to prevent the oxidative stress by inhibiting lipid peroxidation, human lipoxygenase and arresting hydroxyl and superoxide free radicals, decrease lung carcinogenesis in animal studies [42]. The memory enhancing and antioxidant properties of the methanolic extract of *Piper nigrum* were investigated in Alzheimer's disease model in rats [43]. The memory-enhancing effects of the extract were studied by means of *in vivo*. While, the antioxidant activity was evaluated by measuring activities of glutathione peroxidase, catalase, superoxide dismutase, and by measuring the total content of reduced glutathione, malondialdehyde, and protein carbonyl levels in the hippocampus [44]. Administration of the methanolic extract of *Piper nigrum* significantly improved memory performance and exhibited antioxidant potential. These studies suggest that methanolic extract of *Piper nigrum*

ameliorates amyloid beta (1-42)-induced spatial memory deterioration by depletion of the oxidative stress in the hippocampus of rats [3]. The antioxidant effect of three Piper species viz *P. nigrum*, *P. guineense* and *P. umbellatum* was evaluated for the protection of renal, cardiac, and hepatic anti oxidant status in atherogenic diet fed hamsters [42]. Piper species significantly inhibited the atherogenic diet induced increased lipid profile and alteration in antioxidant enzymes activities [45]. Ahmad et al. [47] reported that regenerated tissue of *Piper nigrum* like callus, *in vitro* shoots, roots, *in vitro* plantlets, possesses antioxidants activity which is probably due to the presence of flavonoids and phenolic contents. Piper species significantly inhibited the atherogenic diet induced increased lipid profile and alteration in antioxidant enzymes activities [45].

## 7. ANTI-CANCER ACTIVITY OF BLACK PEPPER

*Piper nigrum* had been reported to inhibit tumors formation in different experimental models [47]. Ahmad et al. [23] reported that piperine reduce the lung cancer by altering lipid peroxidation and by antioxidative protection enzymes activation. Piperine has distinct pharmacological activities along with anti-cancer activity [48]. Piperine was reported to inhibit G1/S transition and the proliferation of human umbilical vein endothelial cells (HUVECs), migration of HUVECs and *in vitro* formation of tubule and angiogenesis induced by collagen and breast cancer cell in chick embryos [49]. Landsron et al. [50] reported that piperine inhibits some of the pro-inflammatory cytokines that are produced by tumour cells, there by interfering with the signalling mechanisms between cancer cells, thereby reducing the chances of tumour progression. The anticancer activity of piperine against many cancer cell lines has been reported earlier. Therefore, the mechanisms of anticancer activity of piperine against both androgen independent and dependent cells of prostate cancer were investigated [51]. Piperine treatment was also found to induce apoptosis, by the activation of caspase-3 and by the cleavage of PARP-1 proteins in different prostate cancer cells like PC-3, DU-145 and LNCaP prostate cancer cells [29]. Treatment with piperine

also found to disrupt the androgen receptor expression in LNCaP prostate cancer cells and cause significant diminution in the level of Prostate Specific Antigen in LNCaP cells [52]. The expression of phosphorylated STAT-3 and Nuclear factor- $\kappa$ B transcription factors were reduced in LNCaP, PC-3 and DU-145 prostate cancer cells after treatment of with piperine [28]. Piperine is non-genotoxic and found to possess anti-mutagenic and anti-tumor influences [3]. Dayem et al. [53] reported that *Piper nigrum* reduced lung cancer by modulating lipid peroxidation and through the activation anti oxidative protection enzyme.

## 8. DIGESTIVE ACTIVITY OF BLACK PEPPER

Many spices are known for their digestive stimulant action [54]. Srinivasan, [2] reported that black pepper enhances digestion by stimulation of the pancreatic enzymes and considerably decreases the food transit time of gastrointestinal tract. Ahmad et al. [23] reported that piperine increases the saliva production and gastric secretions, and increases the production and activation of salivary amylase. Platel and Srinivasan, [55] reported that orally administration of piperine or *P. nigrum* stimulate the liver to the secrete bile acids which in turn play key role in the absorption and digestion of fats

## 9. ANTIDEPRESSANT ACTIVITY OF BLACK PEPPER

Ahmad et al. [23] reported the antidepressant activity of piperine and its possible mechanisms was evaluated in corticosterone-induced model of depression in mice. Depression-like behavior in mice was developed after 3 weeks corticosterone injections. The depression was revealed by the significant reduction in sucrose utilization and augmentation in immobility time in the forced swim test and tail suspension test [55]. Further, the brain-derived neurotrophic factor protein and mRNA levels in the hippocampus were also significantly decreased in corticosterone-treated mice. Bai et al. [57] reported that corticosterone induced the behavioral and biochemical changes after treatment to animals with piperine. These results showed that piperine produces an antidepressant-like effect

in corticosterone-induced model of depression in mice [58].

## 10. INSECTICIDAL PROPERTIES

The phytochemical screening of black pepper fruit shows that it contains 4% alkaloids in the berry [13]. Awoyinka et al. [59] reported that the amide olefinic or alkyl isobutylamides compounds such as piperine, piperettine, tricoastacine, piperuloidin, pipartin and trichonine contribute no small measure. These compounds have been demonstrated to be toxic to fruit flies, adzuki bean weevils, cockroaches and several other insect species [59]. Upadhyay and Jaiswal [60] evaluated the biological activities of *Piper nigrum* oil against *Tribolium castaneum* and found that oil had shown a dose response relationship as the larval and adult mortality increased while the larval survival and adult emergence decreased with increase in the concentration of essential oil. Khani et al. [61] reported that the *P. nigrum* extracts offer a unique and beneficial source of bio-pesticide material for the control of insect pests. The toxic effect of *P. nigrum* was reported against some test insects. *P. nigrum* was shown to be most toxic to *Callosobruchus chinensis*, *Acanthoscelides obtectus*, *C. cephalonica*, *Ephestia cautella* Hubn., followed by *Oryzaephilus surinamensis* (L.), *Sitophilus zeamais* Mosteh, *Rhyzopertha dominica* (Fab.) and *Tribolium castaneum* Herbst. The high toxicity effects of *P. nigrum* essential oils against *S. oryzae* adults and 3rd instar larvae of *C. cephalonica* are attributed to the presence of high concentrations of well-known toxic components piperine. Kraikrathok et al. [62] reported that bio efficacy of some piperaceae plant extracts against *Plutella xylostella* third instars under laboratory conditions and observed that the extracts of *Piper nigrum* plants was dose dependant and correlated to duration of exposure. The hexane extract of *P. nigrum* was active with an LD<sub>50</sub> of 18435 ppm and mode of action of these extracts and effect on other developmental parameters was in progress. Scot et al. [63] reported the efficacy of extracts from two Piperaceae species. *Piper nigrum* and *P. tuberculatum* were evaluated using larvae and adults of the colorado potato beetle *Leptinotarsa decemlineata* and noted that young larvae and neonates were the

most susceptible to 0.05% extract of *P. nigrum* reduced larval survival up to 70% within one week after treatment of potato plants. In the greenhouse, *P. nigrum* at 0.5% was as effective at reducing adult *L. decemlineata* feeding as combinations with 2 separate botanical mixtures, garlic and lemon grass oil. Under field conditions, the residual activity of the *P. nigrum* extracts was less than 3 h. When adult *L. decemlineata* were placed on treated plants exposed to full sunlight for 0, 1.5, and 3 h, leaf damage progressively increased as the main active compound, piperine, was found to degrade by 80% after 3 h. The results suggest that Piper extracts could be used effectively as contact botanical insect control agents to protect potato plants from developing *L. decemlineata* larvae at concentrations less than 0.1%. Paula et al. [64] noted that the natural lipophilic amides piperine and piperiline were isolated from *Piper nigrum* evaluated the contact toxicity of all synthetic amides, and also that of piperine and piperiline, at the dose 10 mg per insect, for the Brazilian economically important insects *Ascia monuste orseis* Latr, *Acanthoscelides obtectus* Say, *Brevicoryne brassicae* L, *Protopolybia exigua* DeSaus and *Cornitermes cumulans* Kollar. The results demonstrated that the insects have different sensitivities to the various amides, with mortality ranging from 0 to 97.5%, according to the compound and insect species. Samuel et al. [65] reported that the larvicidal effects of black pepper (*Piper nigrum* L.) and piperine against insecticide resistant and susceptible strains of *Anopheles* malaria vector mosquitoes and observed that Black pepper and piperine mixtures caused high mortality in the An. Gambiae complex strains, with black pepper proving significantly more toxic than piperine. It is concluded that black pepper shows potential as a larvicide for the control of certain malaria vector species.

## 11. ANTIPLATELET ACTIVITY

Srivastava et al. [66] reported that the valuable component of different piper species is piperine which is mostly responsible for various activities. Park et al. [67] reported that piperine possesses anti-platelet activity. Ahmad et al. [23] noted that the toxic effect of piperine on aggression of platelet in experimental rabbit induced by

different factors which activate platelet, by collagen and thrombin.

## 12. MOLLUSCICIDAL ACTIVITY

Srivastava et al. [66] reported that the effect of sublethal treatment (40% and 80% of 24h LC<sub>50</sub>) of *Piper nigrum* fruit and *Cinnamomum tamala* leaf/bark and their different organic solvent extract, purified fraction singly and binary combination with synergist PB or MGK-264 (1:5) on level of different biochemical parameters viz. protein, amino acid, nucleic acids and phospholipids and rate of lipid peroxidation in nervous tissue of *L. acuminata*. Treatment of 80% of 24h LC<sub>50</sub> of piperine caused maximum reduction in protein (12.95% of control), total free amino acid (10.33% of control), DNA (12.70% of control) and RNA (9.17% of control) levels in nervous tissue of *L. acuminata*. Maximum reduction (18.64% of control) in phospholipid levels and elevation of rate of lipid peroxidation (273.17% of control) were observed in the nervous tissue of snails treated with 80% of 24h LC<sub>50</sub> of piperine. Treatment of 80% of 24h LC<sub>50</sub> of purified fraction of *Cinnamomum tamala* leaf/bark caused significant reduction in protein (41.98% of control), total free amino acid (30.06% of control), DNA (43.71% of control) and RNA (16.42% of control), phospholipid (40.86% of control) level and increase the rate of lipid peroxidation (272.69% of control) in nervous tissue of *L. acuminata*. Binary combinations (1:5) plant products with PB or MGK-264 caused significant decrease in the different biochemical parameters. It is clear from the results that there is a significant elevation in lipid peroxidation levels with a reduction in phospholipid levels of nervous tissue of *L. acuminata* treated with different preparations of *P. nigrum* and *C. tamala* leaf/bark. Phospholipids are needed for the growth of endoplasmic reticulum or other cellular membranes [68]. It has been reported that all classes of phospholipids decrease markedly following high dose piperine treatment [69]. The enhancement of lipid peroxidation might be due to oxidative degradation of polyunsaturated fatty acids of the biomembrane leading to pathological infestation [70]. Formation of activated oxygen can have extremely detrimental consequence not only for phospholipids but also protein, nucleic acids,

polysaccharides and inhibition of vital enzymes [71, 72]. The alkaloid, piperine, found in *P. nigrum* destroys the cytochrom P-450 and inhibits monoxygenase activity [73]. Some workers reported the effect of piperine activity in rat. Johri et al. [74] studied that the effect of piperine on the absorptive function of the intestine. *In vitro* experiments showed an increased rate of lipid peroxidation in the freshly isolated epithelial cells of rat jejunum. These results suggested that piperine may interact with the lipid environment to produce effect which leads to increased permeability of the intestinal cells.

## 13. ANTIREPRODUCTIVE ACTIVITY

Srivastava et al. [75] reported that the anti-reproductive activity of piperine against the snail *Lymnaea acuminata* and observed that piperine caused a significant reduction in the fecundity, hatchability and survival of the snail *Lymnaea acuminata* in each month of the year Nov. 2011 to Oct. 2012. Treatment with the piperine also prolong the hatching time of snails. Sublethal treatment of piperine caused a significant ( $p < 0.05$ ) reduction in protein, amino acids, DNA, RNA and AChE in the ovotestis/nervous tissue of treated snails with respect to control after 96h exposure period. Simultaneously, inhibition in acetylcholinesterase (AChE) activity in nervous tissue was also noted. The active component piperine (*Piper nigrum*) is an effective molluscicide against *L. acuminata*. Constituent of piperine *in vitro* inhibit enzyme activity which is responsible for leukotriene and prostaglandin biosynthesis; 5-lipoxygenase and COX-1 [76]. The cerebral neurosecretory caudo dorsal cells (CDCS) of the fresh water pulmonate snail *Lymnaea stagnalis* control egg lying, an event that involves a pattern of stereotyped behavior [78]. The CDCS synthesize and release multiple peptides, among which is the ovulation hormone (CDCS). It is thought that each peptide controls a specific aspect of the processes involved in egg laying [77].

The synthesis of protein in any of a tissue can be affected in two ways by a chemical, (I) it either affects the RNA synthesis at the transcription stage or (II) it somehow affects the uptake of amino acids in the polypeptide chain. Both these possibilities may account for the lower protein content in the

affected tissue. In the first case, the RNA synthesis would be inhibited resulting in reduced RNA as well protein content. In the second case, only the protein content would be affected [66, 78, 79]. Piperine inhibits P-glycoprotein and the major drug metabolizing enzyme CYP3A4 [81]. It seems that cumulative effect of molluscicide piperine on the level of protein, amino acids and nucleic acids in ovotestis of *L. acuminata* directly/or indirectly CDCs, which release ovulation hormone and ultimately affect the reproduction of snails in each month of the year. The AChE activity is one of the biomarker most frequently used in ecotoxicology. The enzyme is responsible for the breakdown of ACh in cholinergic synapses, preventing continuous nerve firing, which is vital for normal cellular neurotransmitter functioning [81]. The AChE inhibition result in accumulation of acetylcholinesterase at the nerve synapses so that the post synaptic membrane is in a state of permanent stimulation producing paralysis, ataxia and general lack of coordination in neuromuscular system and eventual death [82, 83].

#### 14. COSMOPERINE ACTIVITY

Sabina Corporation [84] reported that cosmoperine prepared from piperine used in cosmetics, a natural bio-enhancer which improve the permeability of active compounds through skin. Cosmoperine activate and stimulate the natural power of skin to absorb nutrients [85, 86]. Cosmoperine isolated from piperine are non-irritant, interacts with the skin quantitatively and qualitatively in various means, furthermore, cosmoperine are pain relieving and causes skin reddening due to vascular engorgement as well as a slight skin tingling sensation.

#### 15. CONCLUSION

In the present review we have made an attempt to congregate the botanical, phytochemical, toxicological information on *Piper nigrum* a medicinal herbs used in the Indian system of medicine, Survey of literature revealed that the presence of alkaloids, lignans, volatile oils and esters in different parts of this plants. Research on alkaloids has gained a special attention in recent

times as several of them have shown promising activities like anti-inflammatory, hepatoprotective, stimulant effect, anti-amoebic and antibacterial etc. This review definitely helps for the researchers as well as practioners, dealing with this plant, to know its proper usage.

#### AUTHORS' CONTRIBUTION

Both authors contributed equally for the success of this review article. The final manuscript has been read and approved by both authors.

#### TRANSPARENCY DECLARATION

The authors declare that there is no conflict of interest regarding the publication of this article.

#### REFERENCES

1. Mathew PJ, Mathew PM, Kumar V. Graph clustering of *Piper nigrum* L. (black pepper). *Euphytica*. 2001; 18: 257-264.
2. Srinivasan. Black pepper and its pungent principle-piperine: a review of diverse physiological effects. *Crit Rev Food Sci Nutr*. 2007; 47(8): 735-748.
3. Damanhoury ZA, Ahmad A. A review on therapeutic potential of *Piper nigrum* L. (black pepper): the king of spices. *Med Aromat Plants*. 2014; 3(3): 161.
4. Singh VK, Singh P, Mishra A, Patel A, Yadav KM. Piperine: delightful surprise to the biological world, made by plant "pepper" and a great bioavailability enhancer for our drugs and great bioavailability enhancer for our drugs and supplements. *World J Pharmac Res*. 2014; 3(6): 2084-2098.
5. Vasavirama K Upender M. Piperine: a valuable alkaloid from piper species. *Int J Pharm Pharm Sci*. 2014; 6(4): 34-38.
6. Awen BZ, Ganapati S, Chandu BR. Influence of *Sapindus mukorossi* on the permeability of ethyl cellulose free film for transe dermal use. *Res J Pharma Biol Chem Sci*. 2010; 1: 35-38.
7. Hussain A, Naz S, Nazir H, Shinwari ZK. Tissue culture of black pepper (*Piper nigrum* L.) in Pakistan. *Pak J Bot*. 2011; 43: 1069-1078.
8. Ahmad N, Fazal H, Abbasi BH, Farooq S, Ali M, Khan MA. Biological role of *Piper nigrum* L. (black pepper): a review. *Asian Pac J Trop Biomed*. 2015: S1945-S1953.

9. Tiwari P, Singh D. Anti-trichomonas activity of *Sapindus saponins*, a candidate for development as microbicidal contraceptive. J Antimicrob Chemother. 2008; 62: 526-534.
10. Dhanya K, Kizhakkayil J, Syamkumar S, Sasikumar B. Isolation and amplification of genomic DNA from recalcitrant dried berries of black pepper (*Piper nigrum* L.). A medicinal spice. Mol Biotechnol. 2007; 7: 165-168.
11. Sujatha R, Luckin CB, Nazeem PA. Histology of organogenesis from callus cultures of black pepper (*Piper nigrum* L.). J Trop Agric. 2003; 41: 16-19.
12. Parganiha R, Verma S, Chandrakar S, Pal S, Sawarkar HA, Kashyap P. *In vitro* anti-asthmatic activity of fruit extract of *Piper nigrum* (Piperaceae). Int J Herbal Drug Res. 2011; 1: 15-18.
13. Fan LS, Muhmad R, Omar D, Rahimani M. Insecticidal properties of *Piper nigrum* fruit extracts and essential oils against *Spodoptera litura*. Int J Agric Biol. 2011; 13: 517-522.
14. Taqvi SI, Shah AJ, Gilani AH. Blood pressure lowering and effects of piperine. J Cardiovasc Pharmacol. 2008; 52: 452-458.
15. Manoharan S, Balakrishnan S, Menon VP, Alias LM, Reena AR. Chemopreventive efficacy of curcumin and piperine during 7,12-dimethylbenz[a]anthracene-induced hamster buccal pouch carcinogenesis. Singapore Med J. 2009; 50: 139-146.
16. Matsuda H, Ninomiya K, Morikawa T, Yasuda D, Yamaguchi I, Yoshikawa M. Protective effects of amide constituents from the fruit of *Piper chaba* on D- galactosamine/TNF-alpha induced cell death in mouse hepatocytes. Bioorg Med Chem Lett. 2008; 18: 2038-2042.
17. Chitlange SS, Payal BS, Sanjay D, Nipanikar, Dheeraj N. Development and validation of RP-HPLC method for quantification of piperine from single herb formulation containing *Piper nigrum* extract. Int J Pharm Pharmacol Sci Res. 2016; 6(2): 16-21.
18. Nair RR, Gupta SD. Somatic embryogenesis and plant regeneration in black pepper (*Piper nigrum* L.): I. Direct somatic embryogenesis from tissues of germinating seeds and ontogeny of somatic embryos. J Hort Sci Biotech. 2003; 78: 416-421.
19. Gupta V, Meena AK, Krishna CM, Rao, MM, Sannd R, Singh H, et al. Review of plants used as kshar of family piperaceae. Int J Ayurveda Med. 2010; 1(2): 2010.
20. Howard RA. Notes on the Piperaceae of lesser antilles. J Arnold Arb. 1973; 54: 377-411.
21. Abbasi BH, Ahmad N, Fazal H, Mahmood T. Conventional and modern propagation techniques in *Piper nigrum*. J Med Plants Res. 2010; 4(1): 7-12.
22. Khusbu C, Roshni S, Anar P, Corol M, Mayuree P. Phytochemical and therapeutic potential of *Piper longum* Linn. a review. Int J Res Ayurveda Pharma. 2011; 2(1): 157-161.
23. Ahmad N, Fazal H, Abbasi BH, Farooq S, Ali M, et al. Biological role of *Piper nigrum* L. (Black pepper): a review. Asian Pac J Trop Biomed. 2012: S1945-S1953.
24. Kumar MA, Sinha A, Verma SC, Gupta MD, Padhi MM. HPTLC Profile of important Indian spices used in ayurvedic formulations. Res J Pharmacogn Phytochem. 2013; 5(4): 188-193.
25. Ganesh P, Kumar RS, Saranraj P. Phytochemical analysis and antibacterial activity of pepper (*Piper nigrum* L.) against some human pathogens. Central Eur J Exp Biol. 2014; 3(2): 36-41.
26. Pelayo VRT, Fernandez MS, Hernandez OC, Torres JM, Garcia JAL. A phytochemical and ethnopharmacological review of the genus *Piper*: as a potent bio-insecticide. Res J Biol. 2016; 4(2): 45-51.
27. Pino JA, Aguero J, Fuentes V. Chemical composition of the aerial parts of *Piper nigrum* L. from Cuba. J Essent Oil Res. 2003; 15: 209-210.
28. Zheng J, Zhou Y, Li Y, Xu DP, Li S, Li HB. Spices for prevention and treatment of cancers. Nutrients. 2016; 8: 495.
29. Rajeswari R. Phytochemical analysis of *Guettarda speciosa* Linn. Asian J Plant Sci Res. 2015; 5(9): 1-4.
30. Khan S, Mirza KJ, Anwar F, Abdin MJ. Development of RAPD markers for authentication of *Piper nigrum* (L.). Environ Int J Sci Tech. 2010; 5: 47-56.
31. Verma VC, Lobkovsky E, Gange AC, Singh SK, Prakash S. Piperine production by endophytic fungus *Periconia* sp. isolated from *Piper longum* L. J Antibiotics. 2011; 64: 427-431.
32. Karsha PV, Laxmi OB. Antibacterial activity of black pepper with special reference to its mode of action on bacteria. Ind J Nat Prod Resour. 2010; 1(2): 2013-215.
33. O'Bryan CA, Pendleton SJ, Philip G. Crandall, Ricke SC. Potential of plant essential oils and their components in animal agriculture - in vitro studies



- on antibacterial mode of action. *Front Vet Sci.* 2015; 2: 35.
34. Khan M, Siddiqui M. Antimicrobial activity of fruits of *Piper longum*. *Nat Prod Rad.* 2007; 6: 111 - 113.
  35. Paulkumar K, Gnanajobitha G, Vanaja M, Rajeshkumar S, Malarkodi C, Pandian K, Annadurai G. *Piper nigrum* leaf and stem assisted green synthesis of silver nanoparticles and evaluation of its antibacterial activity against agricultural plant pathogens. *Scient World J.* 2014: 829894.
  36. Kasote DM, Katyare SS, Hegde MV, Bae H. Significance of antioxidant potential of plants and its relevance to therapeutic applications. *Int J Biol Sci.* 2015; 11(8): 982-991.
  37. Young IS, Woodside JV. Antioxidants in health and disease. *J Clin Pathol.* 2001; 54: 176-186.
  38. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010; 4(8): 118-126.
  39. Marakala. Oxidant - antioxidant imbalance in cancer: a review. *J Evol Res Med Biochem.* 2015; 1(1): 18-23.
  40. Srivastava AK. Effect of certain attractant bait formulations, containing plant molluscicides on the reproduction of *Lymnaea acuminata* with reference to seasonal variation in abiotic factors. Ph.D. Thesis, DDU Gorakhpur University, Gorakhpur, India, 2013.
  41. Shanmugapriya K, Saravana PS, Payal H, Mohammed SP, Williams B. Antioxidant potential of pepper (*Piper nigrum* Linn.) leaves and its antimicrobial potential against some pathogenic microbes. *Ind J Nat Prod Resour.* 2012; 3(4): 570-577.
  42. Chelak SK, Saraf S, Saraf S. Preformulation and formulation study of anticancer principle of piperine. *World J Pharma Res.* 2015; 4(12): 722-737.
  43. Mahady K, Shaker O, Wafay H, Nassar Y, Hassan H, Hussein A. Effect of some medicinal plant extracts on the oxidative stress status in Alzheimer's disease induced in rats. *Eur Rev Med Pharmacol Sci.* 2012; 16(3): 31-42.
  44. Krishna RK, Krishnakumar S, Chandrakala S. Evaluation of antioxidant properties of different parts of *Amorphophallus commutatus*, an endemic aroid of western ghats, south India. *Int J Pharm Bio Sci.* 2012; 3(3): 443-455.
  45. Aqbor GA, Akinfiresoye L, Sortino J, Johnson R, Vinson JA. Piper species protect cardiac, hepatic and renal antioxidant status of atherogenic diet fed hamsters. *Food Chem.* 2012; 34(3): 1354-1359.
  46. Ahmad N, Fazal H, Abbasi BH, Rashid M, Mahmood T, Fatima N. Efficient regeneration and antioxidant potential in regenerated-tissues of *Piper nigrum* L. *Plant Cell Tissue Organ Cult.* 2010; 102: 129-134.
  47. Gaziano R, Moroni G, Buè C, Tony Miele M, Vallebona PS, Pica F. Antitumor effects of the benzophenanthridine alkaloid sanguinarine: evidence and perspectives. *World J Gastroint Oncol.* 2016; 8(1): 30-39.
  48. Paarakh PM, Sreeram DC, Shruthi SD, Ganapathy SPS. In vitro cytotoxic and in silico activity of piperine isolated from *Piper nigrum* fruits Linn. In *Silico Pharmacol.* 2015; 3: 9.
  49. Wang N, Wang JY, Mo SL, Loo TY, Wang DM, Luo HB, et al. Ellagic acid, a phenolic compound, exerts anti-angiogenesis effects via VEGFR-2 signaling pathway in breast cancer. *Breast Cancer Res Treat.* 2012; 134(3): 943-955.
  50. Landskron G, Fuente MD, Thuwajit P, Thuwajit C, Hermoso MA. Chronic inflammation and cytokines in the tumor microenvironment. *J Immunol Res.* 2014: 149185.
  51. Reddy MN, Reddy NR, Jamil K. Spicy anti-cancer spices: A review. *Int J Pharm Pharm Sci.* 2015; 7(11): 1-6.
  52. Zhaomei M, Hachem P, Hensley H, Stoyanova R, Kwon HW, Hanlon AL, et al. Antisense MDM2 Enhances the response of androgen insensitive human prostate cancer cells to androgen deprivation in vitro and in vivo. *Prostate.* 2008; 68(6): 599-609.
  53. Dayem AA, Choi HY, Yang GM, Kim K, Saha SK, Cho SG. The anti-cancer effect of polyphenols against breast cancer and cancer stem cells: molecular mechanisms. *Nutrients.* 2016; 8(9): 581.
  54. Adefegha SA, Oboh G. Phytochemistry and mode of action of some tropical spices in the management of type-2 diabetes and hypertension. *Afr J Pharm Pharmacol.* 2013; 7(7): 332-346.
  55. Platel K, Srinivasan K. Digestive stimulant action of spices: a myth or reality? *Ind J Med Res.* 2004; 119: 167-179.
  56. Singh P, Kumar P, Singh VK, Singh DK. Effect of snail attractant pellets containing plant molluscicides on certain enzyme in the nervous tissue of *Lymnaea acuminata* (Lamarck). *The Bioscan.* 2009; 4(3): 395-398.

57. Bai X, Zhang W, Chen W, Zong W, Guo Z, Liu X. Antihepatotoxic and antioxidant effects of extracts from *Piper nigrum* L. root. *Afr J Biotechnol.* 2011; 10: 267-272.
58. Mao QQ, Huang Z, Siu-P, Xian YF, Chun-Tao C. Protective effects of piperine against corticosterone-induced neurotoxicity in PC12 cells. *Cell Mol Neurobiol.* 2012; 32(4): 531-537.
59. Awoyinka OA, Oyewole IO, Amos BM, Onasoga OF. Comparative pesticidal activity of dichloromethane extracts of *Piper nigrum* against *Sitophilus zeamais* and *Callosobruchus maculatus*. *Afr J Biotech.* 2006; 5: 2446-2449.
60. Upadhyay RK, Jaiswal G. Evaluation of biological activities of *Piper nigrum* oil against *Tribolium castaneum*. *Bull Insectol.* 2007; 60(1): 57-61.
61. Khani M, Awang RM, Omar D. Insecticidal effects of peppermint and black pepper essential oils against rice weevil, *Sitophilus oryzae* L. and rice moth, *Corcyra cephalonica* (St.). *J Med Plants.* 2012; 11(43): 97-110.
62. Kraikrathok C, Ngamsaeng S, Bullangpoti V, Pluempanupat W, Koul O. Bio efficacy of some piperaceae plant extracts against *Plutella xylostella* l. (Lepidoptera: Plutellidae). *Comm Appl Biol Sci.* 2013; 78(2): 305-310.
63. Scott IM, Jensen H, Scott JG, Isman MB, Arnason JT, Philogène BJR. Botanical insecticides for controlling agricultural pests: Piperamides and the colorado potato beetle *Leptinotarsa decemlineata* say (Coleoptera: Chrysomelidae). *Arch Insect Biochem Physiol.* 2003; 54: 212-225.
64. Paula VF, Barbosa LCA, Demuner AJ, Pilo-Veloso D, Picanc MC. Synthesis and insecticidal activity of new amide derivatives of piperine. *Pest Manag Sci.* 2000; 56: 168-174.
65. Samuel M, Oliver SV, Coetzee M, Brooke BD. The larvicidal effects of black pepper (*Piper nigrum* L.) and piperine against insecticide resistant and susceptible strains of *Anopheles* malaria vector mosquitoes. *Parasites Vectors.* 2016; 9: 238.
66. Srivastava P, Kumar P, Singh VK, Singh DK. Effect of *Piper nigrum* and *Cinnamomum tamala* on biochemical changes in the nervous tissue of fresh water snail *Lymnaea acuminata*. *Bioscan.* 2010; 1: 247-256.
67. Park BS, Son DJ, Park YH, Kim TW, Lee SE. Antiplatelet effects of acid amide isolated from the fruits of *Piper longum* L. *Phytomed.* 2007; 14: 853-855.
68. Moore TS. Phospholipid biosynthesis. *Ann Rev Plant Physiol.* 1982; 33: 235-259.
69. Malini T, Arunakaran J, Aruldas MM, Govindarajulu P. Effects of piperine on the lipid composition and enzymes of the pyruvate -malate cycle in the testis of the rat *in vivo*. *Biochem Mol Biol Int.* 1999; 47(3): 537-545.
70. Dillard CJ, Tappel AL. Fluorescent products of mitochondria and microsomes. *Lipids.* 1971; 6: 715-721.
71. Inlay JA, Linn S. DNA damage and oxygen radical toxicity. *Sci.* 1988; 240: 1302.
72. Singh DK, Singh VK, Kumar P. Pestiferous gastropods and their control. LAP Lambert. Academic Publication GmbH and Co. Germany. 2012.
73. Atal CK, Dubey RK, Singh J. Biochemical basis of enhanced drug bioavailability by piperine: evidence that piperine is a potent inhibitor of drug metabolism. *J Pharmacol Exp Ther.* 1985; 232: 258-262.
74. Johri RK, Tusu N, Khajuria A, Zutshi U. Piperine-mediated changes in the permeability of rat intestinal epithelial cells. The status of gamma glutamyl transpeptidase activity, uptake of amino acids and lipid peroxidation. *Biochem Pharmacol.* 1992; 43: 1401-1407.
75. Srivastava AK, Singh DK, Singh VK. Change of seasonal variation and feeding of bait containing piperine on reproduction and certain biochemical changes of fresh water snail *Lymnaea acuminata*. *Front Biol Life Sci.* 2014; 2(3): 53-61.
76. Stohr JR, Xiao PG, Bauer R. Constituents of Chinese piper species and their inhibitory activity on prostaglandin and leukotriene biosynthesis *in vitro*. *J Ethnopharmacol.* 2001; 75: 133-139.
77. Vreugdenhil E, Jackson JF, Bouwmeester T, Smit AB, Van Minnen J, Vanheerikhuizen H, et al. Isolation, characterization and evolutionary aspects of a cDNA clone encoding multiple neuropeptides involved in a stereotyped egg-laying behavior of the fresh water snail *Lymnaea stagnalis*. *J Neurosci.* 1988; 81: 4184-4191.
78. Tariq S, Haqqi M, Usman MA. Effect of thiotepa on RNA and total protein synthesis content in testis of albino rats. *Ind J Exp Biol.* 1977; 15: 804-805.
79. Singh RN, Kumar P, Singh VK, Singh DK. Toxic effects of deltamethrin on the levels of biochemical changes in the snail *Lymnaea acuminata*. *J Pharm Res.* 2010; 3(8): 1739-1742.

80. Bhardwaj RK, Glaeser H, Becquemont L, Koltz U, Guptan SK, Fromm MF. Piperine a major constituent of black pepper, inhibits human P-glycoprotein and CYP3 A4. *Pharmacol Exp Ther.* 2002; 302: 645-650.
81. Elena F, Antonio C. Cyclic AMP signaling in bivalve mollusks: an overview. *J Exp Zool.* 2010; 313: 179-200.
82. Matsumura F. *Toxicology of insecticides.* Plenum press, New York, 2nd ed. 1985.
83. Singh P, Kumar P, Singh VK, Singh DK. Effect of snail attractant pellets containing plant molluscicides on certain enzyme in the nervous tissue of *Lymnaea acuminata* (Lamarck). *Bioscan.* 2009; 4(3): 395-398.
84. Sabinsa Corporation. Sabinsa corporation home page. East Windsor, NJ: Sabinsa Corporation. 2011; Available from: [www.sabinsa.com](http://www.sabinsa.com). [Accessed on 2011 Apr 20].
85. Badmaev V, Majeed M, Norkus EP. Piperine, an alkaloid derived from black pepper, increases serum response of beta-carotene during 14 Days of oral beta-carotene supplementation. *Nutr Res.* 1999; 19: 381-388.
86. Majeed M, Prakash LTHP. An all natural delivery system adjuvant. In *delivery system handbook for personal care and cosmetic products: Technology, applications and formulations.* Meyer RR, ed, William and Andrew Publishing, 2005.