

# Risk Factors of Subependymal Hemorrhage-Intraventricular Haemorrhage in Preterm Infants

Gulbin Shahid <sup>1</sup>, Mahmood Jamal <sup>2</sup>, Yasir Bin Nisar <sup>3</sup>

1. Department of Paediatric Medicine, Children's Hospital, Pakistan Institute of Medical Sciences, Islamabad ;2. Department of Paediatrics, Islamabad Medical and Dental College, Islamabad; 3. Public Health Division , WHO, Geneva.

## Abstract

**Background:** To determine the frequency and peak time of occurrence of subependymal hemorrhage-intraventricular hemorrhage (SEH-IVH) in preterm infants and to identify the predisposing risk factors.

**Methods:** In this descriptive study consecutive live born babies (n=135), between the gestational ages of 28-34 weeks, were enrolled. Information regarding weight, gestational age, mode of delivery and presence of risk factors were recorded. Each newborn underwent cranial ultrasound on designated days and was followed up till the fourth week of life unless expired.

**Results:** The frequency of SEH-IVH was found to be 20.8%. The most vulnerable group in terms of gestational age and weight were infants < 32 wks and weighing < 1.5 kg respectively. The peak age of occurrence of SEH-IVH was first three days of life. Risk factors found to be associated with SEH-IVH were RDS with ventilation, exchange transfusion and thrombocytopenia.

**Conclusion:** The incidence of SEH-IVH in infants between 28-34 wks gestation is significantly high. The crucial period is the first three days of life and respiratory distress syndrome (RDS) with ventilation, exchange transfusion and thrombocytopenia carry the highest risk of hemorrhage. There is a need to emphasize on the prevention of prematurity and optimum perinatal management to minimize the risk of hemorrhage.

**Key words:** Subependymal hemorrhage-intraventricular hemorrhage, Premature infants.

## Introduction

Subependymal haemorrhage - intraventricular hemorrhage is one of several complications associated with prematurity and carries significant morbidity and mortality. It is principally a problem of very low birth weight infants i.e. infants weighing less than 1.5kg and also those delivered before 32-34wks of gestation.<sup>1-4</sup> With improved obstetric and perinatal management, the incidence of SEH-IVH has shown a significant decline. <sup>5-10</sup> Tertiary centres currently report an

incidence between 20-25% in infants born before 34 wks of gestation with figures rising with decreasing weight and gestational age.<sup>2,4,11-25</sup> The incidence of IVH in premature infants between 500g-749g is still very high and is approximately 45% .<sup>26</sup>

Upto 90% of IVH cases evolve in the first 72 hours of postnatal life; 30% in the first 6 hours, 40-60% in the first 24 hours and 60-80% within 48 hours<sup>27-30</sup> Hemorrhage on or after the fourth day of life occurs in 10% of all preterm infants with IVH.<sup>1</sup> Intraventricular hemorrhage has been extensively documented in preterm newborns with respiratory distress syndrome requiring mechanical ventilation.<sup>10,18,31-33</sup> Other well recognized risk factors for IVH include asphyxia, hypercarbia, rapid volume expansion, routine care taking maneuvers, patent ductus arteriosus, high concentration of inspired oxygen, pneumothorax, seizures, anemia, low blood glucose, platelet and coagulation defects.<sup>1,18,31-42</sup> Better understanding of the pathogenesis of IVH has led to several therapeutic interventions aimed at counteracting the known risk factors. Many such interventions like use of antenatal steroids and careful ventilator management have demonstrated a significant reduction in the incidence of IVH.<sup>43-49</sup> In developing countries incidence has been reported to be between 28%-75%.<sup>50-54</sup> More recent data also shows a very wide range between 9%-64.4%.<sup>27, 55-62</sup>

## Patients and Methods

This non-interventional, descriptive, analytic study was carried out in the Department of Neonatology, Children's Hospital, Pakistan Institute of Medical Sciences (P.I.M.S), Islamabad from September 2002 to January 2003. During the study period, all consecutive live births delivered at the MCH Centre with gestational ages between 28-34 weeks were enrolled. Weight was measured without clothes and grouped under four categories, namely 0.75kg-1kg, 1.1kg-1.5kg, 1.6kg-2.0kg, and >2.0kg. Gestational age was estimated by dates i.e. last menstrual period or antenatal ultrasound if performed. In the absence of aforementioned information, Dubowitz scoring was used for estimating the gestational age.<sup>63</sup> Groups assigned for gestational age were 28 wks, 29-30 wks,

31-32 wks and 33-34 wks. Babies with gestational age <28 weeks, with weight <750gms and with lethal malformations were excluded from this study. Cranial ultrasonography of selected patients was performed. Images were obtained through the anterior fontanelle in both coronal and right and left parasagittal planes. Every enrolled infant was followed up for 28 days or till death if earlier. Thirteen babies, without IVH on the first two visits, failed to report for 3<sup>rd</sup> cranial ultrasonography on the designated days. They were reported as well on telephone by parents. These infants underwent cranial ultrasound examination at least three times, (0-3, 7-10, 21-28 days). Admitted babies, however, had daily ultrasounds till they were discharged or expired. In addition to this, ultrasound was also done whenever clinical indications like abrupt fall in hematocrit, shock, bulging fontanel, i.e., change in the level of consciousness or change in the respiratory support without any other explanation, appeared at any time during admission. IVH was classified into four grades (Gd. I-IV) according to the system of grading reported by Papile; Gd I-isolated SEH, Gd II-rupture into ventricle but no ventricular dilatation, Gd III-rupture into ventricle with ventricular dilatation and Gd IV-IVH with parenchymal extension.<sup>64</sup>

Risk factors were defined according to standard recommendations and protocols used in the neonatal unit. Sample size was calculated by using statcalc of software EpiInfo version 6.0 by using descriptive study, random, non-cluster sampling technique. By keeping the population size as 400 (estimated annual deliveries in MCH at the time with gestation of 28-34 weeks) and expected frequencies as 25% and worst expected being 20%, with confidence level of 95%, a sample size of 135 was calculated. Univariate and multivariate analysis was carried out for risk factors to determine their effect on SEH-IVH and Odds ratio with 95% confidence interval were calculated by keeping p-value of < 0.05 as significant.

### Results

Of these 135 babies with gestational age of 28-34 weeks, 60% were male and 40% were females. The mean gestational age was 31.4±2.05 weeks. The mean birth weight of study population was 1.59±0.42 kg. Two third of babies were delivered by SVD whereas about 1/3<sup>rd</sup> (30%) were delivered through C-Section (Table 1). Of the total 135 babies, 28 babies were found to have SEH-IVH on ultrasonography during their neonatal period. Eight babies each had Grade I and Grade II bleed while 5 babies had Grade III and seven had grade IV bleed. Three fourth (21) of

the babies developed SEH-IVH during the 1<sup>st</sup> three days of life. Of these, again, the majority i.e. 12 out of 21 babies had SEH-IVH within 1<sup>st</sup> 24 hours of life. Almost all SEH-IVH occurred within the 1<sup>st</sup> week of life with the exception of one, where the bleeding occurred on 14<sup>th</sup> day of life (Table 2; Figure 1). Two of the studied risk factors i.e. birth asphyxia and hypoglycemia were not found to be associated with SEH-IVH, while all others including RDS, mechanical ventilation, hypoxia, PDA, high FiO<sub>2</sub>, exchange transfusion, anemia and thrombocytopenia were found to be significantly associated with SEH-IVH (Table 3). Thrombocytopenia carried the highest odds ratio, however, RDS with ventilation, exchange transfusion and thrombocytopenia retained their significance independently in multivariate logistic regression (stepwise) model (Table 4).

**Table 1: Comparison of demographic factors of premature infants with and without SEH-IVH (n = 135)**

Variables	SEH-IVH (n = 28)	NO SEH-IVH (n = 107)	OR (95%CI), p-value
Sex			
Male	19 (67.9%)	63 (58.1%)	1.47 (0.57-3.91) 0.51
Female	9 (32.1%)	44 (41.1%)	
Gestational Age			
Median	31.0	32.0	1.57 (0.21-9.95) 0.96
Mean ± SD	30.0 ± 2.0	31.6 ± 2.0	
28 wks	2 (2.1%)	5 (4.7%)	1.92 (0.73-5.00) 0.21 1.34 (0.49-3.58) 0.69 0.35 (0.12-1.00) 0.05
29 to 30 wks	11 (39.3%)	27 (25.2%)	
31 to 32 wks	9 (32.1%)	28 (26.2%)	
33 to 34 wks	6 (21.5%)	47 (43.9%)	
Weight (kg)			
Median	1.26	1.64	0.46 (0.02-3.91) 0.75
Mean ± SD	1.45 ± 0.40	1.63 ± 0.40	
0.75 to 1.0 kg	1 (3.6%)	8 (7.5%)	2.39 (0.95-6.12) 0.06 0.39 (0.12-1.22) 0.11
1.1 to 1.5 kg	17 (60.6%)	42 (39.3%)	
1.6 to 2.0 kg	5 (17.9%)	38 (35.5%)	
More than 2.0 Kg	5 (17.9%)	19 (17.7%)	1.01 (0.29-3.28) 0.79
Mode of delivery			
SVD	18 (64.3%)	73 (68.2%)	0.84 (.32-2.20) 0.86
C-Section	9 (32.1%)	32 (29.9%)	1.11 (0.41-2.94) 0.99
Forceps	1 (3.6%)	2 (1.9%)	1.94 (0.00-28.9) 0.86




Fig 1: Relationship of age with SEH-IVH

Of these, the babies on assisted ventilation with RDS carried the highest risk of SEH-IVH closely followed by babies undergoing exchange transfusion and thrombocytopenia. Other risk factors like hypoxia, high concentration of O<sub>2</sub>, PDA, and anemia lost their significance in multivariate logistic regression analysis. Out of these 28 infants, 20(71.4%) expired and only 8(28.3%) were either discharged or remained admitted beyond 28 days of life. Of this mortality group, majority (12) had grade III and Grade IV bleed while only three babies with Grade I SEH-IVH expired during their stay in the hospital.

Table 2: Frequency of SEH-IVH on days of life (n = 28)

Age in days	Number	Percentage
First day	12	42.8%
Second day	5	17.8%
Third day	4	14.3%
Fourth day	1	3.6%
Fifth day	4	14.3%
Sixth day	1	3.6%
Fourteenth day	1	3.6%

### Discussion

The most common form of brain injury in the premature infant is subependymal hemorrhage-intraventricular hemorrhage. Most tertiary care centres from the United States, Australia and other developed countries currently report an incidence between 20-25% in infants born before 34 wks of gestation.<sup>2, 4, 11-18</sup> Data from the developing countries shows a wide range of incidence of IVH in premature infants. Generally the incidence is very high primarily because of lack of optimum obstetric and perinatal care.<sup>53</sup>

Table 3: Predisposing factors associated with SEH-IVH in premature infants (n = 135)

Variable	SEH-IVH (n = 28)	NO SEH-IVH (n = 107)	X <sup>2</sup>	DF	OR (95% CI), p-value
Birth asphyxia	5 (17.9%)	6 (5.6%)	2.96	1	3.66(0.87-15.2), 0.08
RDS	12(42.9%)	19 (17.8%)	6.55	1	3.47(1.29-9.37) 0.01
Mechanical ventilation	18(64.3%)	15 (14.0%)	27.7	1	11.04(3.91-32.0) <0.00001
Hypoxia	14(50.0%)	18 (16.8%)	11.74	1	4.94(1.84-13.39) <0.00001
PDA	3 (10.7%)	1 (0.9%)	4.37	1	12.72(1.10-331.79) 0.03
High concentration of inspired O <sub>2</sub>	12(42.9%)	12 (11.2%)	13.11	1	5.94(2.06-17.34) <0.00001
Exchange transfusion	15(53.6%)	21 (19.6%)	11.40	1	4.73(1.79-12.57) <0.00001
Anemia	6 (21.4%)	6 (5.6%)	5.04	1	4.59(1.17-18.18) 0.02
Hypoglycemia	6(21.4%)	8 (7.5%)	3.27	1	3.38(0.92-12.23) 0.07
Thrombocytopenia	11(39.3%)	3 (2.8%)	27.97	1	22.43(5.93-114.5) <0.00001

Table 4: Risk Factors for SEH-IVH in premature infants by using multivariate logistic regression

Risk Factors	OR (95% CI)	p-Value
RDS with Ventilation	8.21 (2.30 - 28.33)	< 0.00001
Exchange transfusion	7.79 (2.00 - 29.09)	0.003
Thrombocytopenia	5.09 (1.70 - 14.98)	0.002

The most vulnerable group in present study were infants between 28-32 weeks gestation. This inverse relationship between gestational age and SEH-IVH is expected and in accordance with other studies. Dolfin et al reported a significant difference between the incidence of SEH-IVH in infants whose gestational age was less than or equal to 29 weeks compared to infants between 30-34 weeks gestation. Among infants less

than or equal to 29 weeks gestational age 56% had SEH-IVH while only 8% had bleed in the gestational group of 32-34 weeks.<sup>66</sup>

It was seen in the present study that babies between 1.6-2.0 kg had an incidence of IVH of only 12% increasing to 30% in infants between 1.1-1.5 kg. This increasing trend in the frequency with decreasing weight was expected and in accordance with other studies. However, the prevalence of SEH-IVH for infants between 750 gms to 1.0 kg was rather low (11%) in our study because very few babies fell in this group (7% of study population). It was observed that babies > 2 kg did not follow the inverse relationship between weight and frequency and showed a relatively higher incidence (21%). This can be explained by the fact that these babies had other risk factors in addition to being premature. The most vulnerable group prone to SEH-IVH are infants weighing 1.5 kg or less and the incidence increases with decreasing birth weight.<sup>26,51-67</sup> Studies revealed that more than 95% of the bleeds occurred during the first week of life.<sup>60,66,69,70</sup>

Factors found to be significantly associated with SEH-IVH by univariate analysis were respiratory distress syndrome, artificial ventilation, hypoxia, high concentration of inspired oxygen, patent ductus arteriosus, exchange transfusion, anemia and thrombocytopenia. Two of the study risk factors i.e. birth asphyxia and hypoglycemia were not found to be associated with SEH-IVH. On multivariate logistic model, only three factors retained their significance independently. These factors included RDS requiring artificial ventilation, exchange transfusion and thrombocytopenia. Of these, the babies on assisted ventilation with RDS carried the highest risk of SEH-IVH closely followed by babies undergoing exchange transfusion and thrombocytopenia. Other risk factors like noxious stimulation during routine care giving maneuvers, tracheal suctioning, rapid infusion of colloid, infusion of hyperosmolar solutions like sodium bicarbonate and coagulation disturbances.<sup>70-75</sup>

The pathogenesis of SEH-IVH is complex and multifactorial and influenced by intravascular, vascular and extravascular factors.

The optimum mode of delivery for preterm infants at risk for SEH-IVH is controversial. No consistent advantage has been documented with operative delivery compared with the vaginal route.<sup>76,77</sup>

Severe forms of SEH-IVH i.e. grade III and grade IV, carry a high mortality and morbidity. In the present study 71.4% of infants with SEH-IVH died whereas only 28.6 were either still admitted or discharged.

None of the infants with grade III and grade IV bleeds survived whereas the mortality for grades I & II was 37.5% and 62.5% respectively. This mortality is very high compared to that reported by Murphy.<sup>78</sup> In his study he categorized the mortality of different grades according to the weight of the preterm infants. Mortality of infants <750gms was 12%, 24%, 32% and 45% for grades I-IV respectively whereas for infants between 751-1500gms (weight more comparable to our study) it was 0%, 2%, 8% and 22%. This difference can be explained by better neonatal facilities available in the developed countries.

## Conclusions

1. Almost 1/5<sup>th</sup> of the babies between 28-34 weeks gestation developed bleed, a significantly high prevalence. This incidence increased with a decrease in gestational age and weight.
2. RDS with ventilation, exchange transfusion and thrombocytopenia were found to be major risk factors.
3. The peak time of occurrence of bleed was first three days of life with, first 24 hours being the most crucial.
4. Resource availability appears to influence aggressiveness of intervention and survival. Within our constraints, a number of interventions are possible to tackle the issue.
5. Antenatal steroids, careful management of premature deliveries, appropriate post-partum resuscitation, prevention of hypoxia and hypoglycemia, avoidance of hypertonic solutions like sodium bicarbonate, adequate control of seizures and gentle handling of small preterm babies are some of the factors easily applicable in all places caring for such infants.

## References

1. Volpe JJ. Germinal matrix-intraventricular hemorrhage. In: Volpe JJ, editor. *Neurology of the Newborn*, PA: Saunders Elsevier; 2008:517-88.
2. Kuban KCK. Intracranial hemorrhage. In: Cloherty JP, Stark AR. *Manual of neonatal care*. 4<sup>th</sup> ed. Philadelphia: Lippincott Raven, 1998:505-14.
3. Volpe JJ. Brain injury in the premature infant: Overview of clinical aspects, neuropathology, and pathogenesis. *Seminars in Pediatric Neurology*, 1998;(5:3):135-51.
4. Bolisetty S, Legge N, Bajuk B, Lui K. Preterm infant outcomes in New South Wales and the Australian Capital Territory. *Journal of Paediatrics and Child Health*. 2015 51(7):713-21.
5. Ahmann PA, Lazzara A, Dykes FD, Brann AW. Intraventricular hemorrhage in the high-risk preterm infant: Incidence and outcome. *Annals of Neurology*. 1980 ;7(2):118-24.
6. Papile LA, Burstein J, Burstein R. Evolution of subependymal and intraventricular hemorrhage: a study of infants with low birth weight. *Journal of Pediatrics*, 1978;92(4):529-34.

7. Holt PJ and Allan WC. The natural-history of ventricular dilatation in neonatal intraventricular hemorrhage and its therapeutic implications. *Ann Neurol* 1981;10:293-96.
8. Levene MI, Wigglesworth JS, Dubowitz VI. Cerebral structure and intraventricular hemorrhage in the neonate: a real-time ultrasound study. *Archives of Disease in Childhood*. 1981 ;56(6):416-24.
9. Philip AG, Allan WC, Tito AM, Wheeler LR. Intraventricular hemorrhage in preterm infants: declining incidence in the 1980s. *Pediatrics*. 1989 ;84(5):797-801.
10. Roland EH and Hill A. Intraventricular hemorrhage and posthemorrhagic hydrocephalus. Current and potential future interventions. *Clinics in Perinatology*, 1997;24(3):589-605.
11. Davis JM, Parad RB, Michele T. Pulmonary outcome at 1 year corrected age in premature infants treated at birth with recombinant human CuZn superoxide dismutase. *Pediatrics*. 2003;111(3):469-76.
12. Horbar JD, Badger GJ, Carpenter JH, Fanaroff AA. Trends in mortality and morbidity for very low birth weight infants, 1991–1999. *Pediatrics*. 2002;110(1):143-51.
13. Cust AE, Darlow BA, Donoghue DA. Outcomes for high risk New Zealand newborn infants : a population based, national study. *Archives of Disease in Childhood*,2003 ;88(1):15-22.
14. Larroque B, Marret S, Ancel PY. White matter damage and intraventricular hemorrhage in preterm infants. *Journal of Pediatrics*. 2003 ;143(4):477-83.
15. Hamrick SE, Miller SP, Leonard C. Trends in severe brain injury and neurodevelopmental outcome in premature newborn infants. *Journal of Pediatrics*, 2004;145(5):593-99.
16. Jain NJ, Kruse LK, Demissie K. Impact of mode of delivery on neonatal complications. *Journal of Maternal-Fetal & Neonatal Medicine*. 2009;22(6):491-500.
17. McCrea HJ and Ment LR. The diagnosis, management, and postnatal prevention of intraventricular hemorrhage in the preterms. *Clinics in Perinatology*. 2008;35(4):777-92.
18. Ballabh P. Intraventricular hemorrhage in premature infants: mechanism of disease. *Pediatric Research*, 2010 ;67(1):1-8.
19. Wilcox AJ and Skjaerven R. Birth weight and perinatal mortality: the effect of gestational age. *American Journal of Public Health*. 1992 ;82(3):378-82.
20. Horbar JD, McAuliffe TL, Adler SM. Variability in 28-day outcomes for very low birth weight infants. *Pediatrics*. 1988 ;82(4):554-59.
21. Horbar JD, Badger GJ, Lewit EM. Oxford Network. Hospital and patient characteristics associated with variation in 28-day mortality rates for very low birth weight infants. *Pediatrics*. 1997 ;99(2):149-56.
22. Fanaroff AA, Wright LL, Stevenson DK. Very-low-birth-weight outcomes of the National Institute of Child Health and Human Development Neonatal Research Network. *American Journal of Obstetrics and Gynecology*. 1995;173(5):1423-31.
23. Lee SK, McMillan DD, Ohlsson A, Pendray M, Synnes A. Variations in practice and outcomes in the Canadian NICU network: 1996–1997. *Pediatrics* , 2000;106(5):1070-79.
24. Darlow BA, Cust AE, Donoghue DA. Improved outcomes for very low birthweight infants: evidence from New Zealand national population based data. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2003 88(1):23-28.
25. De Vries LS, van Haastert IL, Rademaker KJ. Ultrasound abnormalities preceding cerebral palsy in high-risk preterm infants. *The Journal of Pediatrics*, 2004 ;144(6):815-20
26. Wilson-Costello D, Friedman H, Minich N, Fanaroff AA. Improved survival rates with increased neurodevelopmental disability for extremely low birth weight infants in the 1990s. *Pediatrics*. 2005;115(4):997-1003.
27. Adegoke SA, Olugbemiga AO, Bankole KP. Intraventricular hemorrhage in newborns weighing < 1500 g. *Annals of Tropical Medicine and Public Health*. 2014 ;7(1):48-51.
28. Kuperman AA, Kenet G, Papadakis E, Brenner B. Intraventricular hemorrhage in preterm infants: coagulation perspectives. *Seminars in Thrombosis and Hemostasis*, 2011 ; 37(7): 730-36.
29. Beverley DW, Chance GW, Inwood MJ, Schaus M, O'Keefe B. Intraventricular haemorrhage and haemostasis defects. *Archives of Disease in Childhood*, 1984 ;59(5):444-48.
30. Qureshi AI, Adil MM, Shafizadeh N. A 2-fold higher rate of intraventricular hemorrhage-related mortality in African American neonates and infants: Clinical article. *Journal of Neurosurgery: Pediatrics*. 2013;12(1):49-53.
31. Haroon A, Maheen H, Salat MS, Dileep D. Risk factors for intraventricular haemorrhage in preterm infants. *JPMA*. 2014;64(1146):112-15.
32. Köksal N, Baytan B, Bayram Y, Nacarküçük E. Risk factors for intraventricular haemorrhage in very low birth weight infants. *Indian Journal of Pediatrics*, 2002 ;69(7):561-64.
33. Aly H, Hammad TA, Essers J. Is mechanical ventilation associated with intraventricular hemorrhage in preterm infants?. *Brain and Development*. 2012 ;34(3):201-05.
34. Lee JY, Kim HS, Jung E, Kim ES, Shim GH, Lee HJ. Risk factors for periventricular-intraventricular hemorrhage in premature infants. *Journal of Korean medical science*. 2010 ;25(3):418-24.
35. Lu HY, Zhang Q, Wang QX, Lu JY. Contribution of histologic chorioamnionitis and fetal inflammatory response syndrome to increased risk of brain injury in infants with preterm premature rupture of membranes. *Pediatric Neurology*. 2016 ;61:94-98.
36. BilićČače I, Milardović A, BabićBožović I, Kolak M. Risk factors of intraventricular hemorrhage in preterm infants in the Neonatal Intensive Care Unit, University Department of Obstetrics and Gynecology, University Hospital Center Rijeka. *Medicina Fluminensis*. 2017;52(4.):515-26.
37. Kim KR, Jung SW, Kim DW. Risk factors associated with germinal matrix-intraventricular hemorrhage in preterm neonates. *Journal of Korean Neurosurgical Society*. 2014 ;56(4):334-37.
38. Lupton BA, Hill A, Whitfield MF, Carter C. Reduced platelet count as a risk factor for intraventricular hemorrhage. *American Journal of Diseases of Children*. 1988 ;142(11):1222-24.
39. Antoniuk S and Da Silva RV. Periventricular and intraventricular hemorrhage in the premature infants. *Revista de Neurologia*. 1999 ;31(3):238-43.
40. Kenny JD, Garcia-Prats JA, Hilliard JL. Hypercarbia at Birth: A Possible Role in the Pathogenesis of Intraventricular Hemorrhage. *Pediatrics*. 1978 ;62(4):465-67.
41. DiSalvo D. The correlation between placental pathology and intraventricular hemorrhage in the preterm infant. The Developmental Epidemiology Network Investigators. *Pediatric Research*, 1998 ;43(1):15-19.
42. Volpe JJ. Intraventricular hemorrhage in the premature infant—current concepts. Part I. *Annals of Neurology*. 1989 ;25(1):3-11.
43. Wei JC, Catalano R, Profit J, Gould JB. Impact of antenatal steroids on intraventricular hemorrhage in very-low-birth weight infants. *Journal of Perinatology* 2016; 36 (5): 352-56
44. Liggins GC, Howie RN. A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory

- distress syndrome in premature infants. *Pediatrics*. 1972 ;50(4):515-25.
45. Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth (Review). *Cochrane Database Syst Rev*. 2006;19:CD004454.
  46. Shalak L and Perlman JM. Hemorrhagic-ischemic cerebral injury in the preterm infant: current concepts. *Clinics in Perinatology*, 2002 29(4):745-63.
  47. Wells JT and Ment LR. Prevention of intraventricular hemorrhage in preterm infants. *Early human development*. 1995 ;42(3):209-33.
  48. Schmid MB, Reister F, Mayer B, Hopfner RJ. Prospective risk factor monitoring reduces intracranial hemorrhage rates in preterm infants. *DtschArztebl Int*. 2013 ;110(29-30):489-96.
  49. Ment LR, Oh W, Ehrenkranz RA, Philip AG, Duncan CC. Antenatal steroids, delivery mode, and intraventricular hemorrhage in preterm infants. *American Journal of Obstetrics and Gynecology*, 1995;172(3):795-800.
  50. Chaudhari S, Kinare AS, Kumar R, Pandit AN. Ultrasonography of the brain in preterm infants and its correlation with neurodevelopmental outcome. *Indian Pediatrics*, 1995 ;32(7):735-42.
  51. Soni JP, Gupta BD, Soni M, Gupta M, Dabi DR, Nema KR. Ultrasonic diagnosis of intracranial hemorrhage in high risk neonates. *Indian Pediatrics*, 1995 ;32:453-56.
  52. Sandler DL, Cooper PA, Bolton KD, Bental RY. Periventricular-intraventricular haemorrhage in low-birth-weight infants at Baragwanath Hospital. *South African Medical Journal*, 1994;84(1):26-29.
  53. Rehman M and Khan GRHI. Prevalence of periventricular/Intraventricular haemorrhage in premature infants. *Pakistan Pediatr J* 1998;22(4):145-47.
  54. Yakoob M. Characteristics of very preterm infants delivered at a tertiary care center in Pakistan. *Mymensingh Med J*. 2014 ;23(3)129-32.
  55. Debbarma R, De A, Debbarma S. Incidence of intracranial haemorrhage in low-birth weight infants and its outcome. *International Journal of Research in Medical Sciences*. 2016;4(10):4279-85.
  56. Mulindwa MJ, Sinyangwe S, Chomba E. Prevalence of intraventricular haemorrhage and associated risk factors in preterm neonates. *Medical Journal of Zambia*. 2012;39(1):16-21.
  57. Afjeh SA, Sabzehei MK, Fallahi M. Outcome of Very Low birth weight infants over 3 years report from an Iranian Center. *Iranian journal of pediatrics*. 2013;23(5):579.
  58. Sajjadian N, Fakhrai H, Jahadi R. Incidence of intraventricular hemorrhage and post hemorrhagic hydrocephalus in preterm infants. *Acta Medical Iranica*, 2010(4):260-62.
  59. Marba S, Caldas JP, Vinagre LE. Incidence of periventricular/intraventricular hemorrhage in very low birth weight infants. *Jornal de pediatria*, 2011;87(6):505-11.
  60. Ajmal F, Shaikat F, Hussein W. Intraventricular Haemorrhage Amongst Preterm Babies- Preventive Role of Dexamethasone. *Proceeding S.Z.P.G.M.I*. 2010;24(2):61-65.
  61. Sabir S, Ahmed Z, Ahmed I, Razzaq A. Frequency of intraventricular hemorrhage in premature infants. *Pakistan Armed Forces Medical Journal*. 2011(4): 61: 157-182.
  62. Rashid J, Shahid M, Anwar S, Sharaf D, Bhatti T. Frequency of Intraventricular Haemorrhage in Preterm Neonates. *Pak J Med Health Sci*, 2010;4(4):57578.
  63. Dubowitz LM, Dubowitz V, Goldberg C. Clinical assessment of gestational age in the newborn infant. *The Journal of Pediatrics*. 1970 ;77(1):1-5.
  64. Avery ME. Resuscitation. In: Avery ME, Tausch HW. *Schaffer's diseases of the newborn* 5<sup>th</sup> ed. Philadelphia: WB Saunders 1984:105.
  65. Kumar A, Bhat BV. Epidemiology of respiratory distress of newborns. *The Indian Journal of Pediatrics*. 1996 ;63(1):93-8.
  66. Dolfin T, Skidmore MB, Fong KW, Hoskins EM. Incidence, severity, and timing of subependymal and intraventricular hemorrhages in preterm infants. *Pediatrics*. 1983 ;71(4):541-6.
  67. Perlman JM, Rollins N, Burns D. Relationship between periventricular intraparenchymal echodensities and germinal matrix-intraventricular hemorrhage in the very low birth weight neonate. *Pediatrics*. 1993 ;91(2):474-80.
  68. Patridge JC, Babcock DS, Steichen JJ. Optimal timing for diagnostic cranial ultrasound in low-birthweight infants: detection of intracranial hemorrhage and intraventricular dilatation. *J Pediatr*, 1983;102:281-87.
  69. Paneth N, Pinto-Martin J, Gardiner J. Incidence and timing of germinal matrix/intraventricular hemorrhage in low birth weight infants. *American Journal of Epidemiology*, 1993 ;137(11):1167-76.
  70. Van De Bor M, Van Bel F, Lineman R. Perinatal factors and periventricular-intraventricular hemorrhage in preterm infants. *American Journal of Diseases of Children*. 1986;140(11):1125-30.
  71. Perlman JM, McMennamin JB, Volpe JJ. Fluctuating cerebral blood-flow velocity in respiratory-distress syndrome: relation to the development of intraventricular hemorrhage. *New England Journal of Medicine*. 1983 ;309(4):204-9.
  72. Goldberg RN, Chung D, Goldman SL, Bancalari E. The association of rapid volume expansion and intraventricular hemorrhage in the preterm infant. *The Journal of pediatrics*. 1980 ;96(6):1060-63.
  73. Andrew M, Castle V, Saigal S, Carter C, Kelton JG. Clinical impact of neonatal thrombocytopenia. *The Journal of Pediatrics*, 1987;110(3):457-64.
  74. Bonifacio L, Petrova A, Nanjundaswamy S, Mehta R. Thrombocytopenia related neonatal outcome in preterms. *The Indian Journal of Pediatrics*. 2007 ;74(3):269-74.
  75. Yulandari I, Rundjan L, Kadim M, Amalia P. The relationship between thrombocytopenia and intraventricular hemorrhage in neonates with gestational age < 35 weeks. *Paediatr Indonesiana*, 2016 ;56(4):242-50.
  76. Anderson GD, Bada HS, Shaver DC, Harvey CJ. The effect of caesarean section on intraventricular hemorrhage in the preterm infant. *American Journal of Obstetrics and Gynecology*. 1992 ;166(4):1091-101
  77. Strauss A, Kirz D, Modanlou HD. Perinatal events and intraventricular/subependymal hemorrhage in the very low-birth weight infant. *American Journal of Obstetrics and Gynecology*. 1985;151(8):1022-27.
  78. Murphy BP, Inder TE, Rooks V. Posthaemorrhagic ventricular dilatation in the premature infant: natural history and predictors of outcome. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2002 ;87(1):37-41.