

Pauly Matrix and Transformation Operators for Dirac System

T. N. Harutyunyan, H. H. Azizyan

Yerevan State University, e-mail: hartigr@yahoo.co.uk
 Armenian State Agrarian University, e-mail: hermineazizyan@mail.ru

Let $\sigma_1 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$, $\sigma_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $\sigma_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ are well known Pauly matrix and $E = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. It is known that the solution $y = \varphi(x, \lambda, \alpha)$ of Cauchy problem

$$\left\{ \sigma_1 \frac{1}{i} \frac{d}{dx} + \sigma_2 p(x) + \sigma_3 q(x) \right\} y = \lambda y, \quad \lambda \in C$$

$$y(0) = \begin{pmatrix} \sin \alpha \\ -\cos \alpha \end{pmatrix},$$

can be represented in the form $\left(\varphi_0(x, \lambda, \alpha) = \begin{pmatrix} \sin(\lambda x + \alpha) \\ -\cos(\lambda x + \alpha) \end{pmatrix} \right)$

$$\varphi(x, \lambda, \alpha) = \varphi_0(x, \lambda, \alpha) + \int_0^x K(x, t) \varphi_0(t, \lambda, \alpha) dt = (E + K) \varphi_0.$$

Operator $E + K$ is called the transformation operator. Under different conditions on scalar functions p and q this operator and his kernel $K(x, t)$ was investigated in different papers (see [1]-[6]).

Theorema. *Let $p, q \in L_{loc}^1(0, \infty)$. Then the kernel $K(x, t)$ and the kernel $H(x, t)$ of inverse operator $\varphi_0(x, \lambda) = \varphi(x, \lambda) + \int_0^x H(x, t) \varphi(t, \lambda) dt$ can be represented in the form*

$$K(x, t) = a\sigma_1 + b\sigma_2 + c\sigma_3 + d \cdot E$$

$$H(x, t) = \tilde{a}\sigma_1 + \tilde{b}\sigma_2 + \tilde{c}\sigma_3 + \tilde{d} \cdot E,$$

where the functions (of two variables (x, t)) a, b, c, d and $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$ are represented by functions p and q .

References

- [1] Gasymov M. G., Levitan B. M., Determination of a differential equation by two of its spectra, *Usp. Mat. Nauk* v.19, N 2, 1964, pp.3-63
- [2] Marchenko V. A., *Sturm-Liouville operators and their applications*, Naukova Dumka, Kiev, 1977.
- [3] Melik-Adamyanyan F. E., On the canonic differential operators in Hilbert space, *Izvest. AN Arm. SSR, Mathematics*, v.XII, N 1, pp.10-30.
- [4] Levitan B. M., Sargsyan I. S., *Sturm-Liouville and Dirac operators*, Nauka, Moscow, 1988.
- [5] Harutyunyan T. N., Transformations operators for canonic Dirac system, *Differentialnie uravneniya*, v.44, N 8, 2008, pp. 1011-1021.
- [6] Albeverio S., Hryniv R., Mikituk Ya., Inverse spectral problems for Dirac operators with summable potentials, *Rus. J. of Math. Phys.* v.12, N 4, 2005, pp.406-423.