

ROMANIAN  
NEUROSURGERY

Vol. XXXIV | No. 2      June 2020

A technical note.  
The role of lilliequist membrane  
fenestration during the pterional  
approach for anterior circulation  
aneurysm clipping

Samer S. Hoz,  
Zaid Aljuboori,  
Zahraa F. Al-Sharshahi



# A technical note. The role of lilliequist membrane fenestration during the pterional approach for anterior circulation aneurysm clipping

Samer S. Hoz<sup>1</sup>, Zaid Aljuboori<sup>2</sup>,  
Zahraa F. Al-Sharshahi<sup>3</sup>

<sup>1</sup> MBChB, FABMS/Neurosurgery, Department of Neurosurgery, Neurosurgery Teaching Hospital, Baghdad, IRAQ

<sup>2</sup> MD, Neurosurgery, Department of Neurosurgery, University of Louisville, Kentucky, USA

<sup>3</sup> MB, BCh BAO (NUI, RCSI), LRCP & SI, Royal College of Surgeons in Ireland (Alumnus), Dublin, IRELAND

## ABSTRACT

The Lilliequist membrane is a critical membrane located at the base of the brain separating the supratentorial from the infratentorial cisterns. The advantages of Lilliequist membrane fenestration as a critical part of the pterional trans-Sylvian approach for ruptured anterior circulation aneurysm clipping is not well established. We demonstrated that the fundamental role of Lilliequist membrane fenestration is brain relaxation through the egress of CSF that is not usually gained from other modalities (e.g., placement of a lumbar drain, fenestration of lamina terminalis) in this specific setting.

## BACKGROUND

Dissection through the arachnoid planes and fenestration of the accessible cisterns are the cornerstones of modern microsurgery while dealing with intracranial lesions.

The Lilliequist membrane is a horizontal arachnoid membrane attached to the dorsum sellae anteriorly, the mammillary body superiorly and the basilar artery posteriorly. It is bounded by the oculomotor nerve on each side. Embryologically, the Lilliequist membrane is considered a remnant of the primary tentorium (1).

The first recognition of the Lilliequist membrane was in 1875 by Key and Retzius (2). Eight decades later, it was rediscovered by an eminent Swedish radiologist named Bengt Lilliequist (1923-1972) as part of his doctoral thesis titled "The subarachnoid cisterns. An anatomical and roentgenologic study" (3). The Lilliequist membrane became a universally accepted medical eponym and officially settled through

---

**Keywords**  
lilliequist membrane  
fenestration,  
pterional approach,  
anterior circulation aneurysm

---



Corresponding author:  
Zahraa Al-Sharshahi

MB, BCh BAO (NUI, RCSI), LRCP & SI,  
Royal College of Surgeons in Ireland  
(Alumnus), Dublin, Ireland

Zahraaalsharshahi@rcsi.com

**Copyright and usage.** This is an Open Access article, distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited.

The written permission of the Romanian Society of Neurosurgery must be obtained for commercial re-use or in order to create a derivative work.

ISSN online 2344-4959  
© Romanian Society of  
Neurosurgery



First published  
June 2020 by  
London Academic Publishing  
[www.lapub.co.uk](http://www.lapub.co.uk)

Yasargil's paper "Anatomical observations of the subarachnoid cisterns of the brain during surgery" in 1976 (34). Since that time, the Lilliequist membrane underwent detailed descriptions in literature based on cadaveric and radiological studies (4-8).

In this article, we describe the technique and benefits of Lilliequist membrane fenestration to achieve brain relaxation during the pterional trans-Sylvian approach for ruptured anterior circulation aneurysm clipping.

#### PERTINENT ANATOMY OF THE LILLIEQUIST MEMBRANE

Intracranially, the arachnoid membranes can be either outer or inner membranes. The outer membranes include both the convexity and the basal arachnoid membranes, while the inner membranes include the trabecular arachnoid membranes which encase the neurovascular bundle within the cisterns (8). The Lilliequist membrane is an inner rather than outer arachnoid membrane (9).

There is some controversy regarding the number of layers of the Lilliequist membrane. The early literature published by Lilliequist and Yasargil described a single-layer membrane (5,10-12). Later studies by Matsuno and Rhoton (13,14) assumed the two-layer model (the diencephalic membrane and the mesencephalic membrane). Recently, the existence of a third layer was described, but it was questionable as to whether this third layer represents a distinct layer or sparse arachnoid trabeculae (8).

From the dorsum sellae, the diencephalic membrane extends superoposteriorly to the mammillary bodies between the chiasmatic and the interpeduncular cisterns and can sometimes be identified radiologically in the sagittal MRI view. The mesencephalic membrane is a thin sheet that extends inferoposteriorly to the pontomesencephalic junction, between the interpeduncular and the prepontine cisterns. Laterally, the Lilliequist membrane is attached to the oculomotor nerve and/or the uncus and tentorium. The Lilliequist membrane is closely related and variably penetrated by the posterior communicating and basilar arteries and their perforating branches (4-8,10-14).

#### THE GENERAL INDICATIONS FOR LILLIEQUIST MEMBRANE FENESTRATION

Lilliequist Membrane fenestration plays an important

role in multiple procedures related to microscopic or endoscopic cranial surgery:

1. In surgery for tumors in the sellar and parasellar regions such as craniopharyngiomas, meningiomas, basilar tip aneurysms, and arachnoid cysts, Lilliequist membrane fenestration provides an effective surgical corridor and promotes brain relaxation through CSF egress. (1,15,17).
2. Lilliequist membrane fenestration is one of the most critical steps for endoscopic third ventriculostomy success.
3. Some reports consider the microscopic fenestration of the basal subarachnoid cisterns as an effective method to decrease the intracranial pressure in acute intra-operative brain swelling and also in severe traumatic brain injury. These reports advocate the inclusion of Lilliequist membrane fenestration as a mandatory procedural step to achieve a good surgical outcome (16,18)

#### THE TECHNIQUE FOR LILLIEQUIST MEMBRANE FENESTRATION DURING THE PTERIONAL APPROACH FOR ANTERIOR CIRCULATION ANEURYSM CLIPPING

The pterional approach is the preferred approach for the clipping of most anterior circulation aneurysms except for the distal anterior cerebral artery aneurysm. Using the trans-Sylvian route, the Lilliequist membrane can be approached laterally. Opening the Lilliequist membrane can be achieved through different corridors; namely, the carotid-oculomotor triangle, the optico-carotid triangle and even through the interoptic space. Fenestration through the carotid-oculomotor space is the most convenient, relatively wide and safe technique.

After dural opening, the Sylvian fissure is widely opened from inside to outside on the frontal side of the Sylvian veins and the proximal segment of the middle cerebral artery until the internal carotid artery bifurcation. Frontal and temporal retractors are gently applied; these retractors should "hold" the brain tissue with intermittent retraction to avoid vascular compromise. The frontal lobe is elevated, and the temporal lobe is retracted gently downward and posteriorly. The carotid cistern is opened and the thickened arachnoid fibers over the origins of the anterior and middle cerebral arteries are released. Once the 3rd nerve is identified, the carotid-oculomotor triangle is dissected and subsequently

the deeper membrane of Lilliequist can be identified. As the exposure is deepened, the surgeon will extend the arachnoid opening laterally over the third nerve. Sectioning of the Lilliequist membrane should be started from its anterior part as there is a large space behind the membrane. Then, the interpeduncular cistern is opened through caudal dissection of the Lilliequist membrane between the oculomotor nerve and the internal carotid artery to release the cerebrospinal fluid. At this point, the posterior communicating and the anterior choroidal arteries may be visualized as they arise from the posterior surface of the internal carotid artery (the posterior communicating artery courses perpendicular to the Lilliequist membrane whereas the anterior choroidal artery courses obliquely into the crural cistern) (12).

The egress of the CSF and the identification of the basilar artery within the inter-peduncular cistern is the critical step that ensures the completion of the fenestration process. Additional steps were suggested by some reports including irrigation and mechanical cleaning of blood clots beyond the Lilliequist membrane to minimize the possibility of postoperative vasospasm and hydrocephaly (15,16).

Sharp dissection is always advised to minimize the risk of uncontrolled bleeding from tearing of the vessel wall. Also, the blunt and vigorous dissection may cause disastrous consequences by injuring the hypothalamus either directly or indirectly, by affecting its blood supply. Structures at risk during this step include the posterior communicating artery or its perforators, the oculomotor nerve or its blood supply and the proximal part of the superior cerebellar artery.

In the case of SAH due to a ruptured aneurysm, several studies have described the effectiveness of Lilliequist membrane fenestration, either alone or in tandem with lamina terminalis fenestration, in reducing the incidence of post-SAH hydrocephalus, (9, 12, 15, 19-21). However, None of these studies focused on the effect of Lilliequist membrane fenestration on brain relaxation specifically, which is crucial in such complex procedures regardless of the presence of hydrocephaly.

Lilliequist membrane fenestration will create a free connection between the supratentorial and infratentorial compartments which represents the crucial advantage of Lilliequist membrane fenestration over the opening of the cisterns or

lamina terminalis; this fact was supported by Yaşargil and Winkler-Lawton et al reports through the "Fifth ventricle theory" (12, 19). The fifth ventricle theory can be described as the following; in cases of aneurysmal SAH, the Lilliequist membrane will entrap blood clots and it will be inflamed and thickened, thus the interpeduncular and prepontine cisterns will be isolated and dilated with CSF creating a "fifth ventricle". At this point, Lilliequist membrane fenestration will result in dramatic CSF egress with striking brain relaxation which renders further surgical steps safer and more effective. Castro-Flores et al have suggested that in several patients, even after the opening of the supratentorial cisterns, the CSF pathway will remain constricted until the Lilliequist membrane is opened (15).


#### AN ILLUSTRATIVE CASE

A 59-year male presented to the emergency department with an altered level of consciousness (Glasgow Coma Scale:9) of sudden onset. Initial resuscitation was done and the patient was transferred to the intensive care unit. A brain CT scan showed left gyrus rectus hemorrhage. Contrasted brain CT scan and CT angiography revealed complex ruptured anterior communicating artery aneurysm with a wide neck, and hypoplastic left Proximal anterior cerebral artery. The anterior communicating artery was vertically oriented, and the anterior cerebral arteries were connected distally to the anterior communicating artery, with two Murphy's teats (Figure 1A).


Surgical clipping through the standard right pterional trans-Sylvian approach was chosen. After dural opening, around 20cc of CSF was drained through a lumbar drain. A microsurgical arachnoid dissection was performed using an operative microscope; starting with the opening the proximal Sylvian fissure and the carotid cisterns; there was minimal egress of CSF.

The Lilliequist membrane was identified and opened lateral to the supraclinoid internal carotid artery through the carotid-oculomotor space. At this point, there was a striking CSF egress that filled the surgical field despite repetitive suctioning. This CSF egress had a profound effect on brain relaxation, provided us a wide operative corridor, and rendered the subsequent steps of aneurysm dissection and clipping more feasible. (Figure 2).

The postoperative course was uneventful and the patient gradually improved and was discharged with no neurological deficits (Figure 1B and C).



**Figure 1.** Illustrative case of a ruptured anterior communicating artery aneurysm. **A:** contrasted CT scan of the brain showing the left gyrus rectus due to a ruptured anterior communicating artery aneurysm. **B:** An intraoperative microscopic view showing a two-clip reconstruction for the aneurysmal neck through the right pterional trans-Sylvian approach. **C:** An early postoperative brain CT scan showing the clipping result.



**Figure 2.** Stages of Lilliequist membrane fenestration through the right pterional trans-Sylvian approach (Intraoperative microscopic view). The Lilliequist membrane identified as a whitish layer lateral to the supraclinoid internal carotid artery through the carotid-oculomotor space. **(A)** Opening the Lilliequist membrane resulted in striking CSF egress that filled the surgical field several times **(B)**. (+): pointer on the Lilliequist membrane, F: Frontal lobe, T: Temporal lobe, C: internal carotid artery, ON: optic nerve.

#### CONCLUSION

Fenestration of the Lilliequist membrane has a fundamental role during the pterional trans-Sylvian

approach for ruptured anterior circulation aneurysm clipping as it allows brain relaxation through egress of CSF that is not usually achieved through other

modalities (e.g, placement of a lumbar drain, fenestration of lamina terminalis). Further studies are needed to verify this observation towards making the fenestration of the Lilliequist membrane a recommended surgical step rather than an optional one.

#### ABBREVIATIONS

MRI: magnetic resonance imaging;

CSF: cerebrospinal fluid, SAH: subarachnoid hemorrhage.

#### DECLARATION

Consent for publication: Our local institutional consent was obtained from the patient.

Competing interests: The authors declare that they have no competing interests.

Funding: Nothing to declare.

#### REFERENCES

1. Froelich SC, Aziz KM, Cohen PD, Loveren HR, Keller JT. Microsurgical and endoscopic anatomy of Lilliequist's membrane: a complex and variable structure of the basal cisterns. *Operative Neurosurgery*. 2008 Jul 1;63(suppl\_1):ONS1-9.
2. Retzius MG, Key A. Studien in der Anatomie des Nervensystems und des Bindegewebes, von Axel Key und Gustaf Retzius,... Samson und Wallin; 1875.
3. Connor Jr DE, Nanda A. Bengt Lilliequist: life and accomplishments of a true renaissance man. *Journal of neurosurgery*. 2017 Feb;126(2):645-9.
4. Matsuno H, Rhoton Jr AL, Peace D. Microsurgical anatomy of the posterior fossa cisterns. *Neurosurgery*. 1988 Jul 1;23(1):58-80.
5. Brasil AV, Schneider FL. Anatomy of Lilliequist membrane. *Neurosurgery* 1993; 32:956 -960.
6. Zhang M, An PC. Lilliequist's membrane is a fold of the arachnoid mater: study using sheet plastination and scanning electron microscopy. *Neurosurgery*. 2000 Oct 1;47(4):902-9.
7. Vinas FC, Panigrahi M. Microsurgical anatomy of the Lilliequist's membrane and surrounding neurovascular territories. *min-Minimally Invasive Neurosurgery*. 2001 Jun;44(02):104-9.
8. Lü J, Zhu XI. Microsurgical anatomy of Lilliequist's membrane. *min-Minimally Invasive Neurosurgery*. 2003 Jun;46(03):149-54.
9. Zhang XA, Qi ST, Huang GL, Long H, Fan J, Peng JX. Anatomical and histological study of Lilliequist's membrane: with emphasis on its nature and lateral attachments. *Child's Nervous System*. 2012 Jan 1;28(1):65-72.
10. Lilliequist B. The anatomy of the subarachnoid cisterns. *Acta radiologica*. 1956 Jul(1-2):61-71.
11. Epstein BS. The role of a transverse arachnoid membrane within the interpeduncular cistern in the passage of pantopaque into the cranial cavity. *Radiology*. 1965 Nov;85(5):914-20.
12. Yasargil MG. *Microneurosurgery*. New York: Thieme Verlag, 1984; 27: 15±53
13. Matsuno H, Rhoton Jr AL, Peace D. Microsurgical anatomy of the posterior fossa cisterns. *Neurosurgery*. 1988 Jul 1;23(1):58-80.
14. Rhoton AL. The posterior fossa cisterns. *Neurosurgery* 2000; 47 (Suppl): 297 ± 297
15. Juan Antonio Castro-Flores., et al. "Efficacy of the Lamina Terminalis Fenestration Associated With the Lilliequist Membrane Fenestration in Reducing Shunt-Dependent Hydrocephalus Following Aneurysm Surgery in the Acute Phase of Aneurysmal Subarachnoid Hemorrhage". *EC Neurology* 2.4 (2015): 162-166.
16. Cherian I, Grasso G, Bernardo A, Munakomi S. Anatomy and physiology of cisternostomy. *Chinese Journal of Traumatology*. 2016 Feb 1;19(1):7-10.
17. Miyajima M, Arai H, Okuda O, Hishii M, Nakanishi H, Sato K. Possible origin of suprasellar arachnoid cysts: neuroimaging and neurosurgical observations in nine cases. *Journal of neurosurgery*. 2000 Jul;93(1):62-7.
18. Abdulqader MN, Al-Tameemi AH, Salih H, Hoz SS, Al Ramadan AH, Salazar LR. Acute intra-operative brain swelling managed effectively with emergency basal cisternostomy: A case report. *Journal of Acute Disease*. 2018 Jan 1;7(1):43.
19. Winkler EA, Burkhardt JK, Rutledge WC, Rick JW, Partow CP, Yue JK, Birk H, Bach AM, Raygor KP, Lawton MT. Reduction of shunt dependency rates following aneurysmal subarachnoid hemorrhage by tandem fenestration of the lamina terminalis and membrane of Lilliequist during microsurgical aneurysm repair. *Journal of neurosurgery*. 2017 Dec 15:1-7.
20. Ultrastrüktürel HS. Ultrastructural changes in the Lilliequist membrane in the hydrocephalic process and its implications for the endoscopic third ventriculostomy procedure. *Turkish neurosurgery*. 2011;21(3):359-66.
21. Wang SS, Zheng HP, Zhang FH, Wang RM. Microsurgical anatomy of Lilliequist's membrane demonstrating three-dimensional configuration. *Acta neurochirurgica*. 2011 Jan 1;153(1):191-200.