



# Pathways of metastatic spread in meningiomas

A. I. Cucu<sup>1</sup>, Dana Mihaela Turliuc<sup>1,2</sup>, Claudia Florida Costea<sup>1,2</sup>,  
B. Costachescu<sup>1,2</sup>, Andreea Eliza Malaimare<sup>3</sup>,  
L. A. Blaj<sup>1</sup>, Violeta Trandafir<sup>1,4</sup>, Cristina Danca<sup>1</sup>,  
I. Poeata<sup>1,2</sup>

<sup>1</sup> "Prof. Dr. N. Oblu" Clinic Emergency Hospital, Iasi, ROMANIA

<sup>2</sup> "Grigore T. Popa" University of Medicine and Pharmacy, Iasi, ROMANIA

<sup>3</sup> Individual Medical Practice, ROMANIA

<sup>4</sup> "SF. SPIRIDON" EMERGENCY CLINICAL HOSPITAL IAȘI, ROMANIA

## ABSTRACT

Meningioma is a common intracranial neoplasm derived from meningotheial cells, and it is generally associated with a benign clinical course. In spite of this, the malignant behaviour of these tumours as the occurrence of extracranial meningioma metastases in different organs is described in literature: lung and pleura, spine and other bones, abdominal organs, lymph nodes or even skin. The aim of this review is to analyse the pathways of metastatic spread of the intracranial meningioma tumour cells towards different organs.

## INTRODUCTION

Meningiomas are the most common central nervous system tumours in adults, making up approximately 30% of all the intracranial tumours, with an increasing incidence (14, 57). According to the mitotic activity and tumour differentiation, the World Health Organization (WHO) grading meningiomas as it follows: grade I, grade II and grade III, the last two being characterized by a more aggressive behaviour, a high risk of recurrence (12, 39) and even metastasis. These extracranial meningioma metastases (EMM) are more frequently associated with atypical meningiomas (Table 1) or anaplastic meningiomas (20, 27, 59). Although typical meningiomas are benign solitary intracranial neoplasms, they can cause extracranial metastasis, a rare phenomenon found only in 0.1% of cases (15, 23, 26, 35, 37). According to epidemiological data, about one in 1000 cases of meningiomas metastasize (35, 56).

The most frequent sites of EMM are: the lung (60%-70%), abdominal viscera (the liver most frequently) (30-40%), pleura (23%), lymph nodes (14-20%) and bones (10%) (3, 6, 21, 38, 42, 50, 60, 66). Other rare localizations were identified in the parotid gland (17), skin (32, 51), deep soft tissue (9, 18, 44, 48, 64, 70), kidney, spleen, thyroid and adrenal gland (64).

## Keywords

meningioma, metastasis,  
malignant,  
WHO classification



Corresponding author:  
**Dana Mihaela Turliuc**

"Grigore T. Popa" University of  
Medicine and Pharmacy,  
Iasi, Romania

turliuc\_dana@yahoo.com

**Copyright and usage.** This is an Open Access article, distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited.

The written permission of the Romanian Society of Neurosurgery must be obtained for commercial re-use or in order to create a derivative work.

ISSN online 2344-4959  
© Romanian Society of  
Neurosurgery



First published  
March 2019 by  
London Academic Publishing  
[www.lapub.co.uk](http://www.lapub.co.uk)

Due to the rare nature of EMM, this is a challenging diagnosis for physicians and pathologists (25). Because the treatment has not yet been standard-

ized and a management protocol has not been developed, the prognosis of these patients remains unknown (3, 42, 50).

TABLE 1. Review of publications on extracranial metastasis of atypical meningiomas

Author	Gender, Age	Intracranial Localization	Metastasis Localization
Andric <i>et al.</i> (5)	F 52	parasagittal	lung
Baek <i>et al.</i> (9)	M 44	nasal septum	chest wall, parietal region
Drummond <i>et al.</i> (19)	M 66	frontal	lung
Doxtader <i>et al.</i> (18)	M 8	parasagittal	subcutaneous tissue of neck, lymph nodes
Kim <i>et al.</i> (41)	M 68	parietal	skull bone
Lanfranchi <i>et al.</i> (43)	M 74	not specified	lung, liver
Pinsker <i>et al.</i> (55)	M 76	frontal	cervical spine

#### METASTATIC SPREAD

Although all types of meningioma, including the grade I meningiomas can metastasize, EMM is more common in the atypical (5%) and anaplastic (30%) histopathological subtypes (7). Nevertheless, some authors believe that these percentages are overestimated, firstly because some of the reported cases probably are metastatic angioblastic meningiomas and not genuine meningioma metastases. Moreover, the angioblastic meningioma is now classified as haemangiopericytoma (25, 28, 34, 46). Thus, considering all these modifications, the 5% rate of metastasis in atypical meningiomas does not coincide with the current WHO classification (25). However, the true prevalence of EMM is unknown at present (64). As for the time interval between the initial diagnosis of the meningioma and that of extracranial metastasis, it varies from 3 months to 30 years (40).

Although throughout time, some authors emphasized tumour necrosis, blood vessel invasion, high cellularity, cellular heterogeneity, high mitosis rate and nuclear pleomorphism in the occurrence of EMM (1, 20, 54), these criteria apply to malignant tumours and do not explain metastatic disease in benign meningiomas (64). In spite of this, the exact etiology of metastatic spread in meningioma is still unknown (7, 22).

Initially, the surgical resection was reported as one of the most important risk factors for iatrogenic metastasis of histologically aggressive meningiomas (2). Hence, it was hypothesised that surgery causes the seeding of the lymphatics and vascular channels (including that of the scalp), resulting in extracranial metastatic disease via haematogenous or lymphatic route (31). Besides, other studies showed that tumour cells can propagate with no previous surgery (25, 36), considering that the invasion of adjacent venous sinuses can lead to extracranial spread of meningioma, especially when there is no history of surgery (23).

**Lung metastasis.** As mentioned before, the lung is the most common site of the EMM. In these cases, the seeded tumour cells access dural venous sinuses and cranial veins, then diffuse through the azygos system into pulmonary circulation and then they can reach either the lung or the pleura (21, 32, 40). Most EMM were found along the jugular vascular drainage, such as in the cervical lymph nodes, parotid and thyroid gland and finally in lung and pleura (64).

**Bone metastasis.** Bone is one of the least common sites of meningioma metastasis (38, 60). In a review of meningioma metastasis in the bone, Khan et al. found out that half of the patients had multiple areas of bone metastatic disease, while the

other half had solitary bone involvement (40). The most common region was the axial skeleton, namely the thoracolumbar spine and sacrum (40), while metastases in the extra-axial bone were rarely identified (22). The most common involvement of the spine can be explained by the connection between the dural venous sinuses and the vertebral venous system (10). This route via the paravertebral venous plexus may also play a role in the metastatic spread across the kidney or adrenal glands (64).

Of the non-axial skeleton, the femur was the most common long bone site of involvement (40), and this metastasis location is more difficult to explain, even if some authors suggested the extension of the tumour via arterial spread (68). The bone metastatic disease is a lytic bone lesion and may result in pathological fractures or collapse of the vertebral bodies (40).

Liver metastasis. As far as the dissemination pathways for hepatic metastasis are concerned, some authors believe that this lies in the dissemination through the vertebral venous system connected with the veins of the thoracoabdominal wall or dissemination of tumour cells of the bone metastasis to the liver (24, 25). The liver can be involved whether the metastases pass through the right atrium towards the inferior vena cava and further into the hepatic veins (64). In literature, cases of hepatic metastasis of meningioma via a ventriculoperitoneal shunt through the cerebrospinal fluid route were cited (51).

Scalp metastasis. Surgical seeding of tumour cells in meningioma surgery is a rare entity and it usually involves soft tissues near the craniotomy site (8). In literature, 19 such cases of scalp metastasis of intracranial meningiomas were reported (8).

Scalp metastasis of meningioma were firstly reported by Harvey Cushing, with an incidence of 1.21% cases in a cohort of 313 patients, which results in 4 patients suffering from this condition. In all of these cases, scalp metastases were found in surgical scar many years after the first surgery (4, 30, 49, 69). Different series reported percentages of 3% in a cohort of 119 patients with WHO grade II and III meningiomas (53) and others of 1.2% (8).

The suggested mechanism is intraoperative seeding and it may apply to all histological grades of meningiomas. Avecillas-Chasin et al. considers that immunosuppression, radiation therapy, CSF fistulae and multiple reoperations with subsequent surgical

wounds problems (8). Nevertheless, the reason for extracranial tissue invasions is still unclear.

Carrying out multiple interventions can be a risk factor for the extracranial spreading of intracranial tumours, in that surgical bone deteriorates the natural barriers for intracranial tumour dissemination, providing access to the blood vessels and lymphatic system (8, 63). The higher-grade histopathological meningioma also contributes to this in case of tumours which have the tendency to produce greater levels of vascular endothelial growth factor and angiogenesis, thus favouring vascular invasion and metastatic behaviour (29).

Piecemeal resection was also presented as a risk factor of extracranial dissemination of tumour cells in WHO grade III meningiomas (8). Although initially, it was suggested that the latency period of meningioma metastasis is influenced by the histological grade (2), the significant variability over this latency period proved that this is not related only to the histological grade of meningioma (2, 16, 30, 45, 47, 52, 61, 62, 65, 67, 69).

With regard to preventing the spread of tumour cells, some authors recommended: air-tight closure of dura mater, changing the gloves and surgical instruments for wound closure after the intracranial phase, replacing the bone flap and saline irrigation, wound abundance before the closure (4, 33).

Treatment strategies. The treatment for meningioma metastasis has yet to be set out. In spite of this, metastectomy seems to improve the prognosis, especially when the tumour is a low-grade meningioma. As for the chemotherapy with hydroxyurea, vincristine, cyclophosphamide and doxorubicin, this has a limited efficiency with progression under chemotherapy (25). Moreover, some immunotherapies with interferon- $\alpha$  and somatostatin analogue were tested lately, but with limited effects (11, 58). Nonetheless, the choice treatment in meningiomas remains the gross total resection and adjuvant radiotherapy (13).

## CONCLUSIONS

Because of the fact that EMM were rarely reported, there are no guidelines regarding the staging or treatment of EMM. Metastatic spread remains a therapeutic challenge and the treatment must be multidisciplinary evaluated. In addition, this data highlights the fact that metastatic dissemination of meningiomas is possible even under conditions of

benign histopathological grades and thus organ donation should be considered.

#### REFERENCES

1. Adlakha A, Rao K, Adlakha H, et al. Meningioma metastatic to the lung. *Mayo Clin Proc.* 1999; 74(11): 1129-1133.
2. Akai T, Shiraga S, Iizuka H, et al. Recurrent meningioma with metastasis to the skin incision--case report. *Neurol Med Chir (Tokyo).* 2004; 44(11): 600-602.
3. Alexandru D, Glantz MJ, Kim L, et al. Pulmonary metastases in patients with recurrent, treatment-resistant meningioma: prognosis and identification by <sup>111</sup>Indium-octreotide imaging. *Cancer.* 2011; 117(19): 4506-4511.
4. Allan RS. Scalp metastasis from glioblastoma. *J Neurol Neurosurg Psychiatry.* 2004; 364 75(4): 559.
5. Andric M, Dixit S, Dubey A, et al. Atypical meningiomas--a case series. *Clin Neurol Neurosurg.* 2012; 114: 699-702.
6. Asioli S, Senetta R, Maldì E, et al. "Benign" metastatic meningioma: clinico-pathological analysis of one case metastasising to the lung and overview on the concepts of either primitive or metastatic meningiomas of the lung. *Virchows Arch.* 2007; 450: 591-594.
7. Ather Enam S, Abdulrauf S, Mehta B, et al. Metastasis in meningioma. *Acta Neurochir.* 1996; 138: 1172-1178.
8. Avelillas-Chasin JM, Saceda-Gutierrez J, Alonso-Lera P, et al. Scalp metastases of recurrent meningiomas: aggressive behavior or surgical seeding? *World Neurosurg.* 2015; 84(1): 121-131.
9. Baek BJ, Shin JM, Lee CK, et al. Atypical primary meningioma in the nasal septum with malignant transformation and distant metastasis. *BMC Cancer.* 2012; 12:275.
10. Batson OV. The Function of the vertebral veins and their role in the spread of Metastases. *Ann Surg.* 1940; 112(1): 138-149.
11. Chamberlain MC, Barnholtz-Sloan JS. Medical treatment of recurrent meningiomas. *Expert Rev Neurother.* 2011; 11: 1425-1432.
12. Cucu AI, Costea CF, Poata I, et al. Prognostic factors in atypical meningioma. *Rom Neurosurg.* 2017; 31 (2): 165-171.
13. Cucu AI, Turliuc MD, Costea CF, et al. Atypical meningiomas of cerebellopontine angle. A five case series. *Rev Med Chir Soc Med Nat Iasi.* 2018; 122:533-545.
14. Cucu AI, Costea CF, Carauleanu A, et al. Meningiomas related to the Chernobyl irradiation disaster in North-Eastern Romania between 1990 and 2015. *Revista de Chimie Bucharest* 2018; 69: 1562-1565
15. D'Aiuto M, Veronesi G, Pelosi G, et al. Two-year survival after multiple bilateral lung metastasectomies for cranial meningioma. *Ann Thorac Surg.* 2005; 80: 1129-1130.
16. Darwish B, Munro I, Boet R, et al. Intraventricular meningioma with drop metastases and subgaleal metastatic nodule. *J Clin Neurosci.* 2004; 11(7): 787-791.
17. Dmytriw AA, Gullane P, Bartlett E, et al. Parotid gland metastasis originating from malignant meningioma. *Clin Imaging.* 2013; 37(4): 740-743.
18. Doxtader EE, Butts SC, Holsapple JW, et al. Aggressive pediatric meningioma with soft tissue and lymph node metastases: a case report. *Pediatr Dev Pathol.* 2009; 12(3): 244-248.
19. Drummond KJ, Bittar RG, Fearnside MR. Metastatic atypical meningioma: case report and review of the literature. *J Clin Neurosci.* 2007; 7: 69-72.
20. Enam SA, Abdulrauf S, Mehta B, et al. Metastasis in meningioma. *Acta Neurochir (Wien).* 1996; 138: 1172-1178.
21. Erman T, Hanta I, Hacıyakupoglu S, et al. Huge bilateral pulmonary and pleural metastasis from intracranial meningioma: a case report and review of the literature. *J Neurooncol.* 2005; 74: 179-181.
22. Estanislau ES, de Carvalho GTC, Reis BL, et al. Malignant meningioma with extracranial metastases. *Arq Neuropsiquiatr.* 2009; 67: 730-732.
23. Fabi A, Nuzzo C, Vidiri A, et al. Bone and lung metastases from intracranial meningioma. *Anticancer Res.* 2006; 26: 3835-3837.
24. Figueroa BE, Quint DJ, McKeever PE, Chandler WF. Extracranial metastatic meningioma. *Br J Radiol.* 1999; 72: 513-516.
25. Forest F, Berremila SA, Gyenes C, et al. Metastatic meningiomas: an unusual clinical and pathological diagnosis with highly variable outcome. *J Neurooncol.* 2014; 120(2): 411-421.
26. Frydrychowicz C, Holland H, Hantmann H, et al. Two cases of atypical meningioma with pulmonary metastases: a comparative cytogenetic analysis of chromosomes 1p and 22 and a review of the literature. *Neuropathology.* 2015; 35: 175-183.
27. Gezen F, Kahraman S, Canakci Z, Bedük A. Review of 36 cases of spinal cord meningioma. *Spine (Phila Pa 1976).* 2000; 25: 727-731.
28. Gordon A, Maloney A. A case of metastasizing meningioma. *J Neurol Neurosurg Psychiatry.* 1965; 28:159.
29. Guevara P, Escobar-Arriaga E, Saavedra-Perez D, et al. Angiogenesis and expression of estrogen and progesterone receptors as predictive factors for recurrence of meningioma. *J Neurooncol.* 2010; 98(3): 379-384.
30. Gunes M, Tugcu B, Gunaldi O et al. Seeding metastasis in the scalp after removal of the meningioma: a case report. *Turkish Neurosurg.* 2005; 15: 136-139.
31. Gyepes MT, D'Angio GJ. Extracranial metastases from central nervous system tumors in children and adolescents. *Radiology.* 1966; 87(1): 55-63.
32. Hu S, Zhang Y, Sun Y, et al. Lung metastases from intracranial bleeding meningioma: A case report. *Medicine (Baltimore).* 2018; 97(16):e0457.
33. Jain N, Mirakhor M, Flynn P, Choudhari K A. Cutaneous metastasis from 409 glioblastoma. *Br J Neurosurg.* 2005; 19(1): 65-68.
34. Kabus D, Sidhu GS, Wiczorek RL, Choi HS. Metastatic meningioma. Hemangiopericytoma or angioblastic meningioma? *Am J Surg Pathol.* 1993; 17: 1144-1150.

35. Kaminski JM, Movsas B, King E, et al. Metastatic meningioma to the lung with multiple pleural metastases. *Am J Clin Oncol*. 2001; 24(6): 579-582.
36. Kamiya K, Inagawa T, Nagasako R. Malignant intraventricular meningioma with spinal metastasis through the cerebrospinal fluid. *Surg Neurol*. 1989; 32: 213-218.
37. Kanzaki R, Higashiyama M, Fujiwara A, et al. Surgical resection of pulmonary metastases from meningioma: report of a case. *Surg Today*. 2011; 41: 995-998.
38. Karasick JL, Mullan SF. A survey of metastatic meningiomas. *J Neurosurg*. 1974; 40(2): 206-212.
39. Kessler RA, Garzon-Muvdi T, Yang W, et al. Metastatic Atypical and Anaplastic Meningioma: A Case Series and Review of the Literature. *World Neurosurg*. 2017; 101: 47-56.
40. Khan I, Amary F, Khoo MM. Meningioma metastasis to the bone - review of a rare finding with emphasis on distribution and imaging appearances. *Clinical Imaging*. 2015; 6: 1134-1139.
41. Kim H, Jung T-Y, Kim I-Y, Lee J-K. Two cases of primary osteolytic intrasosseous meningioma of the skull metastasizing to whole skull and the spine. *J Korean Neurosurg Soc*. 2012; 51: 151-154.
42. Koenig MA, Geocadin RG, Kulesza P, Olivi A, Brem H. Rhabdoid meningioma occurring in an unrelated resection cavity with leptomeningeal carcinomatosis. Case report. *J Neurosurg*. 2005; 102: 371-375.
43. Lanfranchi M, Nikpoor N. Detection of meningioma metastasis to liver and lung using somatostatin receptor scintigraphy. *Clin Nucl Med*. 2013; 38: 668-670.
44. Lee GC, Choi SW, Kim SH, Kwon HJ. Multiple extracranial metastases of atypical meningiomas. *J Korean Neurosurg Soc*. 2009; 45: 107-111.
45. Li Z, Cen Y, Gu M, Wei Y. Giant malignant meningioma invading the calvarial bone and scalp. *J Craniofac Surg*. 2012; 23(2): 599-602.
46. Louis D. WHO classification of tumours of the central nervous system. Lyon: IARC Press, 2007.
47. Lüdemann WO, Obler R, Tatagiba M, Samii M. Seeding of malignant meningioma along a surgical trajectory on the scalp. Case report and review of the literature. *J Neurosurg*. 2002; 97(3): 683-686.
48. McCarthy C, Hofer M, Vlychou M, et al. Metastatic meningioma presenting as a malignant soft tissue tumour. *Clin Sarcoma Res*. 2016; 6: 23.
49. Mentrikoski M, Johnson MD, Korones DN, Scott GA. Glioblastoma multiforme in skin: a report of 2 cases and review of the literature. *Am J Dermatopathol*. 2008; 30(4): 381-384.
50. Merell R. Brain tumors. In: Bope ET, Kellerman RD (eds) *Conn's Current Therapy*. Atlanta: Elsevier, 2017.
51. Moir JA, Haugk B, French JJ. Hepatic metastasis via a ventriculoperitoneal shunt from an intracranial meningioma: case report and review of the literature. *Case Rep Gastroenterol*. 2010; 4: 267-272.
52. Ozer E, Kalemci O, Acar UD, Canda S. Pin site metastasis of meningioma. *Br J Neurosurg*. 2007; 21(5): 524-527.
53. Pasquier D, Bijmolt S, Veninga T, et al. Atypical and malignant meningioma: outcome and prognostic factors in 119 irradiated patients. A multicenter, retrospective study of the Rare Cancer Network. *Int J Radiat Oncol Biol Phys*. 2008; 71(5): 1388-1393.
54. Peker S. Meningioma metastasis. In: Pamir MN, Black PM, Rahlbusch R (eds) *Meningiomas*. München: Elsevier, 2010.
55. Pinsker MO, Buhl R, Hugo H-H, Mehdorn HM. Metastatic meningioma WHO grade II of the cervical spine: case report and review of the literature. *Zentralbl Neurochir*. 2005; 66: 35-38.
56. Pramesh CS, Saklani AP, Pantvaidya GH, et al. Benign metastasizing meningioma. *Jpn J Clin Oncol*. 2003; 33(2): 86-88.
57. Raizer J, Sojka Sherman WJ. Meningiomas. In: Packer R, Schiff D (eds) *Neuro-oncology*. Oxford: John Wiley, 2012.
58. Saraf S, McCarthy BJ, Villano JL. Update on meningiomas. *Oncologist*. 2011; 16: 1604-1613.
59. Scognamiglio G, D'Antonio A, Rossi G, et al. CD90 expression in atypical meningiomas and meningioma metastasis. *Am J Clin Pathol*. 2014; 141: 841-849.
60. Shuangshoti S, Hongsaprabhas C, Netsky MG. Metastasizing meningioma. *Cancer*. 1970; 26(4): 832-841.
61. Singh SK, Ojha BK, Chandra A, et al. Scalp meningioma. *Indian J Dermatol*. 2008; 53(4): 196-198.
62. Spagnuolo E, Calvo A, Erman A, Tarigo A, Mañana G. Recurrent meningiomas with progressive aggressiveness and posterior extracranial extension. *Neurocirugia (Astur)*. 2003; 14(5): 409-416.
63. Subramanian A, Harris A, Piggott K, Shieff C, Bradford R. Metastasis to and from the central nervous system—the "relatively protected site". *Lancet Oncol*. 2002; 3(8): 498-507.
64. Surov A, Gottschling S, Bolz J, et al. Distant metastases in meningioma: an underestimated problem. *J Neurooncol*. 2013; 112: 323-327.
65. Tahir MZ, Shamim MS, Chishti KN. Recurrent atypical meningioma seeding to surgical scar. *Neurol India*. 2009; 57(2): 222-224.
66. Teague SD, Conces DJ Jr. Metastatic meningioma to the lungs. *J Thorac Imaging*. 2005; 20: 58-60.
67. Velnar T, Bunc G. Iatrogenic metastasis of a benign meningioma to the periosteum at the site of previous craniotomy: a case report. *Wien Klin Wochenschr*. 2008; 120: 766-769.
68. Wasserkrug R, Peyser E, Lichtig C. Extracranial bone metastases from intracranial meningiomas. *Surg Neurol*. 1979; 12(6): 480-484.
69. Waterson KW Jr, Shapiro L. Meningioma cutis: report of a case. *Int J Dermatol*. 1970; 9(2): 125-129.
70. Williamson BE, Stanton CA, Levine EA. Chest wall metastasis from recurrent meningioma. *Am Surg*. 2001; 67: 966-968.