



SHORT NOTE

Expanding the Distribution of the Remarkable Ant *Gnamptogenys vriesi* Brandão & Lattke (Formicidae: Ectatomminae): First Record From Brazil

RE VICENTE¹, LP PRADO², RCL SANTOS¹

1 - Universidade Federal de Mato Grosso (UFMT), Cuiabá-MT, Brazil

2 - Museu de Zoologia da Universidade de São Paulo (MZSP), São Paulo-SP, Brazil

Article History

Edited by

Rodrigo M. Feitosa, UFPR, Brazil

Received 24 September 2015

Initial acceptance 30 October 2015

Final acceptance 09 December 2015

Keywords

Amazon, Ants, Biodiversity, Faunistic inventory, Cryptic, Myrmecology.

Corresponding author

Ricardo Eduardo Vicente

Laboratório de Ecologia de

Comunidades do Instituto de Biologia

Universidade Federal de Mato Grosso

Av. Fernando Corrêa da Costa, nº 2367

Boa Esperança, 78060-900

Cuiabá-MT, Brazil

E-Mail: ricardomyrmex@gmail.com

ABSTRACT

Gnamptogenys vriesi Brandão and Lattke is a rarely collected ant, which was originally described based on a worker and a dealated queen collected in Morona-Santiago province, Ecuador. After the original description, only a few specimens were collected in the Ecuadorian territory. In this paper, we report the first record of *G. vriesi* in Brazil, which results in a significant extension of its distribution previously considered restricted to Ecuador. In addition, we provide images and a distribution map for the species.

The ant genus *Gnamptogenys* Roger, 1863 is distributed worldwide, with about 140 valid species (Bolton, 2014; Feitosa, 2015) found from southern United States to northern Argentina and from Southeast Asia to northern Australia (Lattke, 2003; 2004; Lattke et al., 2007). They have relatively small colonies and can nest on the ground, in fallen rotten logs, or even in trees (Lattke, 1994; Lattke et al., 2008), and associated with epiphytes (Gobin et al., 1998; Delabie et al., 2010). Most species of *Gnamptogenys* are generalist predators, whereas several species may have varying degrees of specialization (Lattke, 1995; 2003; Brandão et al., 2009; Brandão et al., 2015), preying on a large diversity of arthropods, such as other ants, beetles and millipedes (Brown, 1993; Brandão et al., 2009; Brandão et al., 2015).

In *Gnamptogenys*, the *minuta*-group comprises nine species (*G. bufonis* (Mann, 1926), *G. caelata* Kempf 1967, *G. falcifera* Kempf 1967, *G. fieldi* Lattke 1990, *G. minuta*

(Emery, 1896), *G. petiscapa* Lattke 1990, *G. simulans* (Emery, 1896), *G. striolata* (Borgmeier, 1957) and *G. vriesi* Brandão & Lattke 1990) distributed only in the Neotropics (Lattke, 1992). *Gnamptogenys vriesi* is a rarely collected species of the *minuta*-group (Lattke, 1992). Biological and behavioral traits of this species are still unknown, but the foveolae that covers its body surfaces are usually filled with earth, as found in *Basiceros* Schulz 1906 and *Stegomyrmex* Emery 1912, suggesting that the species presents cryptobiotic habits (Hölldobler & Wilson, 1986; Brandão & Lattke, 1990; Feitosa et al., 2008). Since the only known specimens of *Gnamptogenys vriesi* are a single worker (holotype) and one dealated queen (paratype) collected in Ecuador, the species has been considered endemic so far (Salazar et al., 2015). In this paper, we provide the first record of *G. vriesi* from Brazil.

A single worker of *G. vriesi* was collected by a pitfall trap in February, 2015, in a forest remnant in the municipality



of Juara, state of Mato Grosso, Brazil (Fig 1). The municipality has an area of 22,666 Km² with 34,87% of its original area deforested (*Instituto Nacional de Pesquisas Espaciais*, 2015). According to the Köppen classification, the climate is tropical humid (Am) with an annual average temperature above 24°C and precipitation ranging from 2,800 to 3,100 mm year (Alvares et al., 2013). There are two well-defined seasons: a rainy season, which occurs between November and April; and a dry season, that occurs between May and October (Vicente et al., 2014). The region is characterized by the transition of Cerrado-Amazon areas (Avila & Kawashita-Ribeiro, 2011), and the predominant vegetation consists of Submontane Open Ombrophilous Forest, Submontane Dense Ombrophilous Forest and ecotonal areas between Seasonal Forest and Dense Ombrophilous Forest (*Instituto Brasileiro de Geografia e Estatística* 2004).

The single specimen of *G. vriesi* collected is deposited at the ant collection of the *Museu de Zoologia da Universidade de São Paulo* (MZSP). The specimen was compared to the holotype worker, and its identity was confirmed by specialists in the group (see acknowledgments). Both specimens (the newly collected and the holotype) were photographed using a Leica Application Suite V3 system, at the MZSP Hymenoptera Lab. The images were processed in Adobe Photoshop 7.0.

All measurements (in mm) were obtained under 60x magnification with a Leica MZ95 stereomicroscope.

Abbreviations used throughout the text are: ML (mandible length): the straight line length of the mandible at full closure. HL (head length): the length of the head capsule excluding the mandibles, in full face view. WL (Weber's length): the diagonal length of mesosoma in profile. PL (petiole length): measured from anterior to posterior inflections of petiole node, in profile. GL (gaster length): the maximum length of gaster excluding sting in profile. TL (total length): the sum of ML, HL, WL, PL and GL.

Previous geographic distribution (holotype worker):

HL 2.2, ML 1, WL 3.2, PL 1.2, GL 4.7, TL 12.3; Fig 2): Ecuador, Morona-Santiago province, Los Tayos, 03°08'S, 78°14'W, 03.vii.1976, Tjitte de Vries col. (worker and dealated queen deposited in MZSP).

New record (worker):

HL 1.8, ML 1, WL 3.0, PL 0.9, GL 4.5, TL 11.2; Fig 2): Brazil, Mato Grosso State, Municipality of Juara, M1L1P7, 11°5'24.73"S 57°43'6.55"O, ii-2015, Santos R.C.L. col. (worker deposited in MZSP).

The specimen studied in this work presents the same diagnostic characters that define *G. vriesi* according to Brandão & Lattke (1990), such as WL>3,00mm; anterior border of clypeus concave; more than 20 ommatidia in the greatest diameter of eye; metanotal groove deep; and declivity of propodeum strongly impressed and concave. The worker collected in Brazil is morphologically similar, but the size is comparatively smaller (TL 11.2) than the holotype (TL 12.3); the coloration and sculpturing of the surface integument of the scape of antennae and legs are shiny and slightly darker; the anterior portion of the frontal lobe is weakly striate while in the holotype worker it is smooth; the region below the propodeal spiracles is smooth while in the holotype, this region is superficially foveolate.

This new record increases to 34, which is the number of *Gnamptogenys* species known from Brazil (Feitosa, 2015), and to 12, which is the number of species of the genus for Mato Grosso state (Economo & Guénard, 2015). As *G. vriesi* has only been collected to date in the Ecuadorian territory, this work extends the distribution of this species to more than 2,000 km. Therefore, the distribution of this species is much broader than was previously known. In fact, few inventories have been conducted between the extremes of the *G. vriesi* occurrence (e.g. Souza et al., 2015; Miranda et al., 2012).

Little is known about the distribution of most ant species due to the scarcity of inventories performed in most biomes (Brandão et al., 2008). In the Amazon, most of these inventories have been concentrated in the eastern and central regions (Vasconcelos et al., 2006; 2010; Oliveira et al. 2009; Bastos & Harada, 2011; Baccaro et al., 2012). Therefore, efforts should be made in the southern Amazon rainforest, as well as in Mato Grosso state, Brazil where the ant fauna is practically unknown (Vicente et al., 2011; 2012), and historically the region has faced strong pressure from the agricultural frontier and intensive anthropogenic disturbances.



Fig 1. Distribution map of *Gnamptogenys vriesi*.




Fig 2. Holotype worker of *Gnamptogenys vriesi* in: full-face view (A), dorsal view (B) and lateral view (C); and newly collected worker in: full-face view (D), dorsal view (E) and lateral view (F).

Acknowledgments

We are thankful to Dr John Lattke and Dr Carlos Roberto F. Brandão for his help with the confirmation of the ant species. To Vinicius S. Ferreira for reviewing a previous version of the manuscript and three anonymous reviewers for pertinent remarks that improved the manuscript. REV and RCLS thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (CAPES) for the doctoral fellowships in Ecologia e Conservação da Biodiversidade Postgraduate Program – UFMT. Is publication 51 in the NEBAM technical series.

References

- Alvares, C.A., Stape, J.L., Sentelhas, P.C., De-Moraes-Gonçalves, J.L. & Sparovek, G. (2014). Köppen's climate classification map for Brazil. Meteorologische Zeitschrift, 22: 711-728. doi: 10.1127/0941-2948/2013/0507
- Ávila, R.W. & Kawashita-Ribeiro, R.A. (2011). Herpetofauna of São João da Barra Hydroelectric Plant, state of Mato Grosso, Brazil. Check List, 7:750-755.
- Baccaro, F.B., Souza, J.L.P., Franklin, E., Landeiro, V.L. & Magnusson, W.E. (2012). Limited effects of dominant ants

- on assemblage species richness in three Amazon forests. *Ecological Entomology*, 37: 1-12. doi: 10.1111/j.1365-2311.2011.01326.x
- Bastos, A.H.S. & Harada, A.I. (2011). Leaf-litter as a factor in the structure of a Ponerine ants community (Hymenoptera: Formicidae: Ponerinae) in an eastern Amazonian rainforest, Brazil. *Revista Brasileira de Entomologia*, 55: 589-596. doi: /10.1590/S0085-56262011000400016
- Bolton, B. (2014). An online catalog of the ants of the world. <http://www.antcat.org/catalog> (accessed date: 20 September 2015).
- Brandão, C.R.F. & Lattke, J.E. (1990). Description of a new Ecuadorean *Gnamptogenys* species (Hymenoptera: Formicidae), with a discussion on the status of the Alfaria group. *Journal of the New York Entomological Society*, 98: 489-494.
- Brandão, C.R.F., Prado, L.P., Ulysséa, M.A., Probst, R.S., Alarcon, V. (2015). Dieta das Poneromorfas Neotropicais. In J.H.C. Delabie, , R.M. Feitosa, J.E. Serrão, C.S.F. Mariano & J.D. Majer (Eds.). As formigas poneromorfas do Brasil (pp. 145-161). Ilhéus: Editus.
- Brandão, C.R.F., Silva, R.R. & Delabie, J.C.H. (2009). Formigas. In: A.R. Panizzi & J.R.P. Parra (Eds.). Bioecologia e nutrição de insetos. Base para o manejo integrado de pragas. Embrapa Informação Tecnológica 323-370.
- Brandão, C.R.F.; Feitosa, R.M., Schmidt, F.A. & Solar, R.R.C. (2008). Rediscovery of the putatively extinct ant species *Simopelta mímina* (Brandão) (Hymenoptera, Formicidae), with a discussion on rarity and conservation status of ant species. *Revista Brasileira de Entomologia* 52: 480-483. doi: 10.1590/S0085-56262008000300026
- Brown, W.L. (1993). Two new species of *Gnamptogenys*, and an account of millipede predation by one of them. *Psyche* 99: 275-289.
- Delabie, J.H.C., Rocha, W.D., Feitosa, R.M., Devienne, P. & Fresneau, D. (2010). *Gnamptogenys concinna* (F. Smith, 1858): nouvelles données sur sa distribution et commentaires sur ce cas de gigantisme dans le genre *Gnamptogenys* (Hymenoptera, Formicidae, Ectatomminae). *Bulletin de la Société entomologique de France*, 115: 269-277.
- Econo, E. & Guénard, B. (2015). Antmaps.org. <http://antmaps.org/index.html>. (accessed date: 7 November, 2015).
- Feitosa, R.M. (2015). Lista das Formigas Poneromorfas do Brasil. In J.H.C. Delabie; R.M. Feitosa; J.E. Serrão; C.S. F. Mariano; J.D. Majer. (Eds.). As Formigas Poneromorfas do Brasil (pp. 95-102) Ilhéus: Editus.
- Feitosa, R.M., Brandão, C.R.F. & Diniz, J.L.M. (2008). Revisionary Studies on the Enigmatic Neotropical Ant Genus *Stegomyrmex* Emery, 1912 (Hymenoptera: Formicidae: Myrmicinae), With the Description of Two New Species. *Journal of Hymenoptera Research*, 17: 64-82.
- Gobin, B., Peetersand, C. & Billen, J. (1998). Colony reproduction and arboreal life in the ponerine ant *Gnamptogenys menadensis*. *Netherlands Journal of Entomology*, 48: 53-63.
- Hölldobler B. & Wilson, E.O. (1986). Soil-binding pilosity and camouflage in ants of the tribes Basicerotini and Stegomyrmecini (Hymenoptera: Formicidae). *Zoomorphology*, 106: 12-20.
- Instituto Brasileiro de Geografia e Estatística. (2004). Mapa da vegetação brasileira. 3ª edição. ftp://ftp.ibge.gov.br/Cartas_e_Mapas/Mapas_Murais/ (accessed date: 20 September 2015).
- Instituto de Pesquisas Espaciais. (2015) Projeto Prodes: Monitoramento da Floresta Amazônica Brasileira por Satélite. <http://www.obt.inpe.br/prodes/index.php> (accessed date: 20 September 2015).
- Lattke, J.E. (1992). Revision of the minuta-group of the genus *Gnamptogenys* (Hymenoptera, Formicidae). *Deutsche Entomologische Zeitschrift*, 39: 123-129.
- Lattke, J.E. (1994). Phylogenetic relationships and classification of ectatommine ants (Hymenoptera: Formicidae). *Entomologica Scandinavica*, 25: 105-119.
- Lattke, J.E. (1995). Revision of the ant genus *Gnamptogenys* in the New World (Hymenoptera: Formicidae). *Journal of Hymenoptera Research*, 4: 137-193.
- Lattke, J.E. (2003). Biogeographic analysis of the ant genus *Gnamptogenys* Roger in south-east Asia-Australasia(Hymenoptera: Formicidae: Ponerinae). *Journal of Natural History*, 37: 1879-1897.
- Lattke, J.E. (2004). A taxonomic revision and phylogenetic analysis of the ant genus *Gnamptogenys* Roger in Southeast Asia and Australasia (Hymenoptera: Formicidae: Ponerinae). *University of California Publications in Entomology*, 122: 1-266.
- Lattke, J.E., Fernández, F. & Palacio, E.E. (2007). Identification of the species of *Gnamptogenys* Roger in the Americas. In R.R. Snelling, B.L. Fisher & P.S. Ward (eds). *Advances In Ant Systematics* (Hymenoptera: Formicidae): Homage To E.O. Wilson – 50 Years Of Contributions. *Memoirs of the American Entomological Institute*, 80.
- Lattke, J.E., Fernández, F., Arias-Penna, T.M., Palacio, E.E., Mackay, W. & Mackay, E. (2008). Género *Gnamptogenys* Roger. In E. Jiménez, F. Fernández, T.M. Arias-Penna & F.H. Lozano-Zambrano. *Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia* (pp. 65-105). Instituto Humboldt, Colombia.
- Miranda, P.N., Oliveira, M.A., Baccaro, F.B., Morato, E.F. & Delabie, J.H.C. (2012) Check list of ground-dwelling ants (Hymenoptera: Formicidae) of the eastern Acre, Amazon, Brazil. *CheckList*, 8 (40): 722.
- Oliveira, P.Y., Souza, J.L.P., Baccaro, F.B. & Franklin, E. (2009). Ant species distribution along a topographic gradient

- in a “terra-firme” forest reserve in Central Amazonia. *Pesquisa Agropecuária Brasileira*, 44: 852-860.
- Salazar, F., Reyes-Bueno, F., Sanmartin, D. & Donoso, D.A. (2015). Mapping continental Ecuadorian ant species. *Sociobiology* 62: 132-162. doi: 10.13102/sociobiology.v62i2.132-162
- Souza, J.L.P., Baccaro, F.B., Landeiro, V.L., Franklin, E., Magnusson, W.E., Pequeno, P.A.C.L. & Fernandes, I.O. (2015). Taxonomic sufficiency and indicator taxa reduce sampling costs and increase monitoring effectiveness for ants. *Diversity and Distributions*, 22: 111-122. doi: 10.1111/ddi.12371
- Vasconcelos, H.L., Vilhena, J.M.S., Facure, C.G. & Albernaz, A.L.K.M. (2010). Patterns of ant species diversity and turnover across 2000km of Amazonian floodplain forest. *Journal of Biogeography*, 37: 432-440. doi: 10.1111/j.1365-2699.2009.02230.x
- Vasconcelos, H.L., Vilhena, J.M.S., Magnusson, W.E. & Albernaz, A.L.K.M. (2006). Long-term effects of forest fragmentation on Amazonian ant communities. *Journal of Biogeography*, 33: 1348-1356. doi: 10.1111/j.1365-2699.2006.01516.x
- Vicente, R.E., Dambroz, J. & Barreto, M. (2011). New distribution record of *Daceton boltoni* Azorsa and Sosa-Calvo, 2008 (Insecta: Hymenoptera) ant in the Brazilian Amazon. *Check List*, 7: 878-879. doi: 10.15560/7.6.878
- Vicente, R.E., Dátilo, W. & Izzo, T.J. (2012). New record of a very specialized interaction: *Myrcidris epicharis* Ward 1990 (Pseudomyrmecinae) and its myrmecophyte host *Myrcia madida* McVaugh (Myrtaceae) in Brazilian Meridional Amazon. *Acta Amazonica*, 42: 567-570. doi: 10.1590/S0044-59672012000400016
- Vicente, R.E., Dátilo, W. & Izzo, T.J. (2014). Differential recruitment of *Camponotus femoratus* (Fabricius) ants in response to ant garden herbivory. *Neotropical Entomology*, 43: 519-525. doi: 10.1007/s13744-014-0245-6

